

🌸 日本数学会

2020年度年会

# 英 文 サ マ リ 集

2020年3月

於 日本大学



# 2020 日本数学会 年会プログラム

期 日 2020年3月16日(月)～3月19日(木)

会 場 日本大学駿河台キャンパス  
〒101-8308 東京都千代田区神田駿河台1-8-14

連絡先 日本大学理工学部数学科  
〒101-8308 東京都千代田区神田駿河台1-8-14  
E-mail nichidai20mar@mathsoc.jp

(会期中) Tel 090-1791-3483

一般社団法人 日本数学会  
Tel 03-3835-3483

	第I会場 タワー・スコラ S101	第II会場 タワー・スコラ S204	第III会場 タワー・スコラ S301	第IV会場 タワー・スコラ S302	第V会場 タワー・スコラ S401	第VI会場 タワー・スコラ S505	第VII会場 理工1号館 141	第VIII会場 理工1号館 144	第IX会場 理工1号館 142
16日 (月)	代 数 学 10:00～12:00 15:30～16:50	幾 何 学 9:40～11:40 14:20～15:45	函数方程式論 9:00～12:00 14:15～16:15	トポロジー 10:00～12:00	応用数学 10:00～11:40 14:20～16:05	統計数学 9:20～11:55 14:15～15:00	函数解析学 10:00～11:45	函 数 論 9:30～11:50 14:15～15:20	数学基礎論 および歴史 9:30～12:00 14:15～15:00
	企画特別講演 13:00～14:00								
	特別講演 14:15～15:15	特別講演 16:00～17:00	特別講演 16:30～17:30	特別講演 14:15～15:15 15:30～16:30	特別講演 16:20～17:20	特別講演 15:15～16:15 16:30～17:30	特別講演 14:15～15:15 15:30～16:30	特別講演 15:35～16:35 16:50～17:50	特別講演 15:15～16:15
17日 (火)	代 数 学 10:00～11:45 13:00～14:15	幾 何 学 9:40～11:50 特別講演 13:15～14:15	函数方程式論 9:00～12:00 特別講演 13:15～14:15	トポロジー 10:00～12:00 特別講演 13:15～14:15	応用数学 9:50～11:15	統計数学 9:10～11:25	函数解析学 10:30～11:45 特別講演 13:15～14:15	函 数 論 9:15～11:45 特別講演 13:15～14:15	数学基礎論 および歴史 9:00～10:30 特別講演 10:45～11:45
	日本数学会賞授賞式(理工学部1号館 CSTホール) …………… (15:00～15:20)								
	総 合 講 演 ( " ) 日本数学会賞春季賞受賞者 …………… (15:30～16:30)								
	向 井 茂(京大数理研) …………… (16:45～17:45)								
懇 親 会(理工学部1号館 カフェテリア) …………… (18:00～20:00)									
18日 (水)	代 数 学 9:30～12:00	幾 何 学 9:40～11:35 14:20～15:40	函数方程式論 9:00～12:00 14:15～16:15	トポロジー 10:00～12:00	応用数学 9:30～12:00 14:30～15:50	統計数学 9:00～12:00 14:15～15:05	函数解析学 9:45～12:00 14:15～15:10	実函数論 9:00～11:50 14:15～15:55	無限可積分系 14:15～16:15
	企画特別講演 13:00～14:00								
	特別講演 14:40～15:40 15:50～16:50	特別講演 16:00～17:00	特別講演 16:30～17:30		特別講演 16:00～17:00	特別講演 15:20～16:20 16:35～17:35	特別講演 15:20～16:20	特別講演 16:15～17:15	特別講演 16:30～17:30
19日 (木)	代 数 学 9:20～12:00 15:25～16:50		函数方程式論 9:00～12:00 14:15～16:15		応用数学 9:15～10:50			実函数論 9:00～12:00 14:15～15:55	無限可積分系 9:45～11:30 14:15～16:00
	企画特別講演 13:00～14:00								
	特別講演 14:15～15:15		特別講演 16:30～17:30		特別講演 11:00～12:00			特別講演 16:15～17:15	特別講演 16:15～17:15



## 目次

総合講演	1
企画特別講演	2
数学基礎論および歴史	4
3月16日(月)	4
3月17日(火)	6
代数学	9
3月16日(月)	9
3月17日(火)	12
3月18日(水)	16
3月19日(木)	20
幾何学	26
3月16日(月)	26
3月17日(火)	29
3月18日(水)	31
函数論	35
3月16日(月)	35
3月17日(火)	38
函数方程式論	41
3月16日(月)	41
3月17日(火)	46
3月18日(水)	50
3月19日(木)	56
実函数論	62
3月18日(水)	62
3月19日(木)	65
函数解析学	70
3月16日(月)	70
3月17日(火)	72
3月18日(水)	73
統計数学	77
3月16日(月)	77
3月17日(火)	80
3月18日(水)	82
応用数学	87

3月16日(月) . . . . .	87
3月17日(火) . . . . .	90
3月18日(水) . . . . .	92
3月19日(木) . . . . .	96
トポロジー . . . . .	99
3月16日(月) . . . . .	99
3月17日(火) . . . . .	101
3月18日(水) . . . . .	103
無限可積分系 . . . . .	106
3月18日(水) . . . . .	106
3月19日(木) . . . . .	107

## 総 合 講 演

3月17日(火) 総合講演会場

2020年度日本数学会賞春季賞

日本数学会賞春季賞受賞者 ..... (15:30~16:30)

Spring Prize Winner

向 井 茂 (京大数理研) 代数多様体とその対称性 —K3曲面とその仲間達を中心に—

..... (16:45~17:45)

Shigeru Mukai (Kyoto Univ.) Algebraic varieties and their symmetry with emphasis on K3 surfaces and their companions

概要 There are many phenomena where algebraic geometry and group theory interplay. Algebraic varieties often have apparent or hidden symmetry such as the Schläfli configuration of 27 lines on a cubic surface or ADE-type degeneration of elliptic curves. In 80s I classified finite groups acting symplectically on a K3 surface, in terms of the Mathieu group  $M_{24}$ , one of the 26 sporadic finite simple groups. In 90s Kondo determined the (infinite) automorphism group of generic Jacobian Kummer surfaces using the Higman–Sims(or Menser) graph. Besides these topics, I will also discuss recent results on the decomposition groups of certain plane curves and a conjecture on the virtual cohomological dimension of the automorphism groups if time permits.

## 企画特別講演

3月16日(月)

### 第V会場

堀山 貴史 (北大 情報) 多面体の展開と折り ..... (13:00~14:00)

Takashi Horiyama (Hokkaido Univ.) Folding and unfolding of polyhedra

概要 A development of a polyhedron is a simple polygon obtained by cutting the surface of the polyhedron and unfolding it into a plane. In this talk, we focus on the problems related to the following two big issues:

- (1) Does every convex polyhedron have a nonoverlapping development?
- (2) Are there any common developments for two (or more) polyhedra?

### 第IX会場

藤田 博司 (愛媛 大理) 超限順序数と連続体問題 ..... (13:00~14:00)

Hiroshi Fujita (Ehime Univ.) Transfinite ordinals and the continuum problem

概要 We will explain how Georg Cantor introduced transfinite ordinals into mathematics. From this point of view, we also explain Cantor's continuum problem. The problem can be understood as the question asking how the continuum is well-ordered. We review some known results about definable well-ordering of the continuum.

3月18日(水)

### 第I会場

富安 亮子 (九大 I M I) 数理結晶学における格子の問題 ..... (13:00~14:00)

Ryoko Oishi-Tomiyasu (Kyushu Univ.) Lattice problems in mathematical crystallography

概要 The concept of lattices was originally introduced and studied by August Bravais in botany and crystallography. Nowadays, lattice problems are also studied for lattice-based crystallography, an important candidate for post-quantum cryptography. In such application, to what extent can abstract results in pure mathematics be useful? Starting from the pioneer works of Bravais, I'd explain about several problems in modern crystallography and examples for actual use of number theory I have obtained so far.

### 第II会場

栗林 勝彦 (信州 大理) 導来ストリングトポロジー — 分類空間の2次元開閉位相的場の理論へ — ..... (13:00~14:00)

Katsuhiko Kuribayashi (Shinshu Univ.) Derived string topology — Toward a two dimensional open-closed topological quantum field theory for classifying spaces —

概要 String topology introduced by Chas and Sullivan gives fruitful structures to the loop homology of orientable closed manifolds, more general Gorenstein spaces whose class contains Poincaré duality space, classifying spaces and Borel constructions. In particular, a result due to Chataur and Menichi asserts that the loop homology of the classifying space of a Lie group is endowed with the structure of a two dimensional topological quantum field theory (TQFT). Guldberg has proved that such a structure is generalized to that of a labeled open-closed TQFT. However, there are few calculations of labeled cobordism operations in the theory. In this talk, after recalling the original string topology and derived one due to Félix and Thomas, we survey computations of string operations for the classifying space of a Lie group. Moreover, we consider the non-triviality of the whistle cobordism operation with a label in the set of maximal rank subgroups of the given Lie group. It turns out that the open TQFT and closed one are not separated in general. A part of this talk is based on joint work with Luc Menichi and Takahito Naito.



3月19日(木)

**第III会場**

野海正俊(神戸大理) 超幾何関数の拡がり ..... (13:00~14:00)

Masatoshi Noumi (Kobe Univ.) Expanse of hypergeometric functions

概要 I will give an overview of various topics related to hypergeometric functions, and discuss the present status and possible future directions in the study of special functions.

**第V会場**

特別招待講演(日本応用数理学会)

内山成憲(首都大東京理) ポスト量子暗号の最近の話題について ..... (13:00~14:00)

Shigenori Uchiyama Recent topics in post-quantum cryptography  
(Tokyo Metro. Univ.)

概要 In recent years, research on public key cryptosystems, which are expected to be secure against attacks using quantum computers called post-quantum cryptography (PQC), has been actively conducted. In this talk, I am going to talk about an overview of the recent research in post-quantum cryptography.

**第VIII会場**

小林和夫(早大教育) 確率偏微分方程式の kinetic 方程式論的考察 ..... (13:00~14:00)

Kazuo Kobayasi (Waseda Univ.) A kinetic approach to stochastic partial differential equations

概要 We study some stochastic partial differential equations, in particular stochastic scalar conservation laws on a bounded domain. We see that these equations appear in the hydrodynamical limit of a stochastic kinetic equation where the interactions in velocity are modelled by a nonlinear operator (Bhatnager-Gross-Krook) or a linear operator (Linear Boltzmann or Fokker-Planck). Moreover we derive a stochastic kinetic equation from the stochastic differential equation called the Langevin equation: Consider the motion of a particle with the position  $X(t)$  and the velocity  $V(t)$ . We assume that  $(X(t), V(t))$  satisfies the Langevin equation which is interpreted as Newton's second law of motion applied to the position  $X(t)$ . Then the stochastic kinetic equation is satisfied with the density function  $f(t, x, v)$ , with respect to the Lebesgue measure  $dx dv$ , of the conditional law of  $(X(t), V(t))$ . Finally we treat the stochastic scalar conservation laws obtained above. We discuss them on a bounded domain. First, we consider a more general equation, that is a quasilinear degenerate parabolic stochastic partial differential equation with a multiplicative noise on the  $N$ -dimensional torus. We propose to construct a sequence of approximations by applying a time splitting method and prove that this converges to a kinetic solution. Secondarily, we discuss a stochastic scalar conservation law with non-homogeneous Dirichlet boundary condition. By introduction a notion of renormalized kinetic solutions in which the kinetic defect measures on the boundary of a domain are truncated, we establish a result of well-posedness of renormalized kinetic solutions.

# 数 学 基 礎 論 お よ び 歴 史

3月16日(月) 第IX会場

9:30~12:00

- 1 依岡輝幸(静岡大理) YPFA は MRP を導く ..... 15  
 宮元忠敏(南山大経営)  
 Teruyuki Yorioka (Shizuoka Univ.) YPFA implies MRP  
 Tadatoshi Miyamoto (Nanzan Univ.)

概要 It is proved that YPFA implies MRP.

- 2 池上大祐(芝浦工大SIT総合研) Generic absoluteness in ZF ..... 15  
 Daisuke Ikegami Generic absoluteness in ZF  
 (Shibaura Inst. of Tech.)

概要 By the discovery of forcing by Cohen, in ZFC, one can show that there is a partial order which forces the Continuum Hypothesis (CH) while there is a partial order which forces the negation of CH. What if we do not assume the axiom of choice and work in ZF? Can we still find such a statement as CH? Note that the formulation of CH and the arguments to force CH heavily depend on the axiom of choice.

In this talk, we show that in ZF, one can find a statement  $\phi$  such that there is a partial order which forces  $\phi$  while there is a partial order which forces the negation of  $\phi$ .

This is joint work with Philipp Schlicht and W. Hugh Woodin.

- 3 薄葉季路(早大理工) On generically extendible cardinals ..... 15  
 Toshimichi Usuba (Waseda Univ.) On generically extendible cardinals

概要 We study generically extendible cardinals. We show that the consistency strength of the generic extendibility of  $\omega_1$  is weak, but  $\omega_3$  is not.

- 4 D. A. Mejia (静岡大理) Lebesgue measure zero modulo ideals ..... 15  
 Diego A. Mejía (Shizuoka Univ.) Lebesgue measure zero modulo ideals

概要 Lebesgue measure zero subsets of the real line can be characterized, in a combinatorial way, through the ideal  $\text{Fin}$  of finite subsets of the natural numbers. But, what happens if we consider this combinatorial statement through an arbitrary ideal on the natural numbers, instead of the ideal  $\text{Fin}$ ? We present a dichotomy deciding whether this new property characterizes Lebesgue measure zero or not, and show the connections of this property with Baire category and with the sigma ideal generated by closed measure zero sets. This is a joint work with Viera Sottova.

- 5 藤田憲悦(群馬大理工) George Boolos' "The Hardest Logic Puzzle Ever" revisited ..... 15  
 Kenetsu Fujita (Gunma Univ.) George Boolos' "The Hardest Logic Puzzle Ever" revisited

概要 Following R. Smullyan, G. Boolos posed the puzzle "The Hardest Logic Puzzle Ever" (1996), and provided an answer in the form of "iff". After that, Roberts (2001) and Rabern–Rabern (2008) showed a simple solution to the puzzle by using the embedded question lemma. We introduce a formalization of such Knights and Knaves puzzles in term of propositional logic, and observe that the two answers by Boolos and Roberts–Rabern are essentially logically equivalent. This elegant method can be applied to an analysis of the puzzle in the fantasy film Labyrinth as well.

- 6 関 隆 宏 (新潟大経営戦略本部) 対偶を持つ対合的部分構造論理に対する Gentzen 流の形式化 ..... 15  
 Takahiro Seki (Niigata Univ.) A Gentzen-style formulation for involutive substructural logics with contra-  
 position

概要 Involutivity, called double-negation axiom in classical logic, is one of the important additional axioms to intuitionistic substructural logics. In this talk, we consider a Gentzen-style formulation for some involutive non-associative substructural logics and show the cut elimination theorem. The logics in our formulation include contraposition axiom, regarded as a restricted associativity.

- 7 田 中 義 人 (九州産大経済) A representation of modal algebras preserving countably many infinitary  
 meets and joins ..... 15  
 Yoshihito Tanaka A representation of modal algebras preserving countably many infinitary  
 (Kyushu Sangyo Univ.) meets and joins

概要 By incorporating the Jónsson–Tarski Theorem with the Rasiowa–Sikorski Lemma, Tanaka–Ono gives an embedding which preserves countably many infinitary meets and joins, from every modal algebra which satisfies certain conditions including a weaker form of complete additivity into the complex algebra of a subframe of the prime filter Kripke frame. In this talk, we give another representation for modal algebras without assuming complete additivity. We show that for every modal algebra, there exists an embedding preserving countably many infinitary meets and joins into the complex algebra of a subframe of the prime filter neighborhood frame.

- 8 大 川 裕 矢 (千葉大融合理工) 部分保存的な文に対する Bennet の結果の一般化 ..... 15  
 倉 橋 太 志 (木更津工高専)  
 Yuya Okawa (Chiba Univ.) Generalizations of Bennet’s result on partially conservative sentences  
 Taishi Kurahashi  
 (Nat. Inst. of Tech., Kisarazu Coll.)

概要 A sentence  $\varphi$  is said to be  $\Gamma$ -conservative over  $T$  if for every  $\Gamma$  sentence  $\theta$ , if  $T + \varphi \vdash \theta$ , then  $T \vdash \theta$ . For  $\Gamma = \Sigma_n$  (resp.  $\Pi_n$ ), let  $\Gamma^d = \Pi_n$  (resp.  $\Sigma_n$ ). In 1979, Guaspari asked that for any reasonable sequence  $\{T_i\}_{i \in \omega}$  of theories, whether there is a  $\Gamma^d$  sentence which is simultaneously independent and  $\Gamma$ -conservative over each  $T_i$ . For two theories, this problem was investigated by Bennet. He completely characterized the existence of simultaneously independent and  $\Sigma_n$ -conservative  $\Pi_n$  sentences over two theories.

We generalized Bennet’s results to the case of theories more than two.

#### 14:15~15:00

- 9 増 田 茂 (流体数理古典理論研) Study of the Eulerian Integrals by Legendre ..... 15  
 Shigeru Masuda Study of the Eulerian Integrals by Legendre  
 (Res. Workshop of Classical Fluid Dynamics)

概要 Legendre issues *Traite des fonctions elliptiques et des integrales euleriennes*, in 1825. In this books he discusses Eulerian Integral with two sorts of integrals, in relation to Euler’s integrals, including his elliptic functions. Legendre complains Euler’s integral, saying “they have never been occupied to make the calculation easy, nor to fix the degree of precision of which it is susceptible,” (§169) and proposes that some functions are explained with the arcs of circle and of the logarithms. (§60, etc.)

- 10 増田 茂 (流体数理古典理論研) The complete functions by Legendre ..... 15  
 Shigeru Masuda The complete functions by Legendre  
 (Res. Workshop of Classical Fluid Dynamics)

概要 In 1825, Legendre defines as the most important function among his elliptic functions, the complete function (=c.f.)  $F^1, E^1, \Pi^1$ . Afterward in modern times, these functions are newly expressed with  $K, E$  and  $\Pi$  such as in the study of a quantum mechanics, where it shows as  $F^1 = K$ . We call  $E$  and  $\Pi$  c.f.s of newly defined versions, then  $E = E^1, \Pi = \Pi^1$ . Legendre uses the c.f. in variously applied arena, such as Geometry, Mechanics, Construction of the Tables, Eulerian Integrals, etc. We discuss the c.f. as the original by Legendre, his objects, effects, and so on.

- 11 真島 秀行 (お茶の水女大\*) 2022年, 関孝和314年祭等に向けて ..... 15  
 Hideyuki Majima (Ochanomizu Univ.\*) Towards the year 2022, the 314th memorial year of SEKI Takakazu

概要 The year 2022 has various meaning for mathematics, mathematical history and mathematical education: the 314th year after his death of SEKI Takakazu (?-1708), the 400th year after the publication of Division-Instruction (Wariznsyo), the 300th anniversary of TAKEBE Katahiro's Tetsujjutsu-Sankei, and so on. We describe research problems towards the year.

### 15:15~16:15 特別講演

- 高瀬 正仁 西欧近代の数学におけるさまざまな泉の造型  
 Masahito Takase Shaping the fountains in Modern Western Mathematics

概要 There are several impressive fountains in the history of the mathematics on which Modern Western Mathematics is founded as follows: Descartes's method of normals, Fermat's method of tangents, Fermat's number theory, Leibniz's analysis of the infinity, Johann Bernoulli's beautiful formula  $\frac{\log \sqrt{-1}}{\sqrt{-1}} = \frac{\pi}{2}$ , Euler's observation on dividing 0 by 0, Euer's discovery of the concept of function, Gauss' idea of transferring number theory from rational integers to Gaussian integers, The concept of an Abelian equation discovered by Abel, Kronecker's "Jugendtraum", Kummer's idea of ideal prime factors, Jacobi's inverse problem of Abelian integrals invented by Jacobi, Riemann's idea of Riemann surfaces, Hartogs' inverse problem invented by K. Oka.

I will explain the significances of returning to the fountains and make a critical observation on contemporary mathematics.

3月17日(火) 第IX会場

### 9:00~10:30

- 12 只木孝太郎 (中部大工) アルゴリズム的ランダムネスによる量子情報理論の精密化 III ..... 15  
 Kohtaro Tadaki (Chubu Univ.) A refinement of quantum information theory by algorithmic randomness  
 III

概要 The notion of probability plays a crucial role in quantum mechanics. It appears as the Born rule. In modern mathematics which describes quantum mechanics, however, probability theory means nothing other than measure theory, and therefore any operational characterization of the notion of probability is still missing in quantum mechanics. In our former works, based on the toolkit of algorithmic randomness, we presented an operational refinement of the Born rule, called the principle of typicality, for specifying the property of the results of quantum measurements in an operational way. In this talk, we refine and reformulate the theory of quantum error-correction based on the principle of typicality, in order to demonstrate how properly our framework works in practical problems in quantum mechanics.

- 13 宮部 賢志 (明大理工) 帰納的推論における収束速度 ..... 15  
 滝坂 透 (国立情報学研)  
 Kenshi Miyabe (Meiji Univ.) The speed of convergence of induction  
 Toru Takisaka  
 (Nat. Inst. of Information)

概要 Consider trying to predict the next bit from a given finite binary string when the sequence is sampled from a computable probability measure on the Cantor space. There exists the best betting strategy among a class of effective ones up to a multiplicative constant, the induced prediction from which is called algorithmic probability or universal induction by Solomonoff. The prediction converges to the true induced measure for sufficiently random sequences. However, the prediction is not computable. We introduce a framework to study the properties of computable predictions. In this framework, we can prove that all sufficiently general computable predictions also converge to the true induced measure. We also discuss the speed of convergence.

- 14 荒武 永史 (京大数理研) Classifying toposes for existentially closed models and finite-generic models ..... 15  
 Hisashi Aratake (Kyoto Univ.) Classifying toposes for existentially closed models and finite-generic models

概要 In the context of first-order categorical logic, Blass and Scedrov constructed classifying toposes for existentially closed models and finite-generic models. We make some categorical analysis of these toposes and deduce model-theoretic results.

- 15 田中 一之 (東北大理工) ブール木に対する固有独立分布について ..... 15  
 Kazuyuki Tanaka (Tohoku Univ.) On eigen-distributions for Boolean trees in the ID case

概要 Our primary objective is to characterize the “eigen-distribution”  $d$  in  $ID(r)$  for a (weighted) Boolean tree (with a fixed probability  $0 < r < 1$  for the root), which achieves the distributional complexity. In this talk, we introduce two new concepts on independent distributions: “proportional” ID (PID for short) and “decent” ID. For a fixed root probability  $0 < r < 1$ , a  $PID(r)$  on a weighted tree is uniquely constructed by a technique “Super-RAT”. As a main theorem, we show that if the eigen-distribution  $d$  in  $ID(r)$  is decent then it is PID, from which we can deduce our previous theorem that for a balanced AND-OR tree (with no weights considered), the eigen-distribution  $d$  in  $ID(r)$  is IID.

- 16 横山 啓太 (北陸先端大) 算術における非構成的論理原理のクリプキモデルを用いた分離 ..... 15  
 藤原 誠 (明大理工)  
 石原 哉 (北陸先端大)  
 根元多佳子 (北陸先端大)  
 鈴木 信行 (静岡大理)  
 Keita Yokoyama (JAIST) Kripke models and separations of logical principles  
 Makoto Fujiwara (Meiji Univ.)  
 Hajime Ishihara (JAIST)  
 Takako Nemoto (JAIST)  
 Nobu-Yuki Suzuki (Shizuoka Univ.)

概要 In this talk, I will explain some ideas of separating non-constructive logical principles (omniscience principles) over Heyting arithmetic by using Kripke models.

- 17 新井 敏康 (東大数理) Some contributions to proof theory ..... 15  
 Toshiyasu Arai (Univ. of Tokyo) Some contributions to proof theory

概要 I will report some recent results in proof theory.

**10:45~11:45 特別講演**

鈴木 信行 (静岡 大理) 中間述語論理における選言特性と存在特性

Nobu-Yuki Suzuki (Shizuoka Univ.) Disjunction and existence properties in intermediate predicate logics

**概要** We discuss relationships between the disjunction property (DP) and existence property (EP) in intermediate predicate logics. These properties are regarded as characteristics of constructivity of intuitionistic logic. Since existence-quantifier could be written as powerful (possibly infinitary) disjunction, we expected that there must be some relationship between them. They are, however, independent in intermediate predicate logics. This result contrasts with those e.g., in intuitionistic and modal arithmetic, and also gives a negative answer to Ono's problem P52 posed in 1987. The key is the quantifier-annihilation axiom Z. This axiom is abnormal in intermediate logics; but it works when we discuss DP and EP in superintuitionistic predicate logics. We introduce a condition, Z-normality, that excludes the Z and is a natural condition for reasonable logics. Then, we can show that EP implies DP for Z-normal logics. This result suggests a reason why Ono's problem P52 has remained open. We also report some recent results related to DP and EP.

**11:45~12:00 数学基礎論および歴史分科会総会****12:00~12:15 歴史部門懇談会**

## 代 数 学

3月16日(月) 第I会場

## 10:00~12:00

- 1 中空大幸 (神戸学院大共通教育センター) ゴーレイ符号の extended doubling とムーンシャイン符号 ..... 10  
Hiroyuki Nakasora (Kobe Gakuin Univ.) The extended doubling of the Golay code and the Moonshine code

概要 In this talk, we present an example of codes all of whose weight classes support 1-designs, with duals whose classes include two that support 2-designs. We can find this example in the triply even binary codes of length 48.

- 2 B. Mühlherr (Univ. Giessen) Locally finite continuations and Coxeter groups of infinite ranks ..... 10  
縫田光司 (東大情報理工) Bernhard Mühlherr (Univ. Giessen) Locally finite continuations and Coxeter groups of infinite ranks  
Koji Nuida (Univ. of Tokyo)

概要 In some previous works on the isomorphism problem in Coxeter groups, certain special kind of subgroups called finite continuations have played a central role. However, the definition of finite continuations assumes finiteness of the rank (cardinality of generating set) of the underlying Coxeter group and is in general not available in the infinite rank cases. In our present work, we propose a generalization of finite continuations to the infinite rank cases (which itself is also defined for arbitrary groups), and study how the previous results in the finite rank cases do or do not extend to the infinite rank cases.

- 3 千吉良直紀 (熊本大理) 群上のある方程式の解集合と群の構造 ..... 10  
Naoki Chigira (Kumamoto Univ.) Solutions of a certain equation on a group and group structure

概要 We consider the set of solutions of two variable equation  $y^{-1}[x, y]^{[x^2, y]} = a$  on a finite group  $G$  and  $a \in G$ . We discuss some properties of a group  $G$  using the number of solutions of the equation.

- 4 小田文仁 (近畿大理工) 斜バーンサイド環とマッキー 2 関手 ..... 10  
竹ヶ原裕元 (室蘭工大) Fumihito Oda (Kindai Univ.) Crossed Burnside rings and Mackey 2-functors  
Yugen Takegahara (Muroran Inst. of Tech.)

概要 Balmer and Dell'Ambrogio introduced Mackey 2-functors. They showed that the 2-endomorphism ring of the identity of a finite group  $G$  in certain 2-category of Mackey 2-motives is isomorphic to the so called crossed Burnside ring of  $G$ . We show that a decomposition of  $\mathcal{M}(G)$  for a Mackey 2-functor  $\mathcal{M}$  into additive categories corresponds to the set of conjugacy classes of perfect subgroups of  $G$ .

- 5 飛田明彦 (埼玉大教育) 有限群の既約指標の次数の積と共役類の長さの積の関係について ..... 10  
Akihiko Hida (Saitama Univ.) On the relation of the product of character degrees and the product of conjugacy class lengths of a finite group

概要 Let  $G$  be a finite group. K. Harada conjectured that the product of degrees of all irreducible character of  $G$  divides the product of lengths of all conjugacy classes of  $G$ . We verify this conjecture for symmetric groups, alternating groups and wreath products.

- 6 飛田明彦 (埼玉大教育) Lower defect groups and vertices of simple modules ..... 10  
 Akihiko Hida (Saitama Univ.) Lower defect groups and vertices of simple modules

概要 Let  $kG$  be the group algebra of a finite group  $G$  over an algebraically closed field  $k$  of positive characteristic  $p$ . Let  $b$  be a block idempotent of  $kG$ . We compare  $p$ -regular lower defect groups of  $b$  with vertices of simple  $kGb$ -modules.

- 7 越谷重夫 <sup>b</sup> 2次特殊アフィン変換群  $\text{Qd}(p)$  に対するスコット加群のブラウアー直既約性 ..... 10  
 (千葉大先進科学センター・千葉大\*)

I. Tuvay

(Mimar Sinan Fine Arts Univ.)

Shigeo Koshitani

(Chiba Univ./Chiba Univ.\*)

Brauer indecomposability of Scott modules for the quadratic group

$\text{Qd}(p)$

Ipek Tuvay

(Mimar Sinan Fine Arts Univ.)

概要 Let  $k$  be an algebraically closed field of prime characteristic  $p$  and  $P$  a finite  $p$ -group. We compute the Scott  $kG$ -module with vertex  $P$  when  $\mathcal{F}$  is a fusion system on  $P$  and  $G$  is Park's group for  $\mathcal{F}$ . Further when  $\mathcal{F}$  is a fusion system of the 2-dimensional special affine groups of transformations  $\text{ASL}(2, p) = \text{Qd}(p) = (\mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z}) : \text{SL}(2, p)$  on a Sylow  $p$ -subgroup  $P$  of  $\text{Qd}(p)$  and  $G$  is Park's group for  $\mathcal{F}$ , we prove that the Scott  $kG$ -module with vertex  $P$  is Brauer indecomposable.

- 8 越谷重夫 <sup>b</sup> 二面体群を Sylow 2-部分群に持つ有限群の主ブロックの splendid Morita 同値について ..... 10  
 (千葉大先進科学センター・千葉大\*)

C. Lassueur (TU Kaiserslautern)

Shigeo Koshitani

(Chiba Univ./Chiba Univ.\*)

Splendid Morita equivalences for principal blocks with dihedral defect groups

Caroline Lassueur (TU Kaiserslautern)

概要 Given a dihedral 2-group  $P$  of order at least 8, we classify the splendid Morita equivalence classes of principal 2-blocks with defect groups isomorphic to  $P$ . To this end we construct explicit stable equivalences of Morita type induced by specific Scott modules using Brauer indecomposability and gluing methods; we then determine when these stable equivalences are actually Morita equivalences, and hence automatically splendid Morita equivalences. Finally, we compute the generalised decomposition numbers in each case.

- 9 越谷重夫 <sup>b</sup> 一般四元数群を Sylow 2-部分群に持つ有限群の主ブロックの splendid Morita 同値について ..... 10  
 (千葉大先進科学センター・千葉大\*)

C. Lassueur (TU Kaiserslautern)

Shigeo Koshitani

(Chiba Univ./Chiba Univ.\*)

Splendid Morita equivalences for principal blocks with generalised quaternion defect groups

Caroline Lassueur (TU Kaiserslautern)

概要 We prove that splendid Morita equivalences between principal blocks of finite groups with dihedral Sylow 2-subgroups realised by Scott modules can be lifted to splendid Morita equivalences between principal blocks of finite groups with generalised quaternion Sylow 2-subgroups realised by Scott modules.



- 10 伊藤 眞麻 (京大情報) 双直交多項式から導かれる平面分割の積型母関数 ..... 10  
 上岡 修平 (京大情報)  
 Mawo Ito (Kyoto Univ.) A product formula for plane partitions derived from a biorthogonal poly-  
 Shuhei Kamioka (Kyoto Univ.) nomial

概要 Plane partition is a two-dimensional array of nonnegative integers which is not increasing in rows and columns. Recently, some new generating functions of plane partitions are studied in view of orthogonal polynomials and integrable systems. We use Kamioka's framework that interpret biorthogonal polynomials in terms of lattice paths and derive some combinatorial identities that generalizes the generating functions derived by MacMahon and Stanley.

#### 14:15~15:15 特別講演

- 田邊 顕一朗 (北大理)<sup>b</sup> 非退化偶格子に付随する頂点代数の不変部分代数の表現  
 Kenichiro Tanabe (Hokkaido Univ.) Representations of some fixed point subalgebra of the vertex algebra  
 associated to a non-degenerate even lattice

概要 The notion of a vertex algebra was introduced by Borchers in 1986 and is a mathematically precise algebraic counterpart of the concept of a 2-dimensional conformal field theory in physics. The vertex algebra  $V_L$  associated to a non-degenerate even lattice  $L$  is a basic example of a vertex algebra. In this talk, we discuss the fixed point subalgebra  $V_L^+$  of  $V_L$  under the action of the automorphism of  $V_L$  induced from the  $-1$  symmetry of  $L$ . The fixed point subalgebras play an important role in the study of vertex algebras, particularly in the construction of interesting examples. For example, the moonshine vertex algebra  $V^\natural$  is constructed from  $V_\Lambda^+$  and its irreducible module where  $\Lambda$  is the Leech lattice.

When  $L$  is positive definite, the representations of  $V_L^+$  have been intensively studied. However, when  $L$  is not positive definite, little is known about the representations of  $V_L^+$  so far. I will talk about the classification of irreducible weak  $V_L^+$ -modules for any non-degenerate even lattice  $L$ .

#### 15:30~16:50

- 11 上田 衛 (京大理) Affine super Yangian ..... 10  
 Mamoru Ueda (Kyoto Univ.) Affine super Yangian

概要 In this talk, we define the affine super Yangian  $Y_{\varepsilon_1, \varepsilon_2}(\widehat{\mathfrak{sl}}(m|n))$  with a coproduct structure. We also obtain an evaluation homomorphism, that is, an algebra homomorphism from  $Y_{\varepsilon_1, \varepsilon_2}(\widehat{\mathfrak{sl}}(m|n))$  to the completion of the universal enveloping algebra of  $\widehat{\mathfrak{gl}}(m|n)$ .

- 12 浦野 慧 (筑波大数理物質) Modular moonshine ..... 10  
 Satoru Urano (Univ. of Tsukuba) Modular moonshine

概要 The original modular moonshine conjectures of Ryba asserted the existence of vertex algebras over finite fields with finite group actions, such that the graded Brauer characters are genus zero modular functions. Borchers and Ryba reinterpreted the conjectures in terms of Tate cohomology of a prime order element of the monster acting on an integral form of the monster vertex algebra. In this talk, I shall briefly explain the modular moonshine conjectures and consider generalized modular moonshine for any element of the monster group.

- 13 S. Carnahan (筑波大数理物質) Monstrous moonshine over the integers ..... 10  
 Scott Carnahan (Univ. of Tsukuba) Monstrous moonshine over the integers

概要 We briefly describe the construction of a self-dual integral form of the "Moonshine Module" vertex operator algebra that has Monster symmetry. This construction resolves the last open assumption in Borchers and Ryba's proof of the Modular Moonshine conjectures. As a corollary, we obtain a positive definite unimodular lattice of rank 196884 with a faithful Monster action.

- 14 川 節 和 哉 (熊本大先導機構) アフィン頂点作用素代数の relaxed 最高ウェイト表現 ..... 10  
 D. Ridout (Univ. of Melbourne)  
Kazuya Kawasetsu (Kumamoto Univ.) Relaxed highest-weight modules over affine vertex operator algebras  
 David Ridout (Univ. of Melbourne)

概要 Recently, irrational vertex operator algebras are becoming more and more important. Affine vertex operator algebras  $L_k(\mathfrak{g})$  with fractional levels  $k$  are one of the most fundamental examples of such algebras. T. Creutzig and D. Ridout investigated a category of *relaxed highest-weight modules* (and their twists) over  $L_k(\mathfrak{sl}_2)$  and proposed a continuous analog of modular invariance of characters and Verlinde formula. Today, we explain recent developments on classification and characters of relaxed highest-weight modules over affine vertex operator algebras for general  $\mathfrak{g}$ .

- 15 島 倉 裕 樹 (東北大情報) On automorphism groups of the holomorphic VOAs associated with Niemeier lattices and the  $-1$ -isometries ..... 10  
 Hiroki Shimakura (Tohoku Univ.) On automorphism groups of the holomorphic VOAs associated with Niemeier lattices and the  $-1$ -isometries

概要 Recently, it has been proved that there exist exactly 70 holomorphic VOAs of central charge 24 whose weight one Lie algebras are non-trivial. One of the next projects is to determine the automorphism groups of holomorphic VOAs of central charge 24. We discuss the 14 cases related to Niemeier lattices and the  $-1$ -isometries.

- 16 林 正 洪 (中華民国中央研究院) On a  $c=33$  extremal VOA ..... 10  
山 内 博 (東京女大現代教養)  
 Ching Hung Lam (Academia Sinica) On a  $c=33$  extremal VOA  
Hiroshi Yamauchi  
 (Tokyo Woman's Christian Univ.)

概要 Tener–Wang and Grady–Tener classified rank two extremal VOAs. In this talk I will construct an example of a  $c=33$  extremal VOA as a framed VOA which fills the last missing piece of their classification. This is a joint work with Ching Hung Lam.

- 17 浅 井 聡 太 (京大数理研) ねじれ類がなす束における広大区間 ..... 10  
 Sota Asai (Kyoto Univ.) Wide intervals in lattices of torsion classes

概要 This talk is based on joint work with Calvin Pfeifer (Bonn). For a fixed abelian length category  $\mathcal{A}$ , the poset  $\text{tors } \mathcal{A}$  of torsion classes in  $\mathcal{A}$  is a lattice. Any interval  $[\mathcal{U}, \mathcal{T}]$  in  $\text{tors } \mathcal{A}$  is a sublattice of  $\text{tors } \mathcal{A}$ , and the difference between the two torsion classes  $\mathcal{U}$  and  $\mathcal{T}$  is described by the subcategory  $\mathcal{W} := \mathcal{U}^\perp \cap \mathcal{T}$ . Motivated by  $\tau$ -tilting reduction of Jasso, we mainly studied the case that  $\mathcal{W}$  is a wide subcategory of  $\mathcal{A}$ ; such  $[\mathcal{U}, \mathcal{T}]$  are called wide intervals. In this talk, I will explain our main result that a wide interval  $[\mathcal{U}, \mathcal{T}]$  is isomorphic to the lattice  $\text{tors } \mathcal{W}$  of torsion classes in the abelian category  $\mathcal{W}$ . If time permits, I would like to talk about some characterization of wide intervals obtained in our work.

3月17日(火) 第I会場

10:00~11:45

- 18 佐 藤 眞 久 (愛知大地域政策) Ware 問題に関連したある種の環と加群の例 ..... 10  
 Masahisa Sato (Aichi Univ.) Some examples of rings and modules relating to Ware's problem

概要 We give some example of rings and modules, which gives a counter example of the assertion that Ware's problem is negative.

- 19 柴田 義大 (山口大創成) Dual square full 加群について ..... 10  
 菊政 勲 (山口大理)  
 倉富 要輔 (山口大理)  
 Yoshiharu Shibata (Yamaguchi Univ.) On dual square full modules  
 Isao Kikumasa (Yamaguchi Univ.)  
 Yosuke Kuratomi (Yamaguchi Univ.)

概要 A module  $M$  is said to be *dual square free* if, for any nonzero module  $N$ , there are no epimorphisms from  $M$  to  $N^2$ . Oppositely, a module  $M$  is called *dual square full* if, for any proper submodule  $X$  of  $M$ , there exist a proper submodule  $Y$  of  $M$  containing  $X$  and an epimorphism from  $M$  to  $(M/Y)^2$ . We will give a decomposition of quasi-discrete modules using the concepts “dual square free” and “dual square full”, and show a condition for a quasi-discrete module to be quasi-projective as its application.

- 20 板場 綾子 (東京理大理) 3次元 quadratic AS 正則環に付随する非可換射影スキームの中心上有限生成について ..... 10  
 Ayako Itaba (Tokyo Univ. of Sci.) On finite generations over centers of non-commutative projective schemes associated to 3-dimensional quadratic AS-regular algebras

概要 In this talk, for generic 3-dimensional quadratic AS-regular algebras  $A = \mathcal{A}(E, \sigma)$ , we show that a non-commutative projective schemes  $\text{Proj}_{\text{nc}} A$  associated to  $A$  is finitely generated over its center if and only if the norm  $\|\sigma\|$  of  $\sigma$  is finite. This result is a generalization of Mori’s result for a 3-dimensional quadratic AS-regular algebra of Type S.

- 21 松野 仁樹 (静岡大理) 点スキームが楕円曲線である 3次元 quadratic AS 正則環の分類 ..... 10  
 Masaki Matsuno (Shizuoka Univ.) The classification of 3-dimensional quadratic AS-regular algebras whose point schemes are elliptic curves

概要 It is known that there is a one-to-one correspondence between 3-dimensional quadratic AS-regular algebras and regular geometric pairs. In this talk, we give a complete list of regular geometric pairs which correspond to 3-dimensional quadratic AS-regular algebras whose point schemes are elliptic curves in  $\mathbb{P}^2$  up to isomorphism.

- 22 Ji-Wei He Hopf dense Galois extensions over a ring ..... 10  
 (Hangzhou Normal Univ.)  
 Haigang Hu (静岡大創造科学技術)  
 Ji-Wei He (Hangzhou Normal Univ.) Hopf dense Galois extensions over a ring  
 Haigang Hu (Shizuoka Univ.)

概要 Let  $R$  be a commutative domain, let  $H$  be a Hopf  $R$ -algebra which is a finitely generated free  $R$ -module, and let  $A$  be an  $R$ -algebra which is also an  $H$ -comodule algebra. We will say that  $A/A^{coH}$  is a Hopf dense Galois extension if the cokernel of the associated canonical map  $\beta : A \otimes_{A^{coH}} A \rightarrow A \otimes_R H$  is quotient finite. It is a generalization of a Hopf dense Galois extension over a field. A weaker version of Auslander theorem holds for Hopf dense Galois extensions over  $R$ . We also give a sufficient condition for that the localization of  $H$  is dual to a finite dimensional group algebra over an algebraic closed field containing  $R$ .

- 23 松井 紘樹 (東大数理) 三角圏のスペクトラムとその可換環論への応用 ..... 10  
 Hiroki Matsui (Univ. of Tokyo) Construction of spectra of triangulated categories and their applications  
 to commutative algebra

概要 Classification of thick subcategories has been one of the main approaches in the studies of triangulated categories. It has been studied so far in many areas. In this century, a beautiful theory, called *tensor-triangular geometry*, is initiated by Balmer. He defined a topological space for a given tensor-triangulated category. Using this topological space, he classified radical thick tensor ideals and this result enables us to do algebro-geometric studies of tensor-triangulated categories. However the Balmer theory is successful, it does not work for triangulated categories that are not tensor triangulated. Such triangulated categories include two of the most important triangulated categories in commutative algebra: the bounded derived category and the singularity category of a commutative noetherian ring. In this talk, I will introduce a way to construct spectra for (not necessarily tensor) triangulated categories and give applications to commutative algebra.

- 24 平野 文菜 (北見工大工) 単純グラフの3種のマッチング数とエッジイデアルの次元について ..... 10  
 松田 一徳 (北見工大工)  
 Ayana Hirano (Kitami Inst. of Tech.) Matching numbers and dimension of edge ideals  
 Kazunori Matsuda  
 (Kitami Inst. of Tech.)

概要 We determine the relationship between the induced matching number, the minimum matching number, the matching number and the dimension of edge ideals.

- 25 宮崎 充弘 (京都教育大) On the traces of the canonical modules of the Ehrhart rings of order  
 J. Page (Univ. of Michigan) and chain polytopes ..... 10  
 Mitsuhiro Miyazaki On the traces of the canonical modules of the Ehrhart rings of order  
 (Kyoto Univ. of Edu.) and chain polytopes  
 Janet Page (Univ. of Michigan)

概要 Let  $R$  be a commutative ring and  $M$  an  $R$ -module. The ideal  $\sum_{\varphi \in \text{Hom}_R(M, R)} \varphi(M)$  is called the trace of  $M$ . It is known that if  $R$  is a Cohen–Macaulay local ring with a canonical module  $\omega$ , then the trace of  $\omega$  is a defining ideal of non-Gorenstein locus of  $R$ . In this talk, we state a criterion of a given Laurent monomial is a member of the radical of the trace of the canonical module of the Ehrhart ring of the order polytope (resp. chain polytope).

- 26 東谷 章弘 (阪大情報) Generalized permutohedron のトーリックイデアル ..... 10  
 大杉 英史 (関西学院大理工)  
 Akihiro Higashitani (Osaka Univ.) Toric ideals of generalized permutohedra  
 Hidefumi Ohsugi  
 (Kwansei Gakuin Univ.)

概要 In this talk, we discuss the toric ideals of generalized permutohedra. More precisely, we prove that the toric ideal of Minkowski sum of unit simplices has a squarefree initial ideal and is generated by quadratic binomials. Moreover, we also prove that Minkowski sums of unit simplices have the integer decomposition property. Those results are a partial contribution to Oda conjecture and Bøgvad conjecture.

## 13:00~14:15

- 27 大杉英史 (関西学院大理工) Unimodular 配置に付随する nef 分割 ..... 10  
 土谷昭善 (東大数理)  
 Hidefumi Ohsugi (Kwansei Gakuin Univ.) Nef-partitions arising from unimodular configurations  
 Akiyoshi Tsuchiya (Univ. of Tokyo)

概要 Reflexive polytopes have been studied from viewpoints of combinatorics, commutative algebra and algebraic geometry. A nef-partition of a reflexive polytope  $\mathcal{P}$  is a decomposition  $\mathcal{P} = \mathcal{P}_1 + \cdots + \mathcal{P}_r$  such that each  $\mathcal{P}_i$  is a lattice polytope containing the origin. Batyrev and van Straten gave a combinatorial method for explicit constructions of mirror pairs of Calabi–Yau complete intersections obtained from nef-partitions. In this talk, by means of Gröbner basis techniques, we give a large family of nef-partitions arising from unimodular configurations.

- 28 土谷昭善 (東大数理) イニシャルイデアルとその深度 ..... 10  
 日比孝之 (阪大情報)  
 Akiyoshi Tsuchiya (Univ. of Tokyo) Initial ideals and their depth  
 Takayuki Hibi (Osaka Univ.)

概要 In this talk, given an arbitrary integer  $d > 0$ , we construct a homogeneous ideal  $I$  of the polynomial ring  $S = K[x_1, \dots, x_{3d}]$  in  $3d$  variables over a field  $K$  for which  $S/I$  is a Cohen–Macaulay ring of dimension  $d$  with the property that, for each of the integers  $0 \leq r \leq d$ , there exists a monomial order  $<_r$  on  $S$  with  $\text{depth}(S/\text{in}_{<_r}(I)) = r$ , where  $\text{in}_{<_r}(I)$  is the initial ideal of  $I$  with respect to  $<_r$ .

- 29 C. McDaniel (Endicott Coll.) Equivariant coinvariant rings of finite groups ..... 10  
 Chris McDaniel (Endicott Coll.) Equivariant coinvariant rings of finite groups

概要 Let  $W$  be a finite group acting linearly on a polynomial ring  $R$ , and let  $R^W \subset R$  be its invariant subring. The equivariant coinvariant ring of  $W$  is defined to be the tensor product  $R \otimes_{R^W} R$ ; if  $W$  is a Weyl group it is isomorphic to the equivariant cohomology ring of the associated flag variety with respect to its natural torus action. In a 1976 paper, J. Watanabe obtained two important results: (1)  $R \otimes_{R^W} R$  is Cohen–Macaulay if and only if  $W$  is generated by pseudo-reflections, and (2) if  $W$  is generated by pseudo-reflections then  $R \otimes_{R^W} R$  is a reduced ring. In this talk, we will discuss these and other more recent related results obtained in part in collaboration with J. Watanabe and L. Smith.

- 30 渡辺純三 (東海大\*) A new definition of the principal radical system and an application to  
 C. McDaniel (Endicott Coll.) Specht ideals of type  $(n-k, k)$  ..... 10  
 Junzo Watanabe (Tokai Univ.\*) A new definition of the principal radical system and an application to  
 Chris McDaniel (Endicott Coll.) Specht ideals of type  $(n-k, k)$

概要 The notion of the principal radical system was introduced by Hochster and Eagon in 1971 and they used it to prove that the determinantal ideals are radical and Cohen–Macaulay. We will simplify the definition to prove that the Specht ideals of type  $(n-k, k)$  are radical in any characteristic and Cohen–Macaulay in characteristic zero, independent of the proofs of Yanagawa and Etingof.

- 31 柴田 孝祐 (岡山大自然) Hilbert series of Cohen–Macaulay Specht ideals ..... 10  
 柳川 浩二 (関西大システム理工)  
 Kohsuke Shibata (Okayama Univ.) Hilbert series of Cohen–Macaulay Specht ideals  
 Kohji Yanagawa (Kansai Univ.)

概要 For a partition  $\lambda$  of positive integer  $n$ , let  $I_\lambda^{\text{Sp}}$  be the ideal of  $R = K[x_1, \dots, x_n]$  generated by all Specht polynomials of shape  $\lambda$ . It is known that if  $R/I_\lambda^{\text{Sp}}$  is Cohen–Macaulay then  $\lambda$  is of the form either  $(n-d, 1, \dots, 1)$ ,  $(n-d, d)$ , or  $(d, d, 1)$ , and it is also known that the converse is true if  $\text{char}(K) = 0$ . In this talk, we compute that the (Castelnuovo–Mumford) regularity of  $R/I_\lambda^{\text{Sp}}$ , when  $\lambda = (n-d, d)$  or  $(d, d, 1)$ . More precisely, the regularities of  $R/I_\lambda^{\text{Sp}}$  for  $(n-d, d)$  and  $(d, d, 1)$  are  $d$  and  $d+1$ , respectively.

- 32 S. Masuti (Chennai Math. Inst.) On the structure of the Sally module and the second normal Hilbert  
 大関 一秀 (山口大創成) coefficient ..... 10  
 M. E. Rossi (Genova Univ.)  
 H. L. Truong (Saarlandes Univ.)  
 Shreedevi Masuti (Chennai Math. Inst.) On the structure of the Sally module and the second normal Hilbert  
 Kazuho Ozeki (Yamaguchi Univ.) coefficient  
 Maria Evelina Rossi (Genova Univ.)  
 Hoang Le Truong (Saarlandes Univ.)

概要 The Sally module was introduced by W. V. Vasconcelos and it is useful to connect the Hilbert coefficients to the homological properties of the associated graded module of a Noetherian filtration. In this report we give a complete structure of the Sally module in the case the second normal Hilbert coefficient attains almost minimal value in an analytically unramified Cohen–Macaulay local ring. As a consequence, in this case we present a complete description of the Hilbert function of the associated graded ring of the normal filtration.

3月18日(水) 第I会場

9:30~12:00

- 33 岩見 智宏 (九工大工)<sup>b</sup> Semistable extremal neighborhoods rigged by framed-form fans of cuspidal  
 type singularities ..... 10  
 Tomohiro Iwami (Kyushu Inst. of Tech.) Semistable extremal neighborhoods rigged by framed-form fans of cuspidal  
 type singularities

概要 For a three-dimensional extremal neighborhood  $(X, C) \subset \mathbb{C}^4$  of semistable type, infinitesimal deformations of  $C$  and the abundance property contribute to the existence of three-dimensional flips for  $(X, C)$  (S. Mori, 1988). Based on these facts, the author gave an analogue of Miyaoka–Yau type inequality for  $(X, C)$  with the associated  $c_3$ , and gave a slight generalization of such an inequality with non-trivial coefficients of  $c_2$ , by the associated Higgs sheaves (Y. Miyaoka, 2009) and formal neighborhoods of  $C$  (M. Ebihara, 1992), for related local automorphisms. In this talk, by framed-form fans (or, T-complexes) as a toric analogue of formal scheme (M. Ishida, 1989–), the author will give: i) a concept of extremal neighborhood rigged by the framed-form fan  $\Gamma := (\Gamma_a, C_a, K_a)$  associated to  $(X, C)$ , denoted as  $(X, \Gamma)$ , ii) infinitesimal deformations of  $(X, \Gamma)$  by the associated Higgs sheaves, and, iii) a proto-type of characterization of Mukai–Umemura 3-folds by ii).

- 34 縫田 光司 (東大情報理工) 楕円曲線上の群構造に関する線型代数のみで完結する証明 ..... 10  
Koji Nuida (Univ. of Tokyo) An elementary linear-algebraic proof for group law on elliptic curves

概要 The group structure on the rational points of elliptic curves plays several important roles, in mathematics and recently also in other areas such as cryptography. However, the famous proofs for the group property (in particular, for its associative law) require somewhat advanced mathematics and therefore are not easily accessible by non-mathematician. This talk introduces a self-contained proof for this property, assuming mathematical knowledge only at the level of basic linear algebra and not requiring heavy computation.

- 35 桜井 真 (開智学園) カイラル代数の拡張と応用について ..... 10  
Makoto Sakurai (Kaichi Gakuen) Extension and applications of chiral algebra theory

概要 The formalism of chiral conformal field theory by Beilinson and Drinfeld was inspired by the perturbative physics of Costello, Nekrasov, and Witten. However, the progress does not acquire sufficient consensus of both mathematicians and physicists. This “chiral algebra theory” was applied from a kind of tensor category theory in the operads to cousins of vertex algebras. In the topological field theory, this theory has a “factorization algebra” theory with a sophisticated theory of higher categories. In my talk, I will report two trials of weakening the toric condition and seeking for the hidden relation to topological vertex theory.

- 36 土基善文 (高知大理工) 非可換複素ケーラー射影平面上の代数曲線について ..... 10  
Yoshifumi Tsuchimoto (Kochi Univ.) On curves on the non-commutative complex Kähler plane

概要 By using a non-commutative analogue of Marsden–Weinstein quotients, we may define non commutative complex Kähler plane. In this talk we talk about curves on it.

- 37 安藤 哲哉 (千葉大理) 半代数多様体上の PSD 錐の理論 ..... 10  
Tetsuya Ando (Chiba Univ.) Theory of PSD cones on semialgebraic varieties

概要 For study of algebraic inequalities, theory of PSD (positive semidefinite) cones provides various new methods. A merits of theory of PSD cones is that we can use many results and ideas of algebraic geometry. Moreover, we can understand algebraic and geometric meanings of inequalities. For an example, we proved that the edge discriminant of cyclic homogeneous inequalities with three variables of degree  $d$  agrees with the discriminant of one variable equations of degree  $d$ . For an another example, we proved that extremal rays are determined by equality conditions. In PSD method, we determine the characteristic variety, and its critical decomposition. Irreducible components of the boundary of the PSD cone are obtained as the dual of a critical set. Using this idea, we determine extremal rays and discriminants.

- 38 南 範彦 (名工大) 一般 Bott 塔を通した, 高次単線織性=低次単有理性よりも強い階層構造  
を与えるある十分条件の適応可能性について ..... 10  
Norihiro Minami (Nagoya Inst. of Tech.) On the applicability of the sufficient criterion for a stronger hierarchy  
of higher uniruledness = lower unirationality via Bott tower

概要 Recently, the author obtained a sufficient criterion for a stronger hierarchy of higher uniruledness = lower unirationality via Bott tower. In this talk, the author clarifies the following points: (1) This sufficient criterion is actually for a stronger hierarchy than the natural hierarchy arising from higher uniruledness = lower unirationality, especially from the view point of Hodge conjecture. (2) Still, this sufficient criterion has some reasonable acceptable applicability. From this reason, the author would like to call those Fano manifolds satisfying this sufficient criterion as Higher Fano manifolds.

- 39 加藤 芳文 (名城大理工) Curvature matrix of the universal bundle of the Grassmann variety ... 10  
 Yoshifumi Kato (Meijo Univ.) Curvature matrix of the universal bundle of the Grassmann variety

概要 We introduce a local coordinates system to the Grassmann variety  $Gr(k; n)$  and express the curvature matrix of the universal bundle  $E$  simply and independently of the choice of local coordinates. We state the relation between the local coordinate system and the Schubert cells of  $Gr(k; n)$ . By using and the expression of the cells, we obtain explicit integral formulas corresponding to Young diagrams.

- 40 加藤 芳文 (名城大理工) An observation on Schubert polynomials ..... 10  
 Yoshifumi Kato (Meijo Univ.) An observation on Schubert polynomials

概要 We introduce a local coordinates system to the Flag variety and construct a vector field which is expressed explicitly with respect to the system. Each Schubert cell becomes a union of some flows of the vector field. We associate a certain diagram to the cell and investigate the relation between the diagrams and Schubert polynomials. We present some conjectures.

- 41 松原 祐貴 (神戸大理) 従順分岐型の幾何学的ラングランズ対応について ..... 10  
 Yuki Matsubara (Kobe Univ.) Tamely ramified geometric Langlands correspondence

概要 We consider the moduli space of logarithmic connections of rank 2 on the projective line minus 5 points with fixed spectral data. We compute the cohomology of such moduli space, and this computation will be used to extend the results of geometric Langlands correspondence due to D. Arinkin to the case where this type of connections have five simple poles on  $P^1$ . In this talk, I will review the geometric Langlands correspondence in the tamely ramified cases, and after that, I will explain how the cohomology of above moduli space will be used.

- 42 白根 竹人 (徳島大理工) Torsion divisors of plane curves and Zariski pairs ..... 10  
 E. Artal Bartolo (Univ. Zaragoza)  
 坂内 真三 (茨城工高専)  
 徳永 浩雄 (首都大東京理)  
 Taketo Shirane (Tokushima Univ.) Torsion divisors of plane curves and Zariski pairs  
 Enrique Artal Bartolo (Univ. Zaragoza)  
 Shinzo Bannai  
 (Ibaraki Nat. Coll. of Tech.)  
 Hiro-o Tokunaga (Tokyo Metro. Univ.)

概要 We study the embedded topology of reducible plane curves having a smooth irreducible component. In previous studies, the relation between the topology and certain torsion classes in the Picard group of degree zero of the smooth component was implicitly considered. We formulate this relation clearly and give a criterion for distinguishing the embedded topology in terms of torsion classes.

- 43 佐藤 宏平 (小山工高専) 3次元アーベル商特異点に対するクレパント特異点解消と足利連分数展開  
 佐藤 悠介 (東大数理) について ..... 10  
 Kohei Sato (Oyama Nat. Coll. of Tech.) On Ashikaga's continued fractions and crepant resolutions for 3-dimensional  
 Yusuke Sato (Univ. of Tokyo) Abelian quotient singularities

概要 The existence of crepant resolutions for three dimensional Gorenstein quotient singularities has proved by Y. Ito, D. G. Markushevich, S. S. Roan on a case-by-case based on the classification table of the finite subgroups of  $SL(3, \mathbb{C})$ .

In this talk, we shall discuss a sufficient condition of the existence of crepant resolutions for Abelian quotient singularities by using Ashikaga's continued fractions. As a corollary, we have an alternative proof of the abelian case of the above results.



- 44 長岡 高広 (京大 理) 超トーリック多様体の普遍被覆とボゴモロフ分解 ..... 10  
Takahiro Nagaoka (Kyoto Univ.) The universal covers of hypertoric varieties and Bogomolov's decomposition

概要 Hypertoric varieties are singular symplectic varieties with good positive weight  $\mathbb{C}^*$ -actions, and they can be studied from the corresponding combinatorial objects (hyperplane arrangements) as toric varieties. In general, for such symplectic varieties, one can consider the universal covering which is also a symplectic variety. In this talk, we describe the universal covers of hypertoric varieties explicitly, and we also establish the analogue of Bogomolov's decomposition for them. Moreover, as a byproduct, we can show that the  $\mathbb{C}^*$ -equivariant isomorphism class of hypertoric varieties is classified by the associated combinatorial objects.

- 45 赤池 広都 (阪大 理)<sup>b</sup> Slope inequalities for irregular cyclic covering fibration ..... 10  
Hirotu Akaike (Osaka Univ.) Slope inequalities for irregular cyclic covering fibration

概要 The relative invariant slope is defined as the ratio of the self intersection number of relative canonical divisor and the relative Euler number. In this time, I studied the slope of primitive cyclic covering fibration with positive relative irregularity. I obtained the lower bound described by the genus of general fiber, covering degree and relative irregularity.

**14:15~14:30 2020年度(第23回)日本数学会代数学賞授与式**

**14:40~15:40 2020年度(第23回)日本数学会代数学賞受賞特別講演**

高橋 亮 (名大 多元数理) 可換環の加群圏と導来圏における生成問題

Ryo Takahashi (Nagoya Univ.) Generation in module categories and derived categories of commutative rings

概要 Let  $R$  be a ring, and let  $M, N$  be  $R$ -modules. It is a natural question to ask whether or how one can build  $M$  out of  $N$  by iteration of fundamental operations such as direct sums, direct summands, extensions etc. This can be thought of not only in module categories but also in derived categories. I will speak about it in the case where the base ring is commutative.

**15:50~16:50 2020年度(第23回)日本数学会代数学賞受賞特別講演**

岡田 拓三 (佐賀大 理工) ファノ多様体の双有理的森ファイバー構造の研究と有理性問題への応用

Takuzo Okada (Saga Univ.) Birational Mori fiber structures of Fano varieties and its application to rationality problems

概要 I will talk about birational geometry of Fano varieties. Fano varieties of Picard number 1 are so called Mori fiber spaces, that is, they appear as outputs of Minimal Model Program (MMP). Outputs of MMP are not necessarily unique and in general a Fano variety can be birationally transformed into other Mori fiber spaces. For a given Fano variety of Picard number 1, we can sometimes determine the Mori fiber structure of it, that is, the Mori fiber spaces birational to the given Fano variety. I will survey results concerning this kind of birational studies on Fano varieties and explain a direct connection to rationality problems of algebraic varieties.

3月19日(木) 第I会場

9:20~12:00

- 46 丸山文綱 Recent progress on Euler–Fermat type theorem on matrix ring …… 10  
 豊泉正男 (東洋大理工)  
 出口洋三  
Fumitsuna Maruyama Recent progress on Euler–Fermat type theorem on matrix ring  
 Masao Toyozumi (Toyo Univ.)  
 Yozo Deguchi

概要 We share new results for Euler- Fermat type theorem on the matrix ring over  $F_p$ .

- 47 吉野聖人 (東北大情報) Non 2-integrable lattices of rank 12 …… 10  
 Qianqian Yang (中国科学技術大)  
Kiyoto Yoshino (Tohoku Univ.) Non 2-integrable lattices of rank 12  
 Qianqian Yang  
 (Univ. of Sci. Tech. China)

概要 For a positive integer  $s$ , a lattice  $L$  is said to be  $s$ -integrable if  $\sqrt{s} \cdot L$  is isomorphic to a sublattice of  $\mathbb{Z}^n$  for some integer  $n$ . Conway and Sloane found two non 2-integrable lattices of rank 12 and discriminant 7. We found two more lattices of rank 12 and discriminant 15. By a joint work with Dr. Qianqian Yang, we derived a necessary condition for the discriminant of a non 2-integrable lattice of rank 12 by embedding a lattice of rank 12 to a unimodular lattice of rank 14. Furthermore, we give a sufficient condition for a non 2-integrable lattice to be indecomposable in terms of quadratic forms, and show that Conway and Sloane's lattices and ours satisfy it.

- 48 星明考 (新潟大理) Norm one tori and Hasse norm principle …… 10  
 金井和貴 (新潟大自然)  
 山崎愛一 (京大理)  
Akinari Hoshi (Niigata Univ.) Norm one tori and Hasse norm principle  
 Kazuki Kanai (Niigata Univ.)  
 Aiichi Yamasaki (Kyoto Univ.)

概要 Hoshi and Yamasaki (2017, Mem. AMS) classified stably/retract  $k$ -rational algebraic  $k$ -tori of dimensions 4 and 5. There exist 710 (resp. 6079) cases of 4-dim. (resp. 5-dim.) algebraic  $k$ -tori and 216 (resp. 3003) cases of them are not retract  $k$ -rational. Using this, we prove:

Theorem (Hoshi, Kanai and Yamasaki, arXiv:1910.01469). Let  $k$  be a field,  $T$  be an algebraic  $k$ -torus of dimension  $n$  and  $X$  be a smooth  $k$ -compactification of  $T$ .

(1) Among 216 cases of not retract rational algebraic  $k$ -tori  $T$  of dimension 4, there exist exactly 2 (resp. 20, 194) algebraic  $k$ -tori with  $H^1(k, \text{Pic } \bar{X}) \simeq (\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$  (resp.  $H^1(k, \text{Pic } \bar{X}) \simeq \mathbb{Z}/2\mathbb{Z}$ ,  $H^1(k, \text{Pic } \bar{X}) = 0$ ).

(2) Among 3003 cases of not retract rational algebraic  $k$ -tori  $T$  of dimension 5, there exist exactly 11 (resp. 263, 2729) algebraic  $k$ -tori with  $H^1(k, \text{Pic } \bar{X}) \simeq (\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$  (resp.  $H^1(k, \text{Pic } \bar{X}) \simeq \mathbb{Z}/2\mathbb{Z}$ ,  $H^1(k, \text{Pic } \bar{X}) = 0$ ).

- 49 星 明 考 (新潟大理) Norm one tori and Hasse norm principle, II ..... 10  
 金 井 和 貴 (新潟大自) .....  
 山 崎 愛 一 (京 大 理) .....  
 Akinari Hoshi (Niigata Univ.) Norm one tori and Hasse norm principle, II  
 Kazuki Kanai (Niigata Univ.) .....  
 Aiichi Yamasaki (Kyoto Univ.) .....

概要 Theorem (Hoshi, Kanai and Yamasaki, arXiv:1910.01469). Let  $2 \leq n \leq 15$  be an integer with  $n \neq 12$ . Let  $k$  be a field,  $K/k$  be a separable field extension of degree  $n$  and  $L/k$  be the Galois closure of  $K/k$ . Let  $G = \text{Gal}(L/k) = nTm$  be a transitive subgroup of  $S_n$  and  $H = \text{Gal}(L/K)$  with  $[G : H] = n$ . Let  $T = R_{K/k}^{(1)}(\mathbb{G}_m)$  be the norm one torus of  $K/k$  of dimension  $n - 1$  and  $X$  be a smooth  $k$ -compactification of  $T$ . Then  $H^1(k, \text{Pic } \overline{X}) \neq 0$  if and only if  $G$  is given as in Table 1. In particular, if  $L/k$  is unramified extension, then  $A(T) = 0$  and  $H^1(k, \text{Pic } \overline{X}) \simeq \text{Sha}(T)$ .

Moreover, we give a necessary and sufficient condition for the Hasse norm principle for  $K/k$  where  $[K : k] = n \leq 15$  and  $n \neq 12$ .

- 50 奥 村 喜 晶 (東 工 大 理) Non-existence of Drinfeld modules with constrained torsions ..... 10  
 Yoshiaki Okumura (Tokyo Tech) Non-existence of Drinfeld modules with constrained torsions

概要 In the arithmetic of function fields, Drinfeld modules play the role that elliptic curves play in the arithmetic of number fields. The aim of this talk is to study a non-existence problem of Drinfeld modules with constraints on torsion points, which is motivated by a conjecture of Rasmussen and Tamagawa on abelian varieties over number fields. We prove the non-existence in the case where the inseparable degree of the base field is not divisible by the rank of Drinfeld modules. Conversely if the rank divides the inseparable degree, then we prove the existence of a Drinfeld module satisfying Rasmussen–Tamagawa-type conditions. We also prove a partial result on the Drinfeld module analogue of the uniform Rasmussen–Tamagawa conjecture.

- 51 木 村 巖 (富山大理工) 素導手円分関数体の相対類数の評価について ..... 10  
 青 山 大 輝 (富山大理工) .....  
 Iwao Kimura (Univ. of Toyama) On an estimate of the relative class number of cyclotomic function field  
 Daiki Aoyama (Univ. of Toyama) of prime conductor

概要 We first present a comparison of two upper bounds of relative class numbers of cyclotomic function fields.

We then give some explicit lower bound of the relative class number of cyclotomic function field of a prime conductor, whose constant field is the field of 3 elements.

These bounds describe the size of determinants in a class number formula of cyclotomic function fields.

- 52 平 林 幹 人 (金 沢 工 大) A generalization of Jakubec’s formula related to the multiplication theorem for Bernoulli polynomials ..... 10  
 Mikihito Hirabayashi A generalization of Jakubec’s formula related to the multiplication theorem for Bernoulli polynomials  
 (Kanazawa Inst. of Tech.) .....

概要 In 2017 Jakubec gave a formula for the relative class number of the  $p$ -th cyclotomic field,  $p$  an odd prime, by using a determinant related to the multiplication theorem for Bernoulli polynomials. We generalize his formula to an imaginary abelian number field, and we also determine the sign of the formula, which he had not given.

- 53 加藤 裕基 (宇部工高専) Homotopy invariant  $K$ -theory of perfectoidification of regular local rings ..... 10  
 Yuki Kato (Ube Nat. Coll. of Tech.) Homotopy invariant  $K$ -theory of perfectoidification of regular local rings

概要 Let  $R$  be a (Noetherian) regular local ring. Then we can obtain there perfectoidification  $R_\infty$  such that  $R_\infty$  is a perfectoid ring which is faithfully flat over  $R$ . If  $R$  is regular coherent ring, then the  $K$ -theory has the homotopy invariant property. In this talk, we consider the étale  $K$ -theory for perfectoid algebras and we compare the étale  $K$ -theory spectrum and homotopy invariant étale  $K$ -theory spectrum of  $R_\infty$ .

- 54 太田 和惟 (慶大理工) Big Heegner points and generalized Heegner cycles ..... 10  
 Kazuto Ota (Keio Univ.) Big Heegner points and generalized Heegner cycles

概要 For Galois cohomology groups of big Galois representations attached to Hida families (of elliptic cuspforms), Howard has constructed the systems of big Heegner points. On the other hand, for members of Hida families there are Euler systems of generalized Heegner cycles, which are constructed by Bertolini–Darmon–Prasanna. Castella recently proved that the systems of big Heegner points interpolate the systems of generalized Heegner cycles. In this talk, we will explain a different approach to show the interpolation property and slightly improve Castella’s work.

- 55 境 優一 Characterization of minimal models by modular linear differential equations of order 4 and their modules ..... 10  
 (九大多重ゼータ研究センター) 永友 清和 (阪大情報)  
 Yuichi Sakai (Kyushu Univ.) Characterization of minimal models by modular linear differential equations of order 4 and their modules  
 Kiyokazu Nagatomo (Osaka Univ.)

概要 We characterize simple vertex operator algebras of CFT type satisfying several conditions which arise from a family of simple Virasoro vertex operator algebras, so-called minimal models.

- 56 武田 渉 (名大多元数理) Factorial functions represented as norm forms ..... 10  
 Wataru Takeda (Nagoya Univ.) Factorial functions represented as norm forms

概要 We consider the number of solutions of  $F(x_1, \dots, x_n) = H_l$ , where  $F$  is a norm form of an order of a number field and  $H_l$  is a variant of factorial. In a previous study, we prove that for any norm form  $F(x_1, x_2)$  of an order of a quadratic field, there exists only finitely many  $l$  such that  $H_l$  is represented as  $F$ . The aim of this talk is to show the same result for any norm form  $F(x_1, \dots, x_n)$  of an order of a number field  $K \neq \mathbf{Q}$ .

- 57 杉山 和成 (千葉工大) 概均質ゼータ関数と Katok–Sarnak 対応について ..... 10  
 Kazunari Sugiyama (Chiba Inst. of Tech.) Prehomogeneous zeta functions and the Katok–Sarnak correspondence

概要 We discuss the Katok–Sarnak correspondence (a lifting map from Maass forms of weight 0 to those of weight 1/2) from the view point of the theory of prehomogeneous zeta functions. A key role is played by the zeta functions associated with the space of symmetric matrices of degree 2 whose coefficients involve periods of Maass wave forms. Analytic properties of these zeta functions have been investigated by Fumihiko Sato, and by using his results and a Weil-type converse theorem for Maass forms, we can construct the lifting.

- 58 伊東良純(千葉大理) On special values at integers of  $L$ -functions of Jacobi theta products of weight 3 ..... 10

Ryojun Ito (Chiba Univ.) On special values at integers of  $L$ -functions of Jacobi theta products of weight 3

概要 In 2012, Rogers–Zudilin gave a hypergeometric expression of the special value at  $s = 2$  of the  $L$ -function of a cusp form of weight 2 by an analytic method. In this talk, by the Rogers–Zudilin method, we express special values at some integers of  $L$ -functions of modular forms of weight 3 that are products of the Jacobi theta series in terms of special values of the Kampé de Fériet hypergeometric function, which is a two-variable generalization of generalized hypergeometric functions.

- 59 鈴木正俊(東工大) ゼータ関数から生ずる積分作用素の族について. II ..... 10

Masatoshi Suzuki (Tokyo Tech) On a family of integral operators arising from zeta functions. II.

概要 We consider a family of integral operators arising from the Riemann zeta function and state an equivalence condition of the Riemann hypothesis in terms of such operators. This equivalence condition is related to a positive property of the Riemann xi-function studied by Lagarias and Garunkstis.

#### 14:15~15:15 特別講演

並川健一(九大数理) 保型形式の明示的構成とその岩澤理論への応用

Kenichi Namikawa (Kyushu Univ.) Explicit constructions of automorphic forms and its applications to Iwasawa theory

概要 The study of  $L$ -functions is one of major theme in number theory, since it is expected to include various arithmetic information. An important method for the study of  $L$ -functions is automorphic representation theory. By using elements in automorphic representations, which are called automorphic forms, we can observe connections between special values of  $L$ -functions and arithmetic phenomena. In this talk, we introduce how automorphic forms are used to study special values of  $L$ -functions. Namely, we discuss Iwasawa theoretic properties of certain  $L$ -functions of degree 4, which are called Asai  $L$ -functions, by constructing automorphic forms in an explicit way.

#### 15:25~16:50

- 60 峰正博(東工大) Moments of  $L$ -functions associated with cubic fields ..... 10

Masahiro Mine (Tokyo Tech) Moments of  $L$ -functions associated with cubic fields

概要 In this talk, we study  $L$ -functions arising from the factorizations of Dedekind zeta-functions of non-Galois cubic fields. We prove asymptotic formulas for their complex moments by applying the probability density function for the value-distribution of such  $L$ -functions. As an application, we obtain a cubic analogue of Siegel's formula on quadratic class number sums.

- 61 スリアジャヤアデイルマ (九大数理・理化学研) 拡張されたセルバーグクラスの  $L$  関数のスティルチェス定数の上から評価  
井上 翔太 (名大多元数理) ..... 10  
S. Saad Eddin (JKU Linz)  
Ade Irma Suriajaya (Kyushu Univ./RIKEN) An upper bound for Stieltjes constants of  $L$ -functions in the extended  
Selberg class  
Shōta Inoue (Nagoya Univ.)  
Sumaia Saad Eddin (JKU Linz)

概要 The coefficients of the Laurent expansion of the Riemann zeta function near  $s = 1$  were first studied by Stieltjes in 1885 followed by many authors, such as Briggs, Mitrović, Matsuoka, and Saad Eddin. For this reason, these coefficients are widely known as “Stieltjes constants”. An explicit upper bound for the Stieltjes constants of Dirichlet  $L$ -functions is also known. We are interested in investigating the Stieltjes constants for  $L$ -functions in the extended Selberg class. In this talk, we show an upper bound we obtained for these coefficients.

- 62 井上 翔太 (名大多元数理) On the value distribution of the Riemann zeta-function on the critical  
line ..... 10  
Shota Inoue (Nagoya Univ.) On the value distribution of the Riemann zeta-function on the critical  
line

概要 There are many remaining problems for the Riemann zeta-function such as the Riemann Hypothesis and the Lindelöf Hypothesis. The speaker derived a result on the value distribution of the Riemann zeta-function on the critical line. This result is an improvement of Jutila’s result and Radziwiłł’s result, and also related with the mean value estimates and the Lindelöf Hypothesis.

- 63 齋藤三郎 (群馬大\*・再生核研) Values of the Riemann zeta function at positive integers by means of  
松浦 勉 (群馬大理工) the division by zero calculus ..... 10  
奥村 博  
Saburo Saitoh (Gunma Univ.\* / Inst. of Reproducing Kernels) Values of the Riemann zeta function at positive integers by means of  
the division by zero calculus  
Tutomu Matsuura (Gunma Univ.)  
Hiroshi Okumura

概要 In this talk, we will consider the values of the Riemann zeta function for any positive integers by means of the division by zero calculus. In particular, the values for odd positive integers were considered as mysterious ones, however, their values may be considered as in even integers case. We will give both analytical formulas and numerical results. A basic reference is H. Okumura and S. Saitoh, Values of the Riemann Zeta Function by Means of Division by Zero Calculus, viXra:1907.0437 but we will talk more up-to-date information. Our purposes of this talk is to introduce of the division by zero calculus and to show its power.

- 64 小川原 弘士 Mahler 関数方程式系の解の代数的独立性について ..... 10  
(熊本大大学教育統括管理運営機構)  
Hiroshi Ogawara (Kumamoto Univ.) On algebraic independence of solutions for systems of algebraic Mahler  
functional equations

概要 We give conditions for algebraic independence of solutions to a certain system of algebraic Mahler functional equations, which are extensions of a result of Amou and Väinänen. As an application of our results, we give algebraic independence of special values of certain power series and infinite products.

65 D. Banerjee ( IISER ) A divisor problem on square free integers ..... 10  
南出 真 (山口大理)  
谷川 好男

Debika Banerjee (IISER) A divisor problem on square free integers  
Makoto Minamide (Yamaguchi Univ.)  
Yoshio Tanigawa

概要 We consider the average of number of divisors of square free integers  $n \leq x$ .

66 飯高 茂 (学習院大\*) (A,B,C) 完全数 ..... 10  
Shigeru Iitaka (Gakushuin Univ.\*) (A,B,C) perfect numbers

概要 Given integers  $A, B, C, D$ , if positive integers  $a$  satisfy  $A\sigma(a) + B\varphi(a) - Ca = D$  then  $a$  are said to be (A,B,C) perfect numbers with constant  $D$ .

## 幾何学

3月16日(月) 第II会場

9:40~11:40

- 1 吉田尚彦(明大理工) Adiabatic limits, theta functions, and geometric quantization ..... 15  
Takahiko Yoshida (Meiji Univ.) Adiabatic limits, theta functions, and geometric quantization

概要 In the geometric quantization of toric manifolds, Baier–Florentino–Mourão–Nunes have given a one-parameter family of compatible complex structures such that the associated Kähler polarizations converge to the real polarization determined by the moment map. One of the generalizations of the Kähler quantization to possibly non-Kähler symplectic manifolds is the  $\text{Spin}^c$  quantization. In this talk, for a non-singular Lagrangian torus fibration on a compact, complete base with prequantum line bundle and a compatible almost complex structure invariant along the fiber, we show that the  $\text{Spin}^c$  quantization converges to the real quantization by the adiabatic(-type) limit. This talk is based on arXiv:1904.04076.

- 2 桑田健(北大理) 正則ベクトル場と  $\mathbb{P}^1$  上の位相的シグマモデル ..... 15  
秦泉寺雅夫(北大理)  
Ken Kuwata (Hokkaido Univ.) Holomorphic vector field and topological sigma model on  $\mathbb{P}^1$  world sheet  
Masao Jinzenji (Hokkaido Univ.)

概要 Witten suggested that fixed-point theorems can be derived by the supersymmetric sigma model on a Riemann manifold  $M$  with potential term induced from Killing vector on  $M$ . One of the well-known fixed-point theorem is the Bott residue formula which represents intersection number of Chern classes of holomorphic vector bundles on a Kähler manifold  $M$  as sum of contributions from fixed point sets of a holomorphic vector field  $K$  on  $M$ . We derived the Bott residue formula by using topological sigma model (A-model) that describes dynamics of maps from  $CP^1$  to  $M$ , with potential term induced from the vector field  $K$ .

- 3 宮武夏雄(阪大理)<sup>b</sup> A direct proof of Hitchin–Kobayashi type correspondences for abelian vortex equations ..... 15  
Natsuo Miyatake (Osaka Univ.) A direct proof of Hitchin–Kobayashi type correspondences for abelian vortex equations

概要 We discuss Hitchin–Kobayashi type correspondences for the abelian vortex equations, i.e., a certain kind of gauge theoretic equations on compact Kähler manifolds associated with linear actions of tori on complex vector spaces. The correspondence itself had already established for further general settings. Further for the abelian gauge group and linear fiber cases it is known that there exists a very simple characterization of the stability condition. We give a direct proof of the correspondences using such a simple characterization of the stability condition. The problem reduces to show the existence and the uniqueness of Kazdan–Warner type equations associated with linear actions of tori on complex vector spaces.

- 4 新田泰文(東京理大理) 偏極トーリック多様体における Yau–Tian–Donaldson 対応 ..... 15  
斎藤俊輔  
(理化学研 AIP・京大高等研)  
Yasufumi Nitta (Tokyo Univ. of Sci.) A uniform version of the Yau–Tian–Donaldson correspondence for polarized toric manifolds  
Shunsuke Saito (RIKEN/Kyoto Univ.)

概要 We solve a uniform version of the Yau–Tian–Donaldson conjecture for polarized toric manifolds.



- 5 長谷川和志 (金沢大人間社会) A construction of a hypercomplex manifold from a quaternionic manifold —the quaternionic/hypercomplex-correspondence— ..... 15  
 V. Cortés (University of Hamburg)

Kazuyuki Hasegawa (Kanazawa Univ.) A construction of a hypercomplex manifold from a quaternionic manifold —the quaternionic/hypercomplex-correspondence—  
Vicente Cortés (University of Hamburg)

概要 Given a quaternionic manifold  $M$  with a certain  $U(1)$ -symmetry, we construct a hypercomplex manifold  $M'$  of the same dimension. This construction generalizes the quaternionic Kähler/hyper-Kähler-correspondence. As an example of this construction, we obtain a compact homogeneous hypercomplex manifold which does not admit any hyper-Kähler structure.

- 6 橋本義規 (東工大理) Kobayashi–Hitchin correspondence and the Quot-scheme limit of Fubini–  
 J. Keller (Aix-Marseille Univ.) Study metrics ..... 15  
Yoshinori Hashimoto (Tokyo Tech) Kobayashi–Hitchin correspondence and the Quot-scheme limit of Fubini–  
 Julien Keller (Aix-Marseille Univ.) Study metrics

概要 We use the Quot-schemes in algebraic geometry to identify the asymptotic slope of Donaldson’s functional, whose critical point is a Hermitian–Einstein metric on a holomorphic vector bundle. This result, together with an extra technical assumption, provides an alternative proof of the Kobayashi–Hitchin correspondence relating the Hermitian–Einstein metrics and stability of vector bundles.

- 7 青井顕宏 (阪大理) アファイン代数多様体上の完備スカラー平坦 Kähler 計量について ..... 15  
 Takahiro Aoi (Osaka Univ.) Complete scalar-flat Kähler metrics on affine algebraic manifolds

概要 The existence of constant scalar curvature Kähler metric is a fundamental problem in Kähler geometry. Let  $(X, L_X)$  be an  $n$ -dimensional polarized manifold and  $D$  be a smooth hypersurface defined by a holomorphic section of  $L_X$ . I will talk about the following result: if  $D$  has a constant positive scalar curvature Kähler metric,  $X \setminus D$  admits a complete scalar-flat Kähler metric, under the following three conditions, (i)  $n \geq 6$  and there is no nonzero holomorphic vector field on  $X$  vanishing on  $D$ , (ii) an average of a scalar curvature on  $D$  denoted by  $\hat{S}_D$  satisfies the inequality  $0 < 3\hat{S}_D < n(n-1)$ , (iii) there are positive integers  $l(> n), m$  such that the line bundle  $K_X^{-l} \otimes L_X^m$  is very ample and the ratio  $m/l$  is sufficiently small.

#### 14:20~15:45

- 8 中村友哉 (早大理工) Jacobi 双垂代数上の Dirac ペア ..... 15  
 Tomoya Nakamura (Waseda Univ.) Dirac pairs on Jacobi bialgebroids

概要 A Dirac pair on a Jacobi bialgebroid is a pair of two Dirac structures constructing a Nijenhuis relation on a Jacobi bialgebroid and is a generalization of it on a Lie bialgebroid. In this talk, we prove a relation between Dirac pairs on Jacobi and Lie bialgebroids and show examples of Dirac pairs on Jacobi bialgebroids.

- 9 折田龍馬 (首都大東京理) Rigid fibers of spinning tops ..... 15  
 Ryuma Orita (Tokyo Metro. Univ.) Rigid fibers of spinning tops

概要 In the talk, we deal with fibers of classical Liouville integrable systems containing the Lagrangian top and the Kovalevskaya top. Especially, we find a non-displaceable fiber for each of them. To prove these results, we use the notion of superheaviness introduced by Entov and Polterovich. This is a joint work with Morimichi Kawasaki (RIMS).

- 10 佐々木 優 (筑波大数理物質) 対蹠集合の連結性と等質性 ..... 15  
 Yuuki Sasaki (Univ. of Tsukuba) Connectedness and homogeneity of antipodal sets

概要 Let  $M$  be a compact Riemannian symmetric space and denote the geodesic symmetry at  $x \in M$  by  $s_x$ . We say that  $x, y \in M$  are antipodal if  $s_x(y) = y$ . A subset  $S$  of  $M$  is an antipodal set if any two points of  $S$  are antipodal. An antipodal set  $T$  is called maximal if there are no antipodal sets including  $T$  properly. In this talk, we introduce a concept of connectedness of two antipodal points and antipodal sets. Using this connectedness, we construct a method to make a bigger antipodal set from a given antipodal set. Also, we construct a method to decide whether a given maximal antipodal set is homogeneous.

- 11 岡田 真央 (東大数理) Local rigidity of certain actions of solvable groups on the boundaries of rank-one symmetric spaces ..... 15  
 Mao Okada (Univ. of Tokyo) Local rigidity of certain actions of solvable groups on the boundaries of rank-one symmetric spaces

概要 We study local rigidity of certain actions of a solvable group on the boundary of a rank-one symmetric space of non-compact type, which is diffeomorphic to a sphere. When the symmetric space is a quaternionic hyperbolic space or the Cayley plane, the action we constructed is locally rigid.

- 12 酒井 高司 (首都大東京理) Natural  $\Gamma$ -symmetric structures on  $R$ -spaces ..... 15  
 P. Quast (Univ. of Augsburg)  
 Takashi Sakai (Tokyo Metro. Univ.) Natural  $\Gamma$ -symmetric structures on  $R$ -spaces  
 Peter Quast (Univ. of Augsburg)

概要 We classify  $R$ -spaces that admit a certain natural  $\Gamma$ -symmetric structure. We further determine the maximal antipodal sets of these structures.

#### 16:00~17:00 特別講演

- 山本 光 (東京理大理) 特殊ラグランジュ部分多様体と平均曲率流とそのミラー  
 Hikaru Yamamoto (Tokyo Univ. of Sci.) Special Lagrangian submanifolds, mean curvature flows and their mirror

概要 A special Lagrangian submanifold was defined by Harvey and Lawson in 1982 as a volume-minimizing Lagrangian submanifold in a Calabi–Yau manifold. Since Strominger, Yau and Zaslow in 1996 gave physical importance to special Lagrangian submanifolds in the context of mirror symmetry, special Lagrangian submanifolds have acquired much attention from both mathematicians and physicists. In 2002, Thomas and Yau conjectured that if a given Lagrangian submanifold is stable (in the sense of their paper) then the mean curvature flow starting from it exists for all time and converges to a special Lagrangian submanifold in its Hamiltonian deformation class. This is the so-called Thomas–Yau conjecture. In 2015, Joyce updated Thomas–Yau conjecture to make the statement more careful, and it is still widely open. Actually, this story is in A-side of mirror symmetry, and there is a corresponding story in B-side. The special object is a deformed Hermitian–Yang Mills connection and a way to get it is a line bundle mean curvature flow. In the former part of this talk, I would like to give an overview of recent development related to Thomas–Yau conjecture, and in the latter part, give that of studies in B-side which are rapidly developed in these several years.

## 3月17日(火) 第II会場

## 9:40~11:50

- 13 盧 玉 峰 (阪 大 理) Singularity theorems on Lorentz–Finsler manifolds ..... 10  
 E. Minguzzi (Univ. Stud. Firenze)  
 太 田 慎 一 (阪大理・理化学研AIP)  
 Yufeng Lu (Osaka Univ.) Singularity theorems on Lorentz–Finsler manifolds  
 Ettore Minguzzi (Univ. Stud. Firenze)  
 Shin-ichi Ohta (Osaka Univ./RIKEN)

概要 Motivated by recent works on singularity theorems by Case for weighted Lorentzian manifolds and by Minguzzi for Finsler spacetimes, we established the theory for most singularity theorems, including Penrose’s, Hawking’s and Hawking and Penrose’s, in the framework of weighted Lorentz–Finsler manifolds. Our results cover Case’s case ( $N \in [n, \infty]$ ) and Woolgar–Wylie’s case ( $N \in (-\infty, 0)$ ) as special cases.

- 14 本 田 淳 史 (横 浜 国 大 理 工) 特異値集合と第一基本形式を保つカスプ辺の双対性 ..... 15  
 直 川 耕 祐 (広 島 工 大)  
 佐 治 健 太 郎 (神 戸 大 理)  
 梅 原 雅 顕 (東 工 大 情 報 理 工)  
 山 田 光 太 郎 (東 工 大 理)  
 Atsufumi Honda Duality on generalized cuspidal edges preserving singular set images and  
 (Yokohama Nat. Univ.) first fundamental forms  
 Kosuke Naokawa  
 (Hiroshima Inst. of Tech.)  
 Kentaro Saji (Kobe Univ.)  
 Masaaki Umehara (Tokyo Tech)  
 Kotaro Yamada (Tokyo Tech)

概要 In the second, fourth and fifth authors’ previous work, a duality on generic real analytic cuspidal edges in the Euclidean 3-space preserving their singular set images and first fundamental forms, was given. Here, we call this an isometric duality. In this paper, we show that this duality extends to generalized cuspidal edges, including cuspidal cross caps, and 5/2-cuspidal edges. When the singular set image has no symmetries and does not lie in a plane, the dual cuspidal edge is not congruent to the original one.

- 15 伊 敷 喜 斗 (筑 波 大 数 理 物 質) Assouad 次元と距離空間の収束について ..... 15  
 Yoshito Ishiki (Univ. of Tsukuba) On the Assouad dimension and convergence of metric spaces

概要 We introduce the notion of pseudo-cones of metric spaces as a generalization of both of the tangent cones and the asymptotic cones. We prove that the Assouad dimension of a metric space is bounded from below by that of any pseudo-cone of it. We exhibit a example containing all compact metric spaces as pseudo-cones, and examples containing all proper length spaces as tangent cones or asymptotic cones.

- 16 伊 敷 喜 斗 (筑 波 大 数 理 物 質) 全体空間と同じ Assouad 次元を持つ部分距離空間の特徴付け ..... 15  
 Yoshito Ishiki (Univ. of Tsukuba) A characterization of metric subspaces of full Assouad dimension

概要 We introduce the notion of tiling spaces for metric space. The class of tiling spaces includes the Euclidean spaces, the middle-third Cantor spaces, and various self-similar spaces appeared in fractal geometry. On a tiling space, we characterize a metric subspace whose Assouad dimension coincides with that of the whole space.

- 17 印南信宏(新潟大理) Finsler 多様体のための正距方位図法 ..... 15  
 Nobuhiro Innami (Niigata Univ.) The azimuthal equidistant projection for a Finsler manifold

概要 Let  $(M, F)$  be a geodesically forward complete Finsler manifold and  $p \in M$ . We observe how the preimage of a curve in  $M$  under exponential map at  $p$  can behave in the tangent space  $T_p M$  at  $p$ , when the curve approaches a conjugate cut point of  $p$  without crossing the cut locus of  $p$ .

- 18 五明工(名大多元数理) 有限グラフの埋め込み不変量の最小化と第1固有値の最大化 ..... 15  
 小林俊公(摂南大理工)  
 近藤剛史(鹿児島大理)  
 納谷信(名大多元数理)

Takumi Gomyou (Nagoya Univ.) Optimal embedding and spectral gap of a finite graph  
 Toshimasa Kobayashi (Setsunan Univ.)  
 Takefumi Kondo (Kagoshima Univ.)  
 Shin Nayatani (Nagoya Univ.)

概要 We introduce an embedding optimization problem for a finite graph. This problem is related to an optimization problem concerning the smallest nonzero eigenvalue of the graph Laplacian. Goering–Helmberg–Wappler introduced a different embedding problem as a dual of the eigenvalue optimization problem. We establish a relation between the optimal values of the two embedding problems. It then follows that the optimal value of our embedding problem is obtained by the optimal value of the eigenvalue optimization problem. Further, we show that our embedding problem is also a dual of the eigenvalue optimization problem. We present examples of graphs for which these optimization problems can be explicitly solved.

- 19 佐藤直飛(北大理) 統計断面曲率とワーブ積統計多様体 ..... 15  
 古畑仁(北大理)  
 長谷川和泉(北教大\*)

Naoto Satoh (Hokkaido Univ.) Statistical sectional curvature and warped product statistical manifold  
 Hitoshi Furuhashi (Hokkaido Univ.)  
 Izumi Hasegawa  
 (Hokkaido Univ. of Edu.\*)

概要 Sectional curvature is one of the most important notions in differential geometry. Statistical sectional curvature can be introduced on a statistical manifold by using the statistical curvature tensor field, which is defined as the arithmetic mean of curvature tensor fields with respect to the pair of dual connections. For a submanifold in a statistical manifold, the Euler inequality, which presents a relation between the mean curvature and the Gaussian curvature in Euclidean geometry, is generalized in terms of the statistical sectional curvature. A doubly umbilical statistical submanifold is characterized as a submanifold such that the equality holds at each point. In particular, we describe the inequality in the case where the ambient space is a standard warped product statistical manifold.

- 20 小澤 龍ノ介 (東北大 AIMR) リッチ曲率が下に有界な有向グラフの幾何解析的性質 ..... 15  
 櫻井 陽平 (東北大 AIMR)  
 山田 大貴 (総合地球環境学研)  
 Ryunosuke Ozawa (Tohoku Univ.) Geometric and analytic properties of directed graphs under lower Ricci  
 Yohei Sakurai (Tohoku Univ.) curvature bound  
 Taiki Yamada  
 (Res. Inst. for Humanity and Nature)

概要 For undirected graphs, the Ricci curvature introduced by Lin–Lu–Yau has been widely studied from various perspectives, especially geometric analysis. In this talk, we discuss generalization problem of their Ricci curvature for directed graphs. We introduce a new generalization by using the mean transition probability kernel which appears in the formulation of the Chung Laplacian. We conclude several geometric and spectral properties of directed graphs under lower Ricci curvature bound extending previous results in the undirected case.

### 13:15~14:15 特別講演

- U. Pinkall ( TU Berlin ) Discretizing fluids into filaments and sheets  
 Ulrich Pinkall (TU Berlin) Discretizing fluids into filaments and sheets

概要 In differential geometry and in mathematical physics “discretization” means replacing a continuum system (like for example a smooth surface in three-space) by a discrete collection of lower-dimensional objects (like curves or even isolated points). It is often possible to capture many important properties of the original continuum system in its discrete equivalent. In many situations relevant for physics simulation the relevant fields (vorticity of a fluid, current or magnetic field in a plasma) exhibit on their own a strong tendency to become concentrated in one-dimensional filaments or two-dimensional sheets. “Decomposing” a given continuous field into its “constituent” filaments or sheets is important for understanding and visualizing a given field. Moreover, simulation algorithms that take into account the natural tendency of the fields to become ”discretized” into filaments or sheets can be highly efficient.

3月18日(水) 第II会場

### 9:40~11:35

- 21 三石 史人 (福岡大理)  $p$  エネルギーのある種のミニ・マックス値とパッキング半径 ..... 15  
 Ayato Mitsuishi (Fukuoka Univ.) Certain mini-max values of the  $p$ -energy and packing radii

概要 Grosjean proved that the  $(1/p)$ -th power of the first non-zero eigenvalue of the  $p$ -Laplacian on a closed Riemannian manifold converges to the twice of the inverse of the diameter of the space, when  $p$  goes to infinity. Before this being proved, Juutinen, Lindqvist and Manfredi proved a corresponding result for the Dirichlet eigenvalues. The speaker extends their results to “ $k$ -th values”. More precisely, we consider “ $k$ -th diameter” as the  $(k+1)$ -th packing radius introduced by Grove–Markvorsen. Furthermore, we introduce certain min-max value related to the  $p$ -energy as a counter-part of “ $k$ -th eigenvalue.” Here, please note that our min-max values are not known whether they are real eigenvalues.

- 22 Cong Hung Mai (京大 理) Quantitative estimates for the Bakry–Ledoux isoperimetric inequality  
 太田 慎一 (阪大理・理化学研AIP) ..... 10  
Cong Hung Mai (Kyoto Univ.) Quantitative estimates for the Bakry–Ledoux isoperimetric inequality  
 Shin-ichi Ohta (Osaka Univ./RIKEN)

概要 We establish a quantitative isoperimetric inequality for weighted Riemannian manifolds with lower weighted Ricci curvature bound. Precisely, we give an upper bound of the volume of the symmetric difference between a Borel set and a sub-level (or super-level) set of the associated guiding function (arising from the needle decomposition), in terms of the deficit in Bakry–Ledoux’s Gaussian isoperimetric inequality. This is the first quantitative isoperimetric inequality on noncompact spaces besides Euclidean and Gaussian spaces. Our argument makes use of Klartag’s needle decomposition (also called localization), and is inspired by the recent work of Cavalletti, Maggi and Mondino on compact spaces.

- 23 赤嶺新太郎 (名大多元数理) 光的点を許容する平均曲率零超曲面に対する Bernstein 型定理 ..... 15  
 本田 淳史 (横浜国大理工)  
 梅原 雅顕 (東工大情報理工)  
 山田光太郎 (東工大 理)  
Shintaro Akamine (Nagoya Univ.) Bernstein-type theorem for zero mean curvature hypersurfaces admit-  
 Atsufumi Honda ting lightlike points  
 (Yokohama Nat. Univ.)  
 Masaaki Umehara (Tokyo Tech)  
 Kotaro Yamada (Tokyo Tech)

概要 Calabi and Cheng–Yau’s Bernstein-type theorem asserts that an entire zero mean curvature graph in Lorentz–Minkowski  $(n+1)$ -space which admits only space-like points is a hyperplane. We give an improvement of the Bernstein-type theorem and show that an entire zero mean curvature graph in consisting only of space-like or light-like points is a hyperplane.

- 24 高田 土満 (東大 数理) Atiyah–Singer の指数定理の無限次元化に向けて ..... 15  
 Doman Takata (Univ. of Tokyo) Towards an infinite-dimensional Atiyah–Singer index theorem

概要 The Atiyah–Singer index theorem states, on a closed manifold, that the analytic index of the Dirac operator, is determined by topological data. Many mathematicians have generalized this theorem, for example Kasparov proved the version of non-compact manifolds equipped with proper and cocompact group actions, in terms of non-commutative geometry. The overall goal of my research is to formulate and prove an infinite-dimensional version of this index theorem. In other words, I would like to give an equivariant index theorem for infinite-dimensional manifolds equipped with proper and cocompact actions of infinite-dimensional Lie groups, in terms of non-commutative geometry. In this talk, I will explain the results so far, focusing on the non-commutative geometrical side. Time permitting, I will mention my ongoing work which connects the results so far and some topological invariants.

- 25 今井 淳 (千葉大 理) 留数としての Willmore エネルギーの拡張 ..... 15  
 Jun O’Hara (Chiba Univ.) Generalization of Willmore energy as a residue

概要 We give generalization of the Willmore energy for an even-dimensional closed submanifold in the Euclidean space as a residue of a meromorphic function obtained from the Riesz energy. We show that it is independent from the Graham–Witten energy when the dimension is 4.

- 26 高津飛鳥 (首都大東京理)<sup>b</sup> 回転対称距離球上の楕円型・放物型方程式の解の凹性 ..... 15  
 石毛和弘 (東大数理)  
 P. Salani (Univ. Firenze)  
 Asuka Takatsu (Tokyo Metro. Univ.) Elliptic and parabolic boundary value problems on rotationally sym-  
 Kazuhiro Ishige (Univ. of Tokyo) metric domains  
 Paolo Salani (Univ. Firenze)

概要 We study power concavity of solutions to elliptic and parabolic boundary value problems on rotationally symmetric, strongly convex open metric balls in a Riemannian manifold. Our results provide a first step to the study of power concavity for Riemannian manifolds, and improve the known results for Euclidean spaces.

- 27 佐々木東容 (早大理工) カस्प付き双曲曲面上の測地カレントの稠密性問題 ..... 15  
 Dounnu Sasaki (Waseda Univ.) Denseness property of geodesic currents on a cusped hyperbolic surface

概要 The space of geodesic currents on a hyperbolic surface were introduced by Bonahon as a generalization of measured geodesic laminations and have been successfully studied in the case that the surface is closed or compact (possibly with boundary). One of useful properties is that the space of geodesic currents on a closed hyperbolic surface can be considered as the measure-theoretic completion of the set of weighted closed geodesics on the surface, but such a property is not shown in the case that the surface has cusps due to infinite geodesics connecting two cusps on the surface. We have proved that the space of geodesic currents on a cusped hyperbolic surface with finite area also has the same denseness property.

#### 14:20~15:40

- 28 藤岡 敦 (関西大システム理工) 余等質性 1 の中心アファイン曲面 ..... 15  
 古畑 仁 (北大理工)  
 Atsushi Fujioka (Kansai Univ.) Centroaffine surfaces of cohomogeneity one  
 Hitoshi Furuhashi (Hokkaido Univ.)

概要 A nondegenerate centroaffine surface of cohomogeneity one can be considered as a corresponding object to a surface of revolution in Euclidean geometry. We obtain a normal form for such a surface to classify such centroaffine minimal surfaces with centroaffine metrics of constant curvature. We also study proper affine spheres from this viewpoint.

- 29 安本真士 (阪市大数学研) 離散 Weierstrass 型の表現公式 ..... 15  
 Masashi Yasumoto (Osaka City Univ.) Discrete Weierstrass-type representations

概要 Over the last three decades, structure-preserving discretizations of differential geometry are rapidly developing. In particular, discretizations of surfaces are highly related to differential geometry itself, integrable systems, complex analysis, discrete differential geometry, and so on. Bobenko–Pinkall showed that there exists a Weierstrass-type representation for discrete minimal surfaces in terms of discrete holomorphic functions. Afterwards other Weierstrass-type representations were discovered. In this talk we introduce Weierstrass representations for discrete surfaces. A common feature of Weierstrass-type representations for discrete surfaces is that they are described by discrete holomorphic functions. We give a unified description for these representation formulae in terms of transformations for Omega-surfaces. This talk is based on joint work with Mason Pember (Politecnico di Torino) and Denis Polly (Technische Universität Wien).

- 30 森本真弘(阪市大理) ヒルベルト空間の PF 部分多様体のオースティア性とアリッド性 ..... 15  
Masahiro Morimoto (Osaka City Univ.) Austere and arid properties for PF submanifolds in Hilbert spaces

概要 Austere submanifolds and arid submanifolds constitute respectively two different classes of minimal submanifolds in finite dimensional Riemannian manifolds. In this talk we introduce these two notions into a class of proper Fredholm (PF) submanifolds in Hilbert spaces, discuss their relation and show many examples of infinite dimensional austere PF submanifolds and arid PF submanifolds in Hilbert spaces.

- 31 佐藤雄一郎(首都大東京理) 擬 Riemann 空間形内の全臍的部分多様体 ..... 15  
Yuichiro Sato (Tokyo Metro. Univ.) Totally umbilical submanifolds in pseudo-Riemannian space form

概要 Totally umbilical submanifolds in pseudo-Riemannian manifolds are defined by the traceless part of the second fundamental form vanishing identically. They are fundamental submanifolds next to totally geodesic submanifolds. In this talk, we classify the congruent class of full totally umbilical submanifolds in non-flat pseudo-Riemannian space forms, and consider its moduli spaces. As a consequence, we obtain that some moduli spaces of isometric immersions between space forms which are of the same constant curvature are non-Hausdorff.

- 32 奥村和浩(旭川工高専) 非平坦複素空間形内の線織実超曲面の曲率テンソルについて ..... 10  
Kazuhiro Okumura The curvature tensor of ruled real hypersurfaces in a nonflat complex  
(Asahikawa Nat. Coll. of Tech.) space form

概要 In this talk, we consider ruled real hypersurfaces in a nonflat complex space form from the viewpoint of  $\phi$ -invariances of curvature tensors of real hypersurfaces. Furthermore we give a new characterization of these real hypersurfaces.

#### 16:00~17:00 特別講演

- 金沢篤(京大理) Kähler モジュライ空間と三角圏の安定性条件  
Atsushi Kanazawa (Kyoto Univ.) Kähler moduli spaces and stability conditions of triangulated categories

概要 For a Kähler manifold, we can consider two moduli spaces, namely the complex moduli space and the Kähler moduli space. The former has been one of the most important subjects in modern mathematics (studied by Riemann, Kodaira–Spencer, Kuranishi and others), and beautiful theories have been developed in a variety of cases. In light of mirror symmetry, we expect that there are equivalently rich theories for the Kähler moduli spaces. However, it turns out that the Kähler structures are much more delicate than the complex structures. In this talk, I will introduce our program to investigate the Kähler moduli spaces via stability conditions of triangulated categories. Roughly speaking, we will consider a Kähler analogue of the Hodge theory.



# 函 数 論

3月16日(月) 第VIII会場

9:30~11:50

- 1 齋藤三郎 (群馬大\*再生核研) Okumura's disc series can beyond the crucial point of Däumler–Puha's horn torus models for the Riemann sphere ..... 15  
 Saburou Saitoh (Gunma Univ.\*/Inst. of Reproducing Kernels) Okumura's disc series can beyond the crucial point of Däumler–Puha's horn torus models for the Riemann sphere

概要 For the statement of the conclusion, we will first recall the basic related backgrounds; division by zero calculus, examples and horn torus models.

- 2 須川敏幸 (東北大情報) 単位円板の自己正則写像からなる連続半群のレゾルベントの幾何的性質  
 M. Elin (ORT Braude Coll.) について ..... 15  
 D. Shoikhet (Holon Inst. of Tech.)  
 Toshiyuki Sugawa (Tohoku Univ.) Geometric properties of the nonlinear resolvent for a continuous semi-  
 Mark Elin (ORT Braude Coll.) group of holomorphic self-maps of the unit disk  
 David Shoikhet (Holon Inst. of Tech.)

概要 Let  $f$  be the infinitesimal generator of a one-parameter semigroup of holomorphic self-mappings of the open unit disk  $\mathbb{D}$ . Our main purpose is to study properties of the family  $\mathcal{R}$  of nonlinear resolvents  $(I + rf)^{-1} : \mathbb{D} \rightarrow \mathbb{D}$ ,  $r \geq 0$ , in the spirit of classical geometric function theory. To make a connection with this theory, we mostly consider the case where  $f(0) = 0$  and  $f'(0) > 0$ . We found, in particular, that  $\mathcal{R}$  forms an inverse Löwner chain of hyperbolically convex functions. Moreover, each element of  $\mathcal{R}$  satisfies the Noshiro–Warschawskii condition. This, in turn, implies that each element of  $\mathcal{R}$  is also an infinitesimal generator of a one-parameter semigroup on  $\mathbb{D}$ . We mention also quasiconformal extension of elements of  $\mathcal{R}$ .

- 3 泉英明 (千葉工大情報) 階関数方程式の拡張次元数解 ..... 15  
 Hideaki Izumi (Chiba Inst. of Tech.) Dimensioned number solutions to iterative functional equations

概要 We will obtain formal solutions to iterative functional equation  $f(f(x)) = x$  by using extended dimensioned numbers.

- 4 牛島 顕 (金沢大理工) 有限余面積フックス群に対する exceptional な基点の存在について ..... 15  
 中西敏浩 (島根大総合理工)  
 Akira Ushijima (Kanazawa Univ.) Existence of exceptional points for cofinite Fuchsian groups  
 Toshihiro Nakanishi (Shimane Univ.)

概要 It is shown by Fera that there exist uncountably many exceptional points for cocompact Fuchsian groups. We generalize this result to the case that the Fuchsian group is cofinite.

- 5 渡邊天鵬 (京大人間環境) 有理写像からなるマルコフ的ランダム力学系のダイコトミー ..... 15  
 角 大輝 (京大人間環境)  
 Takayuki Watanabe (Kyoto Univ.) Dichotomy of Markov random dynamical systems of rational maps  
 Hiroki Sumi (Kyoto Univ.)

概要 We consider random holomorphic dynamical systems on the Riemann sphere whose choices of maps are related to “Markovian” noise. Our motivation is generalizing the theory of i.i.d. random dynamical systems to our setting. We show that a generic such system is either stable on average or chaotic with full Julia set. We also talk about difference between i.i.d. and non-i.i.d. random dynamical systems.

- 6 木坂正史 (京大人間環境) Fatou–Shishikura inequality for transcendental entire functions in class  $\mathcal{S}$  ..... 15  
 Masashi Kisaka (Kyoto Univ.) Fatou–Shishikura inequality for transcendental entire functions in class  $\mathcal{S}$

概要 We discuss the following realizability problem: For given numbers which satisfy the Fatou–Shishikura inequality, is there a transcendental entire functions in class  $\mathcal{S}$  with the given numbers of Fatou components?

- 7 神本 丈 (九大数理) 局所ゼータ関数の有理型解析接続可能領域について ..... 15  
 野瀬 敏洋 (福岡工大)  
 Joe Kamimoto (Kyushu Univ.) On the maximal region to which local zeta functions can be meromorphically extended  
 Toshihiro Nose (Fukuoka Inst. of Tech.)

概要 It is known that local zeta functions associated with real analytic functions can be analytically continued as meromorphic functions to the whole complex plane. But, in the case of general smooth functions, it has been recently shown that there exist specific smooth functions whose local zeta functions have singularities different from poles. In order to understand the situation of the meromorphic extension in the smooth case, we investigate a simple but essentially important case, in which the respective function is expressed as  $u(x, y)x^a y^b + \text{flat function}$ , where  $u(0, 0) \neq 0$  and  $a, b$  are nonnegative integers.

- 8 野瀬 敏洋 (福岡工大) 局所ゼータ関数の極性をもたない特異性について ..... 15  
 神本 丈 (九大数理)  
 Toshihiro Nose (Fukuoka Inst. of Tech.) On non-polar singularities of local zeta functions  
 Joe Kamimoto (Kyushu Univ.)

概要 In this talk, we investigate asymptotic limits of local zeta functions associated with specific (non-real analytic) smooth functions at some singularities along one direction. It follows from these behaviors that these local zeta functions have singularities different from poles, which gives the optimality of the lower estimates of an invariant concerning with meromorphic continuation of local zeta functions in the case of smooth functions of the form investigated in the previous talk.

#### 14:15~15:20

- 9 綾野 孝則 (阪市大数学研) 熱方程式に基づく 2 変数 sigma 関数の級数展開 ..... 15  
 V. M. Buchstaber  
 (Steklov Inst. of Math.)  
 Takanori Ayano (Osaka City Univ.) Series expansion of two-dimensional sigma function based on the heat equations  
 Victor M. Buchstaber  
 (Steklov Inst. of Math.)

概要 Weierstrass gave the heat equations which characterize the elliptic sigma function. Buchstaber and Leykin succeeded in generalizing the theory of the heat equations to the sigma functions of higher genus curves. Eilbeck, Gibbons, Onishi, and Yasuda gave the detailed proof of their theory. They also gave the recurrence relations of the coefficients of series expansion of the two-dimensional sigma function. In this talk, we derive new recurrence relations of series expansion of the two-dimensional sigma function based on the heat equations. Our recurrence relations take the form in which all the parameters of the curve are included. As a corollary, we can find that the two-dimensional sigma function is characterized by a part of the heat equations.

- 10 林本厚志 (長野工高専) Hua 領域の自己同型群と等方性群 ..... 15  
 Atsushi Hayashimoto Automorphism group and isometry group of Hua domains  
 (Nagano Nat. Coll. of Tech.)

概要 For proper holomorphic polynomial mappings between Hua domains, we show that two such mappings are isotropic equivalence if and only if they are automorphic equivalence.

- 11 濱田英隆 (九州産大理工) Distortion theorems, Lipschitz continuity and their applications for Bloch type mappings on bounded symmetric domains in  $\mathbb{C}^n$  ..... 15  
 Hidetaka Hamada Distortion theorems, Lipschitz continuity and their applications for Bloch type mappings on bounded symmetric domains in  $\mathbb{C}^n$   
 (Kyushu Sangyo Univ.)

概要 Let  $\mathbb{B}_X$  be a bounded symmetric domain realized as the unit ball of an  $n$ -dimensional  $JB^*$ -triple  $X = (\mathbb{C}^n, \|\cdot\|_X)$ . In this talk, we give a new definition of Bloch type mappings on  $\mathbb{B}_X$  and give distortion theorems for Bloch type mappings on  $\mathbb{B}_X$ . As a corollary of the distortion theorem, we obtain the lower estimate for the radius of the largest schlicht ball in the image of  $f$  centered at  $f(0)$  for  $\alpha$ -Bloch mappings  $f$  on  $\mathbb{B}_X$ . Next, as another corollary of the distortion theorem, we show the Lipschitz continuity of  $(\det B(z, z))^{1/2n} |\det Df(z)|^{1/n}$  for Bloch type mappings  $f$  on  $\mathbb{B}_X$  with respect to the Kobayashi metric, where  $B(z, z)$  is the Bergman operator on  $X$ , and use it to give a sufficient condition for the composition operator  $C_\varphi$  to be bounded from below on the Bloch type space on  $\mathbb{B}_X$ , where  $\varphi$  is a holomorphic self mapping of  $\mathbb{B}_X$ .

- 12 I. Graham (Univ. of Toronto) Loewner chains, Bloch mappings and Pfaltzgraff–Suffridge extension operators on bounded symmetric domains ..... 15  
 濱田英隆 (九州産大理工)  
 G. Kohr (Babeş-Bolyai Univ.)  
 Ian Graham (Univ. of Toronto) Loewner chains, Bloch mappings and Pfaltzgraff–Suffridge extension operators on bounded symmetric domains  
 Hidetaka Hamada  
 (Kyushu Sangyo Univ.)  
 Gabriela Kohr (Babeş-Bolyai Univ.)

概要 Let  $Y$  be a complex Banach space and let  $\mathbb{B}_Y$  be the open unit ball of  $Y$ . In this talk, we consider a generalization of the Pfaltzgraff–Suffridge extension operator on bounded symmetric domains  $\mathbb{B}_X$  in  $\mathbb{C}^n$ , and show that this operator extends the first elements of Loewner chains from  $\mathbb{B}_X$  to the first elements of Loewner chains on  $\mathbb{D}_\alpha$ , when  $\alpha \geq n/(2c(\mathbb{B}_X))$ , where  $\mathbb{D}_\alpha \subset \mathbb{B}_X \times \mathbb{B}_Y$  is a certain domain with  $\mathbb{B}_X \times \{0\} \subset \mathbb{D}_\alpha$  and  $c(\mathbb{B}_X)$  is a constant defined by the Bergman metric on  $X$ . Next, we also show that normalized locally univalent I-Bloch mappings, which have finite trace order on  $\mathbb{B}_X$ , are mapped into R-Bloch mappings on  $\Omega_\alpha$  by this extension operator when  $\alpha \geq 1/2$ , where  $\Omega_\alpha \subset X \times Y$  is a bounded balanced convex domain such that  $\mathbb{B}_X \times \{0\} \subset \Omega_\alpha \subset \mathbb{D}_\alpha$ .

### 15:35~16:35 特別講演

- 小森洋平 (早大教育) 双曲コクセター群の増大度  
 Yohei Komori (Waseda Univ.) Growth of hyperbolic Coxeter groups

概要 In this talk I will give an overview of recent progress on arithmetic aspects of growth related to hyperbolic Coxeter groups.

**16:50~17:50 2019年度(第18回)日本数学会解析学賞受賞特別講演**

角 大 輝 (京大人間環境) ランダム複素力学系における様々なランダム性誘起現象とそのメカニズム  
 Hiroki Sumi (Kyoto Univ.) Various randomness-induced phenomena and their mechanisms in random holomorphic dynamical systems

概要 We consider random holomorphic dynamical systems. In particular, we see various randomness-induced phenomena in random holomorphic dynamical systems which cannot hold in the deterministic random holomorphic dynamical systems.

3月17日(火) 第VIII会場

**9:15~11:45**

- 13 田 島 慎 一 (新 潟 大\*) 孤立特異点を持つ complete intersection に沿う対数的ベクトル場の構成  
 渋 田 敬 史 (九州産大理工) 法 ..... 15  
 鍋 島 克 輔 (徳 島 大 理 工)  
 Shinichi Tajima (Niigata Univ.\*) A computation method of logarithmic vector fields associated to isolated  
 Takafumi Shibuta complete intersection singularities  
 (Kyushu Sangyo Univ.)  
 Katsusuke Nabeshima  
 (Tokushima Univ.)

概要 Logarithmic vector fields of a complete intersection are important objects with many applications. However, its computation is hard, as a number of open questions and challenges indicate. In this talk logarithmic vector fields associated to isolated complete intersection singularities are considered. A computation method for computing the logarithmic vector fields is introduced.

- 14 篠 原 知 子 (産業技術高専) 複素2変数有理写像の不定点の局所安定集合 ..... 15  
 Tomoko Shinohara Local stable set of an indeterminate point of rational mappings of two  
 (Tokyo Metro. Coll. of Ind. Tech.) complex variables

概要 In this talk, we study local dynamics of a rational mapping of two complex variables at an indeterminate point. It is known that for some Newton's method of two variables there exist a Cantor bouquet and a local stable set at an indeterminate point. We define the order of a rational mapping at an indeterminate point and give a sufficient condition that the rational mapping becomes to be a horizontal-like mapping. By using this order, we construct a local stable set of an indeterminate point for general rational mappings.

- 15 阿 部 幸 隆 (富 山 大 理 工) Isogenies between commutative complex Lie groups ..... 10  
 Yukitaka Abe (Univ. of Toyama) Isogenies between commutative complex Lie groups

概要 We study isogenies between connected commutative complex Lie groups.

- 16 阿 部 幸 隆 (富 山 大 理 工) Meromorphic function fields closed by partial derivatives ..... 15  
 Yukitaka Abe (Univ. of Toyama) Meromorphic function fields closed by partial derivatives

概要 We characterize meromorphic function fields closed by partial derivatives in  $n$  variables.

- 17 小池 貴之 (阪市大理) Hermitian metrics on the anti-canonical bundle of the blow-up of the projective plane at nine points ..... 15  
 Takayuki Koike (Osaka City Univ.) Hermitian metrics on the anti-canonical bundle of the blow-up of the projective plane at nine points

概要 We investigate Hermitian metrics on the anti-canonical bundle of a rational surface obtained by blowing up the projective plane at nine points. For that purpose, we pose a modified variant of an argument made by Ueda on the complex analytic structure of a neighborhood of a subvariety by considering the deformation of the complex structure.

- 18 足立 真訓 (静岡大理) 複素多様体内の弱擬凸領域に対する Diederich–Fornæss 指数と Stein 性  
 Jihun Yum (Pusan Nat. Univ.) 指数 ..... 15  
 Masanori Adachi (Shizuoka Univ.) The Diederich–Fornæss and Steinness indices in complex manifolds  
 Jihun Yum (Pusan Nat. Univ.)

概要 The Diederich–Fornæss and Steinness indices are estimated for weakly pseudoconvex domains in complex manifolds by using the D’Angelo 1-form of the boundary. In particular, CR invariance of these indices is shown when the domain is Takeuchi 1-convex.

- 19 稲山 貴大 (東大数理) 多重標準束の順像層における擬ノルムについて ..... 15  
 Takahiro Inayama (Univ. of Tokyo) Pseudonorms on direct images of pluricanonical bundles

概要 We investigate Stein morphisms and the pseudonorms on direct images of pluricanonical bundles. Our main goal in this talk is to show that the pseudonorms determine holomorphic structures of Stein morphisms. One important technique is an  $L^{2/m}$  variant of the Ohsawa–Takegoshi extension theorem.

- 20 細野 元気 (東北大理) 最良係数の  $L^2$  拡張定理の証明の簡略化とその応用 ..... 15  
 Genki Hosono (Tohoku Univ.) A simplified proof of the optimal  $L^2$  extension theorem and its application

概要 I will give a simplified proof of an optimal version of the Ohsawa–Takegoshi  $L^2$ -extension theorem. In the proof, I use a method of Berndtsson–Lempert and skip some argument by the method of McNeal–Varolin. As an application, I will explain a result on extensions from possibly non-reduced varieties.

- 21 大沢 健夫 (名大多元数理)<sup>b</sup> Application of the  $L^2$  method to the Levi problem on complex manifolds ..... 15  
 Takeo Ohsawa (Nagoya Univ.) Application of the  $L^2$  method to the Levi problem on complex manifolds

概要  $L^2$  methods for extending holomorphic sections of semipositive bundles are applied to show the holomorphic convexity of complex spaces under several conditions.

**13:15~14:15 特別講演**

松村 慎一 (東北大理)<sup>b</sup> “非負曲率”を持つ射影多様体に対する構造定理について

Shin-ichi Matsumura (Tohoku Univ.) On structure theorems for projective manifolds with certain non-negative curvature

**概要** The study of certain positively curved varieties, which are often formulated with positivity of bisectional curvatures, tangent bundles, or anti-canonical divisors, occupies an important place in the theory of classification of varieties. One of the central problems in this study is to determine the structure of semi-positively curved varieties in terms of naturally associated fibrations such as Albanese maps, Iitaka fibrations, or maximally rationally connected fibrations. In this talk, I would like to discuss structure theorems for projective varieties (more generally compact Kähler manifolds) satisfying various positivity conditions. More specifically, I will explain structure theorems of maximally rationally connected fibrations for varieties satisfying the following conditions: (1) Projective manifolds with tangent bundle admitting positively curved singular hermitian metrics. (2) Projective manifolds with semi-positive holomorphic sectional curvature. (3) Projective KLT pairs with nef anti-canonical bundle. The proof is based on singular hermitian metrics on vector bundles, analytic positivity of direct image sheaves, the theory of foliations, and so on. A part of this talk is a joint work with F. Campana (Lorraine), J. Cao (Jussieu), M. Iwai (Tokyo), G. Hosono (Tohoku).

# 函数方程式論

3月16日(月) 第Ⅲ会場

9:00~12:00

- 1 井上 公人 (九大 I M I) 行列値 Bratu 方程式の指数行列解とベキ級数解 ..... 10  
 Hiroto Inoue (Kyushu Univ.) The exponential matrix solution and power series solution of the matrix-valued Bratu equation

概要 The Bratu equation is a nonlinear ordinary differential equation arising originally from the Frank–Kamenetskii’s combustion theory. In a mathematical contexts, the Bratu equation appears as the simplest case of the finite Toda lattice, whose an elaborate theory for the initial value problem exists. It also appears as the radial part of the 2-dimensional Liouville equation. The boundary value problem of the Liouville equation is called the Liouville–Bratu–Gelfand problem and its interesting solution structures are widely studied. In this talk, we define a matrix-valued extension of the Bratu equation and give its power series solution.

- 2 小森 大地 (北大 理) Čech–Dolbeault cohomology を用いた無限階擬微分作用素からその表象への射の構成 ..... 10  
 Daichi Komori (Hokkaido Univ.) The construction of the morphism of sheaves from pseudodifferential operators to their symbols via Čech–Dolbeault cohomology

概要 For the study of the sheaf  $\mathcal{E}_X^{\mathbb{R}}$  of pseudodifferential operators, Kataoka and Aoki established the symbol theory  $\mathfrak{S}/\mathfrak{N}$  of  $\mathcal{E}_X^{\mathbb{R}}$  by the aid of the Radon transformations. They constructed the isomorphism  $\sigma : \mathcal{E}_{X, z^*}^{\mathbb{R}} \rightarrow \mathfrak{S}_{z^*}/\mathfrak{N}_{z^*}$  of stalks. However, they hadn’t shown the equivalence of  $\mathcal{E}_X^{\mathbb{R}}$  and its symbol class, since the morphism of stalks constructed by them cannot be directly extended to the one of sheaves. In this talk we construct the morphism of sheaves from  $\mathcal{E}_X^{\mathbb{R}}$  to  $\mathfrak{S}/\mathfrak{N}$ , which is an isomorphism of sheaves, by using the theory of Čech–Dolbeault cohomology introduced by Honda, Izawa and Suwa.

- 3 鈴木 正俊 (東工大 理) ある偏微分方程式系と Hermite–Biehler class の整関数 ..... 10  
 Masatoshi Suzuki (Tokyo Tech) On a system of partial differential equations and entire functions of Hermite–Biehler class

概要 We present an initial value problem for a system of first-order partial differential equations in two variables and explain the relationship with entire functions of Hermite–Biehler class.

- 4 田原 秀敏 (上智大 理工) 特異点をもつ一階非線型偏微分方程式の解の一意性について ..... 10  
 Hidetoshi Tahara (Sophia Univ.) Uniqueness of the solution of nonlinear singular first order partial differential equations

概要 In this talk, I will consider a class of nonlinear singular first order partial differential equations, and show the uniqueness of the solution under a very weak assumption.

- 5 只野 之英 (東大 数理) 正方格子上の離散シュレディンガー作用素の連続極限について ..... 10  
 中村 周 (学習院大 理)  
 Yukihide Tadano (Univ. of Tokyo) On a continuum limit of discrete Schrödinger operators on square lattices  
 Shu Nakamura (Gakushuin Univ.)

概要 We consider a continuum limit problem of Schrödinger operators. The motivation comes from the two questions: How to formulate the continuum limit problem from the view point of spectral theory, and whether the formulated problem is solved. In this talk, we obtain positive answers to both questions, and we see that the continuum limit holds for a large class of potentials including uniformly continuous bounded potentials and the harmonic potentials.

- 6 伊藤 健一 (東大数理) Hypergeometric expression for resolvent of the discrete Laplacian in low  
A. Jensen (Aalborg Univ.) dimension ..... 10  
Kenichi Ito (Univ. of Tokyo) Hypergeometric expression for resolvent of the discrete Laplacian in low  
Arne Jensen (Aalborg Univ.) dimension

概要 We discuss an explicit formula for resolvent of the discrete Laplacian on the square lattice, and compute its asymptotic expansions around thresholds in low dimensions. We obtain an expression of the resolvent in a general dimension employing the Appell–Lauricella hypergeometric function of type  $C$  outside a disk encircling the spectrum. In low dimensions it reduces to a generalized hypergeometric function, for which certain transformation formulas are available.

- 7 朴 佳南 (神戸大理) ある拡張された  $q$ - $P_{VI}$  系と対称性 ..... 10  
Kanam Park (Kobe Univ.) A certain generalization of  $q$ -Painlevé VI system and its symmetry

概要 We define a monodromy preserving deformation which has a special solution in terms of a generalization of  $q$ -hypergeometric functions. Also, we obtain its affine Weyl group symmetry. In this talk, we introduce the monodromy preserving deformation and give its symmetry.

- 8 高橋 甫宗 (近畿大理工) On the WKB theoretic transformation to the boosted Airy equation  
..... 10  
Toshinori Takahashi (Kindai Univ.) On the WKB theoretic transformation to the boosted Airy equation

概要 In this talk, we consider the WKB theoretic transformation of a certain Schrödinger equation to the boosted Airy equation. As a result, we find that the transformation series can be constructed and that the relation between their WKB solutions formally holds.

- 9 青木 貴史 (近畿大理工) Voros coefficients at the origin and at the infinity of the generalized  
内田 匠風 (近畿大総合理工) hypergeometric differential equation with a large parameter ..... 10  
Takashi Aoki (Kindai Univ.) Voros coefficients at the origin and at the infinity of the generalized  
Shofu Uchida (Kindai Univ.) hypergeometric differential equation with a large parameter

概要 The Voros coefficients at the origin and at the infinity are defined and their explicit forms are given for the generalized hypergeometric differential equation of  ${}_pF_q$  with a large parameter.

- 10 山根 英司 (関西学院大理工)  $\mu$ -Camassa–Holm 方程式に関する初期値問題の実解析的時間大域解 ... 10  
Hideshi Yamane Analytic global-in-time solutions to the Cauchy problem for the  $\mu$ -  
(Kwansei Gakuin Univ.) Camassa–Holm equation

概要 We solve the Cauchy problem for the  $\mu$ -Camassa–Holm integro-partial differential equation introduced by Khesin–Lenells–Misiolek in the analytic category. We show the existence of a unique global-in-time analytic solution.



- 11 石橋和葵 (広島商船高専) 半分線形差分方程式に対する Moore 型非振動定理 ..... 10  
 Fentao Wu (東北師範大)  
 Lin She  
 Kazuki Ishibashi (Hiroshima Nat. Coll. of Maritime Tech.) Moore-type nonoscillation theorems for half-linear difference equations  
 Fentao Wu (Northeast Normal Univ.)  
 Lin She

概要 In this talk, we consider the half-linear difference equation  $\Delta(r_n \Phi_p(\Delta x_n)) + c_n \Phi_p(x_{n+1}) = 0$ , where  $r_n, c_n$  are real-valued sequences,  $r_n > 0$  for  $n \in \mathbb{N} \cup \{0\}$ , and  $\Phi_p(x) = |x|^{p-2}x$  with  $p > 1$  and  $\mathbb{N}$  is the set of natural numbers. The purpose of this talk is to give new criteria which guarantee that all non-trivial solutions of the half-linear difference equation are nonoscillatory. We obtain the desired results by using the method of Riccati technique.

- 12 谷川智幸 (阪府大理) Riccati 方程式を用いる半分線形微分方程式の非振動解の研究 ..... 10  
 Tomoyuki Tanigawa (Osaka Pref. Univ.) A study of nonoscillatory solutions of half-linear differential equations  
 by Riccati equations

概要 In this talk an attempt is made to depict a clear picture of the overall structure of nonoscillatory solutions of the second order half-linear differential equation (E)  $(p(t)\varphi_\alpha(x'))' + q(t)\varphi_\alpha(x) = 0$ , where  $\alpha > 0$  is a constant,  $p(t)$  and  $q(t)$  are positive continuous functions on  $[0, \infty)$ , and  $\varphi_\gamma(u) = |u|^\gamma \text{sgn } u$ ,  $u \in \mathbb{R}$ ,  $\gamma > 0$ . A special mention should be made of the fact that all possible types of nonoscillatory solutions of (E) can be constructed by solving the Riccati type differential equations associated with (E).

- 13 柴田徹太郎 (広島大工) Simple proof of stationary phase method and application to oscillatory  
 加藤圭一 (東京理大理) bifurcation problems ..... 10  
 Tetsutaro Shibata (Hiroshima Univ.) Simple proof of stationary phase method and application to oscillatory  
 Keiichi Kato (Tokyo Univ. of Sci.) bifurcation problems

概要 We consider the global and local behavior of bifurcation curves of semilinear eigenvalue problems, in which some special oscillatory nonlinearities and bifurcation parameter  $\lambda$  are included. Let  $\alpha = \|u_\lambda\|_\infty$  be the maximum norm of the solution  $u_\lambda$  associated with  $\lambda$ . Then in our situation,  $\lambda$  is a continuous function of  $\alpha > 0$ . Therefore, we write  $\lambda = \lambda(\alpha)$ . The goal of this talk is to establish the precise asymptotic formulas for  $\lambda(\alpha)$  as  $\alpha \rightarrow \infty$  and  $\alpha \rightarrow 0$  with the exact second and third terms by using time-map method and stationary phase method.

- 14 上村豊 (東京海洋大海洋) エネルギー依存無反射逆散乱理論と方法 ..... 10  
 Yutaka Kamimura (Tokyo Univ. of Marine Sci. and Tech.) Energy dependent reflectionless inverse theory and method

概要 We give a reflectionless inverse scattering theory on an energy dependent Schrödinger equation and, based on it, develop an inverse scattering method for an isospectral flow for the equation.

## 14:15~16:15

- 15 竹内 慎吾 (芝浦工大システム理工) Lyapunov-type inequalities for a Sturm–Liouville problem of the one-dimensional  $p$ -Laplacian ..... 10  
 渡辺宏太郎 (防衛大)  
Shingo Takeuchi (Shibaura Inst. of Tech.) Lyapunov-type inequalities for a Sturm–Liouville problem of the one-dimensional  $p$ -Laplacian  
 Kohtaro Watanabe (Nat. Defense Acad. of Japan)

概要 We will give Lyapunov-type inequalities for a Sturm–Liouville problem of the one-dimensional  $p$ -Laplacian. In the linear case of  $p = 2$ , Borg and Ha showed the Lyapunov-type inequalities by the self-adjoint property of an operator and the Green function for a boundary value problem. However, no one has ever extended those inequalities to the nonlinear case  $p \neq 2$ . Using another approach and the generalized trigonometric functions, we will establish those inequalities. Moreover, an inequality we obtained yields the best constant for the Sobolev embedding  $W_0^{1,p} \hookrightarrow L^\infty$ .

- 16 森 竜樹 (武蔵野大工) Parametric representation of a sheet constructed by all solution to a nonlocal Allen–Cahn equation ..... 10  
 久藤 衡介 (早大理工)  
 宮本 安人 (東大数理)  
 辻川 亨 (宮崎大工)  
 四ツ谷 晶二 (龍谷大\*)  
Tatsuki Mori (Musashino Univ.) Parametric representation of a sheet constructed by all solution to a nonlocal Allen–Cahn equation  
Kousuke Kuto (Waseda Univ.)  
 Yasuhito Miyamoto (Univ. of Tokyo)  
 Tohru tsujikawa (Univ. of Miyazaki)  
 Shoji Yotsutani (Ryukoku Univ.\*)

概要 We are interested in the Neumann problem of a 1D stationary Allen–Cahn equation with a nonlocal term. We have obtained the global bifurcation diagram of stationary solutions, which includes the secondary bifurcation from the odd symmetric solution due to the symmetric breaking effect. Furthermore, we derive the stability/instability of all symmetric solutions. However, stability/instability of asymmetric solutions is not clarified. In this talk, we give new representation formula of a sheet consisted of all solutions to investigate stability/instability of asymmetric solutions.

- 17 梅津健一郎 (茨城大教育) Global exact multiplicity of positive solutions for an indefinite sublinear Robin problem ..... 10  
 U. Kaufmann (Univ. Nacional de Córdoba)  
 H. Ramos Quoirin (Univ. de Santiago de Chile)  
Kenichiro Umezū (Ibaraki Univ.) Global exact multiplicity of positive solutions for an indefinite sublinear Robin problem  
 Uriel Kaufmann (Univ. Nacional de Córdoba)  
 Humberto Ramos Quoirin (Univ. de Santiago de Chile)

概要 In this talk we consider an indefinite sublinear elliptic problem with a Robin boundary condition. We aim to provide a precise description of the positive solutions set for this problem, especially proving a global exact multiplicity result for positive solutions. Our methods rely on a priori bounds for positive solutions, the linear stability argument, bifurcation approach, and sub and supersolutions.

- 18 佐藤 洋平 (埼玉大理工) Localized solutions of nonlinear Schrödinger systems with critical frequency for infinite attractive case ..... 10  
Xiaojun Chang  
(Northeast Normal Univ.)
- Yohei Sato (Saitama Univ.) Localized solutions of nonlinear Schrödinger systems with critical frequency for infinite attractive case  
Xiaojun Chang  
(Northeast Normal Univ.)

概要 We consider the singular perturbation problem of the nonlinear Schrödinger system for infinite attractive case. We construct localized solutions concentrating around a local minimum of some function  $b(V_1(x), V_2(x))$  which depends on the two potential functions  $V_i$  ( $i = 1, 2$ ). In particular we also consider the case where local minimum value of  $b(V_1(x), V_2(x))$  is 0. We make clear that, when dimension  $N = 1, 2$ ,  $b(V_1(x), V_2(x)) = 0$  if and only if  $V_1(x) = 0$  and  $V_2(x) = 0$ , when  $N = 3$ ,  $b(V_1(x), V_2(x)) = 0$  if and only if  $V_1(x) = 0$  or  $V_2(x) = 0$ .

- 19 L. Cavallina (東北大情報) 二相等周問題について ..... 10  
A. Henrot  
(Inst. Elie Cartan de Lorraine・Univ. de Lorraine)
- 坂口 茂 (東北大情報) On the two-phase isoperimetric problem  
Lorenzo Cavallina (Tohoku Univ.)  
Antoine Henrot  
(Inst. Elie Cartan de Lorraine/Univ. de Lorraine)
- Shigeru Sakaguchi (Tohoku Univ.)

概要 In this talk, we will deal with a variation of the classical isoperimetric problem in dimension  $N \geq 2$  for a two-phase piecewise continuous density whose discontinuity interface is a given hyperplane. We consider a weighted perimeter functional with three different weights, one for the hyperplane and one for each of the two open half-spaces in which  $\mathbb{R}^N$  gets partitioned. We then consider the problem of characterizing the sets  $\Omega$  that minimize this weighted perimeter functional under the additional constraint that the volumes of the portions of  $\Omega$  in the two half-spaces are given. We will provide a complete classification of the minimizers depending on the various parameters of the problem.

- 20 坂口 茂 (東北大情報) 2相熱伝導体における定温度を持つ界面の特徴付け ..... 10  
L. Cavallina (東北大情報)  
宇田川 誠一 (日大医)
- Shigeru Sakaguchi (Tohoku Univ.) A characterization of the interface with constant temperature in two-phase heat conductors  
Lorenzo Cavallina (Tohoku Univ.)  
Seiichi Udagawa (Nihon Univ.)

概要 We consider the Cauchy problem for the heat diffusion equation in the whole Euclidean space consisting of two media with different constant conductivities, where initially one has temperature 0 and the other has temperature 1. Suppose that the interface is connected and uniformly of class  $C^6$ . We show that if the interface has a time-invariant constant temperature, then it must be a hyperplane.

- 21 塚本 悠暉 (東工大理) 与えられた関数を平均曲率に持つ曲面の存在性について ..... 10  
Yuki Tsukamoto (Tokyo Tech) Existence of a prescribed anisotropic mean curvature problem

概要 We consider a prescribed mean curvature problem. In particular, we want to construct a surface whose mean curvature vector coincides with the normal component of a given vector field. We show that the problem has a solution near a graphical minimal surface if the prescribed vector field is sufficiently small in a dimensionally sharp Sobolev norm.

- 22 吉澤研介(東北大理)<sup>b</sup> Existence and non-existence of elastic graphs with the symmetric cone obstacle ..... 10  
 Kensuke Yoshizawa (Tohoku Univ.) Existence and non-existence of elastic graphs with the symmetric cone obstacle

概要 This talk is concerned with the variational problem for the bending energy defined on symmetric graphs under the unilateral constraint. In this talk, assuming that the obstacle function satisfies the symmetric cone condition, we prove (i) uniqueness of minimizers, (ii) loss of regularity of minimizers, and give (iii) complete classification of existence and non-existence of minimizers in terms of the size of obstacle.

**16:30~17:30 2019年度(第18回)日本数学会解析学賞受賞特別講演**

- 坂井秀隆(東大数理) Painlevé 方程式の世界  
 Hidetaka Sakai (Univ. of Tokyo) The world of the Painlevé equations

概要 More than a century has passed since the Painlevé equations appeared as equations that defines the special functions next to elliptic functions and hypergeometric functions. I have been studying the extension of the Painlevé equations to discrete dynamical systems or higher dimensional cases. In this talk we will see the extended world of the Painlevé equations.

3月17日(火) 第Ⅲ会場

**9:00~12:00**

- 23 濱本直樹(阪市大理) ソレノイダル場に対する Rellich 不等式の最良定数について ..... 10  
 Naoki Hamamoto (Osaka City Univ.) Sharp Rellich inequality for vector-valued functions under the solenoidal condition

概要 We show Rellich inequality with sharp constant for vector valued function under solenoidal (namely divergence-free) condition. This is a continuation of the preceding work by N. Hamamoto on sharp Rellich–Leray inequality for axisymmetric divergence-free vector fields. Our main result asserts that the sharp constant can be derived without the axisymmetry condition which was used in the preceding work for technical reason. We remove such a symmetry condition and compute the best constant. Our technique is based on an appropriate  $L^2(\mathbb{S}^{N-1})$  decomposition of solenoidal fields into radial and spherical directions, which does not depend on the choice of polar angular coordinates.

- 24 佐野めぐみ(広島大工) 改良型 Hardy–Sobolev 不等式に関連した最小化問題 ..... 10  
 Megumi Sano (Hiroshima Univ.) Minimization problem associated with an improved Hardy–Sobolev type inequality

概要 We consider the existence and the non-existence of a minimizer of minimization problems associated with an improved Hardy–Sobolev type inequality. Only for radial functions, the improved Hardy–Sobolev inequality is equivalent to the classical Hardy–Sobolev inequality via some transformation. Therefore we completely know the existence and the non-existence of a minimizer of our minimization problem only for radial functions via the transformation. In this talk, we consider it for all functions without the transformation. Especially, in contrast to the classical results, we show the existence of a minimizer for our minimization problem with the Hardy–Sobolev critical exponent on a bounded domain.

- 25 勝 呂 剛 志 (東 北 大 理)<sup>b</sup> ある拡張エントロピーに対する Shannon の不等式と不確定性原理への応用 ..... 10  
 Takeshi Suguro (Tohoku Univ.) Shannon's inequality for a generalized entropy and an application to the uncertainty principle

概要 We consider Shannon's inequality for the Rényi entropy, which is a generalization of the Boltzmann–Shannon entropy. By using some relative entropies, we identify the sharp constant and the extremal of this inequality. Moreover, we derive an extension of the Heisenberg uncertainty principle.

- 26 川 上 翔 矢 (埼 玉 大 理 工) O'Hara エネルギーの変分公式の評価について ..... 10  
 長 澤 壯 之 (埼 玉 大 理 工)  
 Shoya Kawakami (Saitama Univ.) Estimates on variational formulae of O'Hara's energies  
 Takeyuki Nagasawa (Saitama Univ.)

概要 O'Hara's energies, introduced by O'Hara, were proposed to answer the question, "What is the canonical knot in a given knot class?". One of O'Hara's energies is named the Möbius energy as a result of its Möbius invariance. Several researchers have derived the first variational formula of O'Hara's energies in the terms of Cauchy's principal value integral. Subsequently, for a certain family of O'Hara's energies which includes the Möbius energy, Ishizeki–Nagasawa found expressions for the first and second variational formulae in terms of absolutely convergent integrals. Their argument was based on an appropriate decomposition of the energies; however, such an approach is not applicable for all of O'Hara's energies. In this talk, we present an alternative method for deriving variational formulae enabling us to handle all of O'Hara's energies.

- 27 石 関 彩 (千 葉 大 理) 分解された Möbius エネルギーの上界・下界と連続度評価 ..... 10  
 長 澤 壯 之 (埼 玉 大 理 工)  
 Aya Ishizeki (Chiba Univ.) Upper and lower bounds and modulus of continuity of decomposed  
 Takeyuki Nagasawa (Saitama Univ.) Möbius energies

概要 The Möbius energy is one of knot energies, and is named after its Möbius invariant property. It is known that the energy has several different expressions. One is in terms of the cosine of conformal angle, and this expression is called the cosine formula. Another is the decomposition into Möbius invariant parts, called the decomposed Möbius energy, which was proved by the authors. Hence the cosine formula is sum of the decomposed energies. There is a question. Can each of decomposed energy be estimated by the cosine formula? Here we give an affirmative answer: upper and lower bound, and modulus of continuity of decomposed parts can be evaluated in terms of the cosine formula.

- 28 長 澤 壯 之 (埼 玉 大 理 工) 一般回転数の平面閉曲線に対する非局所曲率流の漸近解析 ..... 10  
 中 村 恒 平 (埼 玉 大 理 工)  
 Takeyuki Nagasawa (Saitama Univ.) Asymptotic analysis for non-local curvature flows for plane curves with  
 Kohei Nakamura (Saitama Univ.) general rotation number

概要 Several non-local curvature flows for plane curve with general rotation number are considered here. The flows include the arc-preserving flow and the length-preserving flow. The research for the curve with the rotation number one has been well studied. In particular, when the initial curve is strictly convex, the flow converges to a circle as time to infinity. Even if the initial curve is not strictly convex, a global solution, if exists, converges to a circle. Here we deal with curves with general rotation number, and show not only a similar result for global solutions but also a blow-up criteria, estimates of the blow-up time and blow-up rate from below. We use a geometric quantity which has never been considered before.

- 29 古川 賢 (東大数理) 動的境界条件付き高階楕円型方程式の可解性について ..... 10  
梶原 直人 (東京理大理工)  
Ken Furukawa (Univ. of Tokyo) On the solvability of higher-order elliptic equations  
Naoto Kajiwara (Tokyo Univ. of Sci.)

概要 In this talk, we give a sufficient condition for the solvability of higher-order elliptic equations in maximal  $L^p$ - $L^q$  regularity settings. First, conditions for coefficients and so-called Lopatinskii–Shapiro conditions are introduced. Furthermore, the asymptotic Lopatinskii–Shapiro condition is required to deal with the boundedness of the symbol operators. Next, we show our method to solve our problem.

- 30 本多 泰理 (東洋大情報連携) Mathematical analysis on a target detection model ..... 10  
Hirokata Honda (Toyo Univ.) Mathematical analysis on a target detection model

概要 This study aims to discuss the global existence of a target detection model with a moving target. We formulate the problem of a target detection model in drug delivery, which describes the behavior of bio-nanomachines, attractant, and repellent in a spatially 2-dimensional domain. Then, we consider the linear problem and provide the global solution under the smallness condition on the initial data. Furthermore, we obtain the strong solution in anisotropic Sobolev–Slobodetskii spaces.

- 31 井上 順平 (電通大) 拡散ロジスティック方程式における最適棲息分布と  $L^1$  非有界な定常解列  
久藤 衡介 (早大理工) の存在について ..... 10  
Junpei Inoue (Univ. of Electro-Comm.) On the optimal distribution and the existence of an  $L^1$ -unbounded se-  
Kousuke Kuto (Waseda Univ.) quence of steady states for the diffusive logistic equation

概要 We discuss a stationary diffusive logistic equation on a ball. This talk focuses on an open question that showing the upper bound of the ratio of a total population to total resources. In one-dimensional case, Bai–He–Li (2015) settled that the supremum is equal to 3 by finding a special sequence of diffusion coefficients and carrying functions, and moreover, the first speaker recently obtained profiles of solutions corresponding to the maximizing sequence. A new question is the following: What happens in higher-dimensional cases? This talk shows that the supremum is infinite on the spherical symmetry domain.

- 32 下條 昌彦 (岡山理大理) 特異被食–捕食モデルにおける捕食者の侵入速度 ..... 10  
郭 忠勝 (淡江大)  
Yu-Shuo Chen (淡江大)  
Masahiko Shimojyou Spreading speed of a singular prey-predator type reaction-diffusion sys-  
(Univ. of Sci.) tem  
Jong-Shenq Guo (Tamkang Univ.)  
Yu-Shuo Chen (Tamkang Univ.)

概要 We consider the dynamical behaviors of a singular predator-prey model and give some new results on the spreading speed of the predator.

- 33 下條 昌彦 (岡山理大理) 特異被食–捕食モデルの進行波 ..... 10  
郭 忠勝 (淡江大)  
Yu-Shuo Chen (淡江大)  
Masahiko Shimojyou Traveling wave solution to a singular prey-predator reaction diffusion  
(Univ. of Sci.) system  
Jong-Shenq Guo (Tamkang Univ.)  
Yu-Shuo Chen (Tamkang Univ.)

概要 We give some results on the existence vs non-existence of traveling waves connecting the predator-free state to the co-existence state, and the existence vs non-existence of spatially periodic traveling waves to the singular predator-prey reaction diffusion system.

- 34 岡 大 将 (東 北 大 理) Space-time homogenization for the fast diffusion equation ..... 10  
 赤 木 剛 朗 (東 北 大 理)  
 Tomoyuki Oka (Tohoku Univ.) Space-time homogenization for the fast diffusion equation  
 Goro Akagi (Tohoku Univ.)

概要 In this talk, we shall discuss a space-time homogenization problem for the fast diffusion equation with periodically oscillating (in space and time) coefficients. Main results consist of a homogenization theorem, i.e., convergence of solutions as the period of coefficients goes to zero, and a characterization of the homogenized equation. In particular, homogenized matrices are described in terms of solutions to cell-problems and depend on the log-ratio of spatial and temporal periods of the coefficients.

- 35 中 村 謙 太 (東 北 大 理)  $p$ -Sobolev 流の時間大域解の存在について ..... 10  
 三 沢 正 史 (熊本大先端科学)  
 T. Kuusi (Univ. of Helsinki)  
 Kenta Nakamura (Tohoku Univ.) Global existence for the  $p$ -Sobolev flow  
 Masashi Misawa (Kumamoto Univ.)  
 Tuomo Kuusi (Univ. of Helsinki)

概要 In this talk, we study a doubly nonlinear parabolic equation arising from the gradient flow for  $p$ -Sobolev type inequality, referred as  $p$ -Sobolev flow from now on, which includes the classical Yamabe flow on a bounded domain in Euclidean space in the special case  $p = 2$ . We present the global existence of the  $p$ -Sobolev flow.

- 36 M. Fuest (Paderborn Univ.) Asymptotic behavior in a chemotaxis-consumption model with realistic  
 J. Lankeit (Paderborn Univ.) boundary conditions for the oxygen ..... 10  
 水 上 雅 昭 (東 京 理 大 理)  
 Mario Fuest (Paderborn Univ.) Asymptotic behavior in a chemotaxis-consumption model with realistic  
 Johannes Lankeit (Paderborn Univ.) boundary conditions for the oxygen  
 Masaaki Mizukami  
 (Tokyo Univ. of Sci.)

概要 This talk considers global existence and asymptotic behavior in a chemotaxis-consumption system under realistic boundary conditions for the oxygen. In previous works a chemotaxis-consumption system under the Neumann boundary condition for the oxygen is mainly considered, and it is shown that solutions of the problem converge to constant steady states by Tao–Winkler (2012); however this result does not describe pattern formation of species. Thus it might be important to consider a chemotaxis-consumption system under realistic boundary conditions for the oxygen. This talk shows solutions of the problem converge to non-constant steady states.

- 37 鈴 木 貴 (阪 大 M M D S) A parabolic concavity maximum principle ..... 5  
 Takashi Suzuki (Osaka Univ.) A parabolic concavity maximum principle

概要 Positive solutions to the parabolic equation on strictly convex domain preserve log concavity of the initial value.

## 13:15~14:15 特別講演

- 蘆田 聡平 (学習院大理) 電子のハミルトニアンの固有値の正確な下界評価  
Sohei Ashida (Gakushuin Univ.) Accurate lower bounds for eigenvalues of electronic Hamiltonians

概要 Electronic Hamiltonians are differential operators depending on relative positions of nuclei as parameters. When we regard an eigenvalue of an electronic Hamiltonian as a function of relative positions of nuclei, minimum points correspond to shapes of molecules. Upper bounds for eigenvalues are obtained by variational methods. However, since only relative energy is relevant to the physical information as minimum points, physical information can not be obtained by variational methods only. Therefore, lower bounds are helpful for physical information to be available. In this talk we discuss the various methods for lower bounds of eigenvalues. In particular, lower bounds for eigenvalues of sums of lower semibounded self-adjoint operators are introduced. Some computations for systems of one electron and several protons are shown.

3月18日(水) 第III会場

## 9:00~12:00

- 38 國谷 紀良 (神戸大システム情報) 異なる境界条件下での拡散を伴う SIR 感染症モデルの閾値定理 ..... 10  
Toshikazu Kuniya (Kobe Univ.) Threshold theorem for an SIR epidemic model with diffusion under the different boundary conditions

概要 In this study, we are concerned with a threshold theorem for an SIR epidemic model with diffusion. We consider two different boundary conditions: (homogeneous) Dirichlet and (homogeneous) Neumann boundary conditions. We show that if the basic reproduction number  $\mathcal{R}_0$  satisfies  $\mathcal{R}_0 < 1$ , then the disease-free equilibrium  $E_0$  of the model is globally asymptotically stable, whereas if  $\mathcal{R}_0 > 1$ , then  $E_0$  is unstable and a positive endemic equilibrium  $E^*$  exists under an additional condition. We show that, under the Neumann boundary conditions,  $\mathcal{R}_0$  does not depend on the shape of the spatial domain  $\Omega \subset \mathbb{R}^2$ , whereas, under the Dirichlet boundary conditions,  $\mathcal{R}_0$  depends on the shape of  $\Omega$ . More precisely, we show that such  $\mathcal{R}_0$  can attain its maximum when  $\Omega$  is a square domain.

- 39 鄭 大樹 (名大多元数理) The role of forward self-similar solutions in the Cauchy problem for semi-linear heat equations with exponential nonlinearity ..... 10  
Daesu Jeong (Nagoya Univ.) The role of forward self-similar solutions in the Cauchy problem for semi-linear heat equations with exponential nonlinearity

概要 In this talk, we consider the Cauchy problem for semi-linear heat equations with exponential nonlinearity. The main purpose of this talk is to prove the existence of solutions lying on the borderline between global existence and blow-up infinite time. The existence has been shown for semi-linear heat equations with power type nonlinearity. We explain the main strategy to prove the existence. By using the definition of exponential function, we approximate the solution to exponential type equation by that to power type equation. Then we can use directly the knowledge for power type equation.

- 40 比佐 幸太郎 (東北大大理) Existence of solutions for an inhomogeneous fractional semilinear heat  
石毛 和弘 (東大数理) equation ..... 10  
高橋 仁 (東工大情報理工)  
Kotaro Hisa (Tohoku Univ.) Existence of solutions for an inhomogeneous fractional semilinear heat  
Kazuhiro Ishige (Univ. of Tokyo) equation  
Jin Takahashi (Tokyo Tech)

概要 We obtain necessary conditions and sufficient conditions on the existence of solutions to the Cauchy problem for a fractional semilinear heat equation with an inhomogeneous term. We identify the strongest spatial singularity of the inhomogeneous term for the solvability of the Cauchy problem.



- 41 M. R. Haque (東北大理)<sup>b</sup> Critical existence to a convection-diffusion equation in a uniformly local  
猪奥倫左(東北大理) lebesgue space ..... 10  
小川卓克(東北大理)  
佐藤龍一(東北大理)  
Md Rabiul Haque (Tohoku Univ.) Critical existence to a convection-diffusion equation in a uniformly local  
Norisuke Ioku (Tohoku Univ.) lebesgue space  
Takayoshi Ogawa (Tohoku Univ.)  
Ryuichi Sato (Tohoku Univ.)

概要 We establish the local in time existence of classical solutions to the Cauchy problem of the convection-diffusion equations in uniformly local Lebesgue spaces. For the proof, we use uniformly local  $L^p$ - $L^q$  estimate for some convolution operators and the Banach fixed point theorem.

- 42 石田祥子 (千葉大理) Weak stabilization in Keller–Segel systems with degenerate diffusion · · 10  
横田智巳 (東京理大理)  
Sachiko Ishida (Chiba Univ.) Weak stabilization in Keller–Segel systems with degenerate diffusion  
Tomomi Yokota (Tokyo Univ. of Sci.)

概要 We focus on stabilization in Keller–Segel systems with degenerate diffusion. As to the known studies on the non-degenerate diffusion systems, Cao (2015) and Cieřlak–Winkler (2017) obtained that  $u(t) \rightarrow \bar{u}_0 := \frac{1}{|\Omega|} \int_{\Omega} u_0$  in  $L^\infty(\Omega)$  as  $t \rightarrow \infty$  for small initial data. Moreover, for chemotaxis–Stokes systems with degenerate diffusion, Winkler (2015) found that  $u(t) \rightarrow \bar{u}_0$  weakly\* in  $L^\infty(\Omega)$  as  $t \rightarrow \infty$  by using a proof by contradiction. Although their argument by contradiction actually can be applied to our system, we present the same stabilization with direct proof. Moreover our proof can be applied to chemotaxis–Navier–Stokes systems, tumor invasion systems and generalized parabolic equations with  $L^1$ -conservation law.

- 43 藤井幹大 (九大数理) Global solutions to the dissipative quasi-geostrophic equation with dis-  
persive forcing ..... 10  
Mikihiro Fujii (Kyushu Univ.) Global solutions to the dissipative quasi-geostrophic equation with dis-  
persive forcing

概要 We consider the initial value problem for the dispersive 2D quasi-geostrophic equation with critical and supercritical dissipation  $(-\Delta)^\alpha \theta$  ( $0 < \alpha \leq 1$ ). We establish a unique global solution for a given initial data  $\theta_0$  which belongs to the scaling subcritical Sobolev space  $H^s(\mathbb{R}^2)$  ( $s > 2 - \alpha$ ) if the size of dispersion parameter is sufficiently large. This phenomenon is so-called the global regularity. We also obtain the relationship between the initial data and the dispersion parameter, which ensures the existence of the global solution.

- 44 黒木場正城 (室蘭工大)<sup>b</sup> Keller–Segel 方程式の移流拡散方程式への零緩和時間極限について · · · · 10  
小川卓克  
(東北大 RACMaS · 東北大理)  
Masaki Kurokiba Singular limit problem for the Keller–Segel system and drift-diffusion  
(Muroran Inst. of Tech.) system  
Takayoshi Ogawa  
(Tohoku Univ./Tohoku Univ.)

概要 We consider a singular limit problem for the Cauchy problem of the Keller–Segel equation in a critical function space. We show that a solution to the Keller–Segel system in a scaling critical function space converges to a solution to the drift-diffusion system of parabolic-elliptic type (the simplified Keller–Segel model) in the critical space strongly as the relaxation time  $\tau \rightarrow \infty$ . For the proof of singular limit problem, we employ generalized maximal regularity for the heat equation and use it systematically with the sequence of embeddings between the interpolation spaces  $\dot{B}_{q,\sigma}^s(\mathbb{R}^n)$  and  $\dot{F}_{q,\sigma}^s(\mathbb{R}^n)$ .

- 45 吉川周二 (大分大理工) 二次元 Cahn–Hilliard 方程式に対する構造保存型離散近似とその誤差評価  
 ..... 10

Shuji Yoshikawa (Oita Univ.) Error estimates of structure-preserving discrete approximations for the Cahn–Hilliard equation in two space dimension

概要 We introduce two structure-preserving finite difference schemes in two-dimensional space with the Voronoi mesh. By applying the energy method to these schemes, we show the unique existence of a solution for these and error estimates between these and strict solutions.

- 46 古屋貴士 (名大多元数理) Direct and inverse scattering problems for the local perturbation of an open periodic waveguide in the half plane ..... 10

Takashi Furuya (Nagoya Univ.) Direct and inverse scattering problems for the local perturbation of an open periodic waveguide in the half plane

概要 We consider the direct and inverse scattering problem of the local perturbation of an open periodic waveguide in the half plane. Our first aim is to show the well-posedness of the direct scattering problem under the suitable radiation condition for this setting. Our second aim is to solve the inverse scattering problem of determining the support of the perturbation function from scattering measurements.

- 47 肥田野久二男 (三重大教育) Global existence for null-form wave equations with data in a Sobolev space of lower regularity and weight ..... 10

横山和義 (北海道科学大工) space of lower regularity and weight  
 Kunio Hidano (Mie Univ.) Global existence for null-form wave equations with data in a Sobolev space of lower regularity and weight  
 Kazuyoshi Yokoyama (Hokkaido Univ. of Sci.)

概要 Assuming initial data have small weighted  $H^4 \times H^3$  norm, we show global existence of solutions to the Cauchy problem for systems of quasi-linear wave equations in 3D satisfying the null condition of Klainerman. Compared with the work of Christodoulou, our result assumes smallness of data with respect to  $H^4 \times H^3$  norm having a lower weight. Our proof uses the ghost weight energy method due to Alinhac. In comparison with the proofs of Klainerman and Hörmander, we can limit the number of occurrences of the generators of hyperbolic rotations or dilations in the course of a priori estimates of solutions. This limitation allows us to obtain global solutions for radially symmetric data, when a certain norm with considerably low weight is small enough.

- 48 Tadahiro Oh (Univ. of Edinburgh) 空間 2 次元 2 次の非線形項と粗いノイズを持つ確率波動方程式の初期値問題の適切性について ..... 10

岡本葵 (信州大工) On the well-posedness for the quadratic stochastic nonlinear wave equation with a rough noise in two dimensions  
 Tadahiro Oh (Univ. of Edinburgh)  
 Mamoru Okamoto (Shinshu Univ.)

概要 We study the two-dimensional stochastic nonlinear wave equation (SNLW) with a quadratic nonlinearity, forced by a fractional derivative (of order  $\alpha > 0$ ) of a space-time white noise. In particular, we show that the well-posedness theory breaks at  $\alpha = \frac{1}{2}$  for SNLW, while the threshold is  $\alpha = 1$  for the stochastic nonlinear heat equation (SNLH). This provides an example showing that SNLW behaves less favorably than SNLH.

- 49 水谷治哉 (阪大理工) Hardy ポテンシャルを伴う分数べき Schrödinger 作用素のレゾルベント評価と Strichartz 評価 ..... 10

Xiaohua Yao (華中師範大) Resolvent and Strichartz estimates for fractional Schrödinger operators with Hardy potentials  
 Haruya Mizutani (Osaka Univ.)  
 Xiaohua Yao (Central China Normal Univ.)

概要 We discuss recent progress on uniform resolvent estimates of Kato–Yajima type and Strichartz estimates for fractional or higher-order Schrödinger operators with Hardy potentials involving subcritical coupling constants related to the best constant in the fractional Hardy inequality.

- 50 田中智之<sup>b</sup> Global well-posedness for the wave equation with a time-dependent scale  
 (名大多元数理・中大理工・理化学研AIP・慶大理工) invariant damping and a cubic convolution ..... 10  
 池田正弘  
 (理化学研AIP・慶大理工)  
 若狭恭平(釧路工高専)  
 Tomoyuki Tanaka Global well-posedness for the wave equation with a time-dependent scale  
 (Nagoya Univ./Chuo Univ./RIKEN/Keio Univ.) invariant damping and a cubic convolution  
 Masahiro Ikeda (RIKEN/Keio Univ.)  
 Kyouhei Wakasa  
 (Kushiro Nat. Coll. of Tech.)

概要 In this talk, we consider global well-posedness for the wave equation with a time-dependent scale invariant damping, i.e.,  $\frac{2}{1+t}\partial_t u$  and a cubic convolution  $(|x|^{-\gamma} * u^2)u$ , where  $0 < \gamma < n$ . For a power type nonlinearity, the work of D'Abbico, Lucente and Reissig shows that a critical exponent, which divides global existence and blow-up for small solutions, is shifted because of the presence of the damping term. Our aim of this work is to determine two types of critical exponents of the problem with the cubic convolution. The one is for compactly supported initial data. The second is a critical exponent about the spatial decay condition on the data.

- 51 隠居良行(東工大理) 準線形消散型弾性波方程式に対する初期値問題の解の時間大域挙動につ  
 竹田寛志(福岡工大) いて ..... 10  
 Yoshiyuki Kagei (Tokyo Tech) Large time behavior of global solutions to nonlinear elastic wave equa-  
 Hiroshi Takeda (Fukuoka Inst. of Tech.) tions with strong damping term

概要 In this talk we consider the Cauchy problem of quasi-linear elastic wave equations with the strong damping term. We prove the existence of the global solution with decay property and its asymptotic profile is given by the diffusion waves as time tends to infinity. Our proof is based on the combination of the semi-group theory and the energy method.

#### 14:15~16:15

- 52 加藤勲(京大理) On the 3D Zakharov system with radial initial data ..... 10  
 木下真也(Univ. Bielefeld)  
 Isao Kato (Kyoto Univ.) On the 3D Zakharov system with radial initial data  
 Shinya Kinoshita (Univ. Bielefeld)

概要 We study the Cauchy problem for the Zakharov system in three spatial dimensions. We prove the global well-posedness at the critical space with radially symmetric and small initial data by  $U^2, V^2$  method introduced by Koch and Tataru. We follow the argument by Kato and Tsugawa, which was proved small data G.W.P. at the critical space for four and higher spatial dimensions. However, we need spherically symmetric data to obtain well-posedness at the critical space.

- 53 清水一慶(京大理) Local well-posedness for Schrödinger maps with helicity terms ..... 10  
 Ikkei Shimizu (Kyoto Univ.) Local well-posedness for Schrödinger maps with helicity terms

概要 We consider the initial-value problem for Schrödinger maps with helicity terms. We prove the local well-posedness in two different settings: the general case and the topologically-free case. In the former case, we establish local well-posedness by applying the energy method of McGahagan with improving her argument. In the later case, we show additional properties for solutions, such as blow-up criterion, quantitative bounds and difference estimates. The key idea is the reduction of the problem to a kind of nonlinear Schrödinger equation, so-called the modified Schrödinger map equation. For its analysis, the treatment for magnetic potentials is required.

- 54 Chunhua Li (Yanbian Univ.) Large time asymptotics for a cubic nonlinear Schrödinger system in one space dimension ..... 10  
 西井良徳(阪大 理)  
 佐川侑司  
 砂川秀明(阪大 理)  
 Chunhua Li (Yanbian Univ.) Large time asymptotics for a cubic nonlinear Schrödinger system in one space dimension  
 Yoshinori Nishii (Osaka Univ.)  
 Yuji Sagawa  
 Hideaki Sunagawa (Osaka Univ.)

概要 We consider a two-component system of cubic nonlinear Schrödinger equations in one space dimension. We show that each component of the solutions to this system behaves like a free solution in the large time, but there is a crucial restriction between the profiles of them. This turns out to be a possible consequence of non-trivial long-range nonlinear interactions.

- 55 駒田 洸一 (東北大理・九大数理) Existence of blow-up solutions to nonlinear Schrödinger equations with anisotropic fourth-order dispersion ..... 10  
 Koichi Komada Existence of blow-up solutions to nonlinear Schrödinger equations with anisotropic fourth-order dispersion  
 (Tohoku Univ./Kyushu Univ.)

概要 We consider the Cauchy problem for nonlinear Schrödinger equations with anisotropic fourth-order dispersion. In this talk, we give sufficient conditions for the existence of blow-up solutions. In the proof, we use the localized virial identity which was introduced by T. Ogawa and Y. Tsutsumi (1991) to prove the existence of blow-up solutions to NLS with radial data. Since in our problem, equations have no radially symmetric structure in full spatial dimensions because of the anisotropic fourth-order term, we need to modify the argument.

- 56 佐藤拓也(東北大 理)<sup>b</sup>  $L^2$ -decay for the one dimensional dissipative nonlinear Schrödinger equation in a critical exponent ..... 10  
 小川卓克(東北大 理)  
 Takuya Sato (Tohoku Univ.)  $L^2$ -decay for the one dimensional dissipative nonlinear Schrödinger equation in a critical exponent  
 Takayoshi Ogawa (Tohoku Univ.)

概要 We consider the Cauchy problem for the dissipative nonlinear Schrödinger equation with cubic nonlinearities in one space dimension. For the dissipative nonlinear term, the cubic nonlinearity in one space dimension is the threshold to exhibit the  $L^2$ -decay of solutions. We prove the existence for the global analytic solution and show the  $L^2$ -decay of the solution in the critical exponent.

- 57 鈴木敏行(神奈川大 工) Nonlinear Schrödinger equations with an inverse-square potential and a repulsive harmonic oscillator ..... 10  
 Toshiyuki Suzuki (Kanagawa Univ.) Nonlinear Schrödinger equations with an inverse-square potential and a repulsive harmonic oscillator

概要 We consider the Cauchy problems for nonlinear Schrödinger equations with an inverse-square potential  $a|x|^{-2}$  and a repulsive harmonic oscillator  $-\omega^2|x|^2$ . Carles (2003) proved the well-posedness and solve the scattering problems for  $a = 0$ . We consider these problems for  $a \neq 0$  by applying the pseudo-conformal type transform.

- 58 浜野 大 (埼玉大理工) ポテンシャル項を持つ非線形シュレディンガー方程式の定常問題について  
池田 正弘 (理化学研 AIP・慶大理工) .....

Masaru Hamano (Saitama Univ.) For a stationary problem of the nonlinear Schrödinger equation with a  
Masahiro Ikeda (RIKEN/Keio Univ.) potential term

概要 We deal with the nonlinear Schrödinger equation with a linear potential term. First, we consider a minimization problem with respect to the elliptic equation corresponding to the NLS. In order to apply to time behavior of solutions to the NLS, the minimization problem is not characterized by the Nehari functional but the virial functional. We prove existence of a minimizer of the problem in mass-supercritical and energy-subcritical case. Then, we consider time behavior of solutions to the NLS with initial data, whose action is less than the minimum value.

- 59 深谷 法良 (東京理大理) Instability of algebraic standing waves for nonlinear Schrödinger equations with double power nonlinearities ..... 10  
林 雅行 (京大数理研)  
Noriyoshi Fukaya (Tokyo Univ. of Sci.) Instability of algebraic standing waves for nonlinear Schrödinger equations with double power nonlinearities  
Masayuki Hayashi (Kyoto Univ.)

概要 We consider nonlinear Schrödinger equations with double power nonlinearities. This equation has two types of standing waves. One decays exponentially, and the other only decays algebraically. In this talk, we prove instability and strong instability of standing waves including algebraic standing waves. Our results extend previous results by Ohta (1995). He studied stability/instability of standing waves with exponential decay in one-dimensional case. We improve the instability results in previous works in one-dimensional case, and moreover establish new instability results in multi-dimensional cases. The key point in our approach is to take advantage of variational characterization of algebraic standing waves.

#### 16:30~17:30 特別講演

- 久藤 衡介 (早大理工) Cross-diffusion limit in the stationary SKT model  
Kousuke Kuto (Waseda Univ.) Cross-diffusion limit in the stationary SKT model

概要 This talk is concerned with the global bifurcation structure of coexistence steady-states to the Shigesada–Kawasaki–Teramoto model with cross-diffusion (so-called the SKT model). In 1999, Lou–Ni showed that the asymptotic behavior of coexistence steady-states as one of the cross-diffusion terms tends to infinity can be classified into two types (the segregation type and the shrink type). For the segregation type, the set of solutions to the corresponding limiting system (the 1st limiting system) has been revealed mainly by Lou–Ni–Yotsutani. This talk focuses on the shrink type and study the corresponding limiting system (the 2nd limiting system). We obtain the global bifurcation structure of positive solutions to the 2nd limiting system. Furthermore, by the perturbation of solutions of two limiting systems, we construct the bifurcation branch of coexistence steady-states to the SKT model in a case when one of the cross-diffusion terms is sufficiently large.

## 3月19日(木) 第III会場

## 9:00~12:00

- 60 坂本 祥太 (東 北 大 理) 非切断ボルツマン方程式の摂動初期値問題及び初期値境界値問題の時間  
Renjun Duan 大域解 ..... 10  
(Chinese Univ. of Hong Kong)  
Shuangqian Liu  
(Central China Normal Univ./Jinan Univ.)  
R. M. Strain (Univ. Pennsylvania)  
Shota Sakamoto (Tohoku Univ.) Solutions to initial and initial-boundary value problems of the non-cutoff  
Renjun Duan Boltzmann equation near an equilibrium  
(Chinese Univ. of Hong Kong)  
Shuangqian Liu  
(Central China Normal Univ./Jinan Univ.)  
Robert M. Strain (Univ. Pennsylvania)

概要 We consider initial and initial-boundary value problems of the non-cutoff Boltzmann equation near the Maxwellian equilibrium on the three-dimensional torus. Especially we considered the specular boundary reflection condition and the inflow boundary condition for the IBVP. Using the Wiener space, we have succeeded in proving the unique global existence of solutions to these problems with energy estimates. Moreover, decay rates of these solutions are also shown.

- 61 齋藤 平和 (東京理大基礎工) On elliptic problems associated with two-phase incompressible flows in  
Xin Zhang (早 大 理 工) unbounded domains ..... 10  
Hirokazu Saito (Tokyo Univ. of Sci.) On elliptic problems associated with two-phase incompressible flows in  
Xin Zhang (Waseda Univ.) unbounded domains

概要 In this talk, we consider elliptic problems associated with two-phase incompressible flows in unbounded domains. First, we prove the unique solvability of the strong elliptic problem (SEP). Next, by means of the solutions of (SEP), we prove the unique solvability of the weak elliptic problem (WEP). The result for (WEP) enables us to show the maximal regularity for a linearized system of the two-phase Navier–Stokes equations and some decomposition of Lebesgue spaces similar to the Helmholtz–Weyl decomposition.

- 62 顧 仲陽 (東 大 数 理) Continuous alignment of vorticity direction prevents the blow-up of the  
儀我 美一 (東 大 数 理) Navier–Stokes flow under the no-slip boundary condition ..... 10  
許 本源 (東 大 数 理)  
Zhongyang Gu (Univ. of Tokyo) Continuous alignment of vorticity direction prevents the blow-up of the  
Yoshikazu Giga (Univ. of Tokyo) Navier–Stokes flow under the no-slip boundary condition  
Pen-Yuan Hsu (Univ. of Tokyo)

概要 This talk is concerned with a regularity criterion based on vorticity direction for the Navier–Stokes equations in a three-dimensional bounded domain under the no-slip boundary condition. It asserts that if the vorticity direction is uniformly continuous in space uniformly in time, then there is no type I blow-up. A similar result has been proved for a half space by Y. Giga, P.-Y. Hsu and Y. Maekawa (2014). The difficulty of extending this result to bounded domains lies in establishing a good  $L^\infty$  theory for the Navier–Stokes equations in a domain with curved boundary. This is achieved based on the  $L^\infty$  theory developed by K. Abe and Y. Giga (2013).

- 63 中村 憲史 (筑波大F-MIRAI) 3次元半空間における双曲型 Navier–Stokes 方程式の線型化問題について  
 小林 孝行 (阪大基礎工) ..... 10  
 久保 隆徹 (お茶の水女大基幹)  
 Kenji Nakamura (Tsukuba Univ.) Linearized problem of the hyperbolic type Navier–Stokes equations in  
 Takayuki Kobayashi (Osaka Univ.) the three dimensional half-spaces  
 Takayuki Kubo (Ochanomizu Univ.)

概要 In this talk, we show the local-in-time unique existence of the linearized problem of the hyperbolic type Navier–Stokes equations in the three dimensional half-spaces under Dirichlet boundary condition. To do this, we add perturbation to the problem and show the unique existence of the solution to the perturbed equations. Then, by using a priori estimates, we can pass to the limit and thus we obtain the solution to the original equations.

- 64 高橋 知希 (名大多元数理) Attainability of a stationary Navier–Stokes flow around a rigid body  
 rotating from rest ..... 10  
 Tomoki Takahashi (Nagoya Univ.) Attainability of a stationary Navier–Stokes flow around a rigid body  
 rotating from rest

概要 In this talk, we consider the large time behavior of a viscous incompressible flow around a rotating rigid body. In particular, assume that both a compact rigid body in  $\mathbb{R}^3$  and a viscous incompressible fluid that occupies the outside are at rest, and that the rigid body gradually increases the angular velocity and moves at a constant angular velocity after a certain time. Then we show that the fluid motion converges to the steady solution obtained by Galdi (2003) at time  $t \rightarrow \infty$ .

- 65 岡田 晃 (京大人間環境) Necessary and sufficient condition for the local existence of solution in  
 the Serrin class of the Navier–Stokes equations ..... 10  
 Akira Okada (Kyoto Univ.) Necessary and sufficient condition for the local existence of solution in  
 the Serrin class of the Navier–Stokes equations

概要 Consider the Cauchy problem of the Navier–Stokes equations with initial data in the Besov space. We construct the local solution in the Serrin class with the initial data in the Besov space. Conversely, if the local solution belongs to the Serrin class, the initial data necessarily belongs to the Besov space. As a result, we are successful to determine the necessary and sufficient condition for the local existence of solution in the Serrin class.

- 66 岡部 考宏 (阪大基礎工) Annihilation of slow-decay factors of the Navier–Stokes flow by the ex-  
 L. Brandolese (Univ. Lyon 1) ternal force ..... 10  
 Takahiro Okabe (Osaka Univ.) Annihilation of slow-decay factors of the Navier–Stokes flow by the ex-  
 Lorenzo Brandolese (Univ. Lyon 1) ternal force

概要 We consider the Navier–Stokes equations on the whole space  $R^n$ . We discuss for any small initial data there exists an external force  $f$  which generates a rapid energy decay  $\|u(t)\|_2 = o(t^{-\frac{n+2}{4}})$  as  $t \rightarrow \infty$ .

- 67 鶴見 裕之 (早大理工) The two-dimensional stationary Navier–Stokes equations in toroidal Besov  
 spaces ..... 10  
 Hiroyuki Tsurumi (Waseda Univ.) The two-dimensional stationary Navier–Stokes equations in toroidal Besov  
 spaces

概要 We consider the stationary Navier–Stokes equations in the two-dimensional torus. For every  $\varepsilon > 0$  and  $(p, q) \in ([1, 2) \times [1, \infty]) \cup (\{2\} \times [1, 2])$ , we show the existence, uniqueness, and continuous dependence of solutions in homogeneous toroidal Besov spaces  $\dot{B}_{p+\varepsilon, q}^{-1+\frac{2}{p}}(\mathbb{T}^2)$  (if  $p = 2$ , in  $L^{2+\varepsilon}(\mathbb{T}^2)$ ) for given small external forces in  $\dot{B}_{p+\varepsilon, q}^{-3+\frac{2}{p}}(\mathbb{T}^2)$ . In addition, for the rest case of  $p$  and  $q$ , we show the ill-posedness caused by the discontinuity of the solution map.

- 68 津田和幸 (阪大基礎工) The time periodic problem of the Navier–Stokes equations in a bounded domain with moving boundary ..... 10  
 R. Farwig (TU Darmstadt)  
 小藺英雄 (早大理工)  
 D. Wegmann (TU Darmstadt)  
 Kazuyuki Tsuda (Osaka Univ.) The time periodic problem of the Navier–Stokes equations in a bounded domain with moving boundary  
 Reinhard Farwig (TU Darmstadt)  
 Hideo Kozono (Waseda Univ.)  
 David Wegmann (TU Darmstadt)

概要 The time periodic problem of the Navier–Stokes equations on a non-cylindrical space-time domain is studied. Motivated by a recent result by J. Saal (2006) on maximal regularity for this kind of system we construct time periodic solutions in  $L^q$ -spaces provided the bounded domain moves periodically with small amplitude and the given periodic external force is small. The proof is based on new decay estimates for the solution operator of parabolic evolution equations corresponding to the non-cylindrical space-time domain problem.

- 69 柴田良弘 (早大理工) 外部領域における第2ヘルムホルツ分解について ..... 10  
 Yoshihiro Shibata (Waseda Univ.) On the second Helmholtz decomposition in an exterior domain

概要 I would like to talk about the second Helmholtz decomposition in an exterior domain  $\Omega$  in the  $N$  dimensional Euclidean space  $R^N$ . This decomposition is based on the unique existence theorem for the weak Dirichlet problem. To obtain the unique existence theorem, the key is to definition of the underlying space  $\hat{H}_{q,0}^1(\Omega)$ .

- 70 柴田良弘 (早大理工) On the isothermal compressible multi-component mixture flow: the local existence and maximal  $L_p$ - $L_q$  regularity of solutions ..... 10  
 Yoshihiro Shibata (Waseda Univ.) On the isothermal compressible multi-component mixture flow: the local existence and maximal  $L_p$ - $L_q$  regularity of solutions

概要 I will talk about the local wellposedness of the isothermal compressible multi-component mixture flow in the  $L_p$ - $L_q$  maximal regularity class. The equations are derived by the Maxwell–Stefan–Navier–Stokes theory, and the equations consist of the Navier–Stokes equations describing the compressible viscous fluid flow and diffusion equations. Since the diffusion part has 1 more unknown functions, we use the transformation of unknown functions introduced by V. Giovangigli to derive the Navier–Stokes equations coupled with a system of reaction diffusion equations. I consider the equations in a bounded domain with non-slip condition for the velocity field and homogeneous Neumann condition for the diffusion part. The proof is based on the maximal  $L_p$ - $L_q$  regularity theorem for the linearized equations.

- 71 柴田良弘 (早大理工) On the maximal  $L_p$ - $L_q$  regularity of solutions to a general linear parabolic system ..... 10  
 Yoshihiro Shibata (Waseda Univ.) On the maximal  $L_p$ - $L_q$  regularity of solutions to a general linear parabolic system

概要 I will talk about the  $L_p$ - $L_q$  maximal regularity theorem for a general linear parabolic system with Neumann type boundary conditions in a uniform  $C^2$  domain. This equations arise in a mathematical study of isothermal compressible multi-component mixture flow based on the Maxwell–Stefan theory. The proof is based on the  $\mathcal{R}$ -solver for the corresponding generalized resolvent problem and Weis’ operator valued Fourier multiplier theorem.



- 72 柴田良弘 (早大理工)  $\mathcal{R}$ -solver と周期解 ..... 10  
 Yoshihiro Shibata (Waseda Univ.) On the  $\mathcal{R}$ -solver and periodic solutions

概要 I introduce the notion of  $\mathcal{R}$  solver for the generalized resolvent problem and explain how to prove the maximal regularity theorem for the high frequency part of the periodic solutions of the system of the parabolic equations with non-homogeneous boundary conditions, which includes Stokes system with the help of the transference theorem.

- 73 柴田良弘 (早大理工) Navier–Stokes 方程式の自由境界問題の周期解 ..... 10  
 Yoshihiro Shibata (Waseda Univ.) On the periodic solutions for free boundary problem of the Navier–Stokes equations

概要 I would like to talk about the existence of periodic solutions for free boundary problem of the Navier–Stokes equations. We consider the case where unknown time dependent domain is bounded close to a ball and the support of time periodic external force is in this ball.

#### 14:15~16:15

- 74 清水扇丈 (京大人間環境) 半空間での放物型方程式の初期値境界値問題に対する最大  $L^1$  正則性 .. 10  
 小川卓克 (東北大理)  
 Senjo Shimizu (Kyoto Univ.) Maximal  $L^1$ -regularity for the parabolic initial-boundary value problem  
 Takayoshi Ogawa (Tohoku Univ.) in the half-space

概要 End-point maximal  $L^1$ -regularity for the parabolic initial-boundary value problem is considered in the half-space. For the inhomogeneous boundary data of both the Dirichlet and the Neumann type, maximal  $L^1$ -regularity for the initial-boundary value problem of parabolic equation is established in time end-point case upon the Besov space as well as the optimal trace estimates. We derive the almost orthogonal properties between the boundary potentials of the Dirichlet and the Neumann boundary data and the Littlewood–Paley dyadic decomposition of unity.

- 75 岩瀬司 (東北大理) Forward self-similar solutions for compressible Navier–Stokes equations ..... 10  
 P. Germain (New York Univ.)  
 Tsukasa Iwabuchi (Tohoku Univ.) Forward self-similar solutions for compressible Navier–Stokes equations  
 Pierre Germain (New York Univ.)

概要 We construct forward self-similar solutions for compressible Navier–Stokes equations with radial symmetry. Some of these solutions are smooth, and the others exhibit a singularity due to vacuum.

- 76 中里亮介 (東北大理) Hall 効果を持つ圧縮性磁気粘性流体方程式系の解の時間大域適切性と時間減衰評価に関して ..... 10  
 川島秀一 (早大理工)  
 小川卓克 (東北大理)  
 Ryosuke Nakasato (Tohoku Univ.) Global well-posedness and time-decay estimates for the compressible  
 Shuichi Kawashima (Waseda Univ.) Hall-magnetohydrodynamic system  
 Takayoshi Ogawa (Tohoku Univ.)

概要 We study the initial value problem for the compressible hall-magnetohydrodynamic system (hereinafter referred to hall-MHD) in the whole space. We first focus on the solution of the linearized system that are close to some constant state  $(\bar{\rho}, 0, \bar{B})$  with a positive constant  $\bar{\rho}$  and nonzero vector  $\bar{B}$  at infinity. For general systems which include the standard linearized compressible magnetohydrodynamic system, Umeda–Kawashima–Shizuta (1984) obtained the pointwise estimates of solutions in the Fourier space. In this talk, we shall explain the results on global well-posedness and time-decay estimates for the hall-MHD around  $(\bar{\rho}, 0, \bar{B})$ . Furthermore, we shall show the pointwise estimate for the linearized hall-MHD that is the same as the result of Umeda–Kawashima–Shizuta. This is based on the joint work with Profs. Shuichi Kawashima and Takayoshi Ogawa.

- 77 小池 開 (慶大理工・理化学研 AIP) 1次元圧縮性粘性流体中を運動する振り子の長時間挙動について ..... 10  
 Kai Koike (Keio Univ./RIKEN) Long-time behavior of a pendulum in a 1D viscous compressible fluid

概要 We consider the motion of a pendulum in a 1D viscous compressible fluid. The main theorem states that the displacement  $X(t)$  of the pendulum from the equilibrium position decays with the power law  $t^{-3/2}$ . This partially explains the numerically observed long-time behavior of a pendulum in the BGK gas [Tsuji and Aoki, *J. Comput. Phys.*, 250: 574–600, 2013]. The basic tool for the analysis is the pointwise estimates of Green's function as in the case of a freely moving point mass [Koike, arXiv:1904.00992, 2019]. We also need to take into account a cancellation effect due to the oscillatory nature of the problem to obtain the presumably optimal decay rate.

- 78 渡邊 圭市 (早大理工) Global solvability of the Navier–Stokes–Korteweg equations with a non-decreasing pressure in  $L^p$ -framework ..... 10  
 Keiichi Watanabe (Waseda Univ.) Global solvability of the Navier–Stokes–Korteweg equations with a non-decreasing pressure in  $L^p$ -framework

概要 In this talk, we consider the isentropic Navier–Stokes–Korteweg equations with a non-decreasing pressure on the whole space  $\mathbb{R}^d$ ,  $d \geq 2$ , where the system describes the motion of compressible fluids such as liquid-vapor mixtures with phase transitions including a variable internal capillarity effect. We prove the global well-posedness of the system for small initial data belonging to  $H^{s+1,q}(\mathbb{R}^d) \times H^{s,q}(\mathbb{R}^d)^d$  supposing that  $s > d/q$  if  $1 < q \leq d$  and that  $s \geq 1$  if  $d < q < \infty$ . The proof relies on the maximal regularity property for the negative of the Laplace operator.

- 79 鈴木 政尋 (名工大) 気体放電を記述する Morrow モデル I: 安定解析 ..... 10  
 谷 温之 (慶大\*)  
 Masahiro Suzuki (Nagoya Inst. of Tech.) The Morrow model of gas discharge I: Stability analysis  
 Atusi Tani (Keio Univ.\*)

概要 We study an initial boundary value problem to the Morrow model over a bounded interval. This problem has a trivial stationary solution for any voltage, which is adopted as a boundary condition. We show that there exists a threshold of voltage at which the trivial solution becomes from stable to unstable. This threshold can be understood as a sparking voltage for the happening of gas discharge.

- 80 鈴木 政尋 (名工大) 気体放電を記述する Morrow モデル II: 大域分岐解析 ..... 10  
 W. Strauss (Brown Univ.)  
 Masahiro Suzuki (Nagoya Inst. of Tech.) The Morrow model of gas discharge II: Global bifurcation  
 Walter Strauss (Brown Univ.)

概要 We consider the steady states of a gas between two parallel plates that is ionized by a strong electric field so as to create a plasma. We use global bifurcation theory to prove that there is a curve of such states with the following property. The curve begins at the sparking voltage and either the particle density becomes unbounded or the curve ends at the anti-sparking voltage.

- 81 鈴木 政 尋 (名 工 大) プラズマ物理に現れる Boltzmann の関係式の正当化 ..... 10  
 E. Grenier (ENS de Lyon)  
 Yan Guo (Brown Univ.)  
 B. Pausader (Brown Univ.)  
 Masahiro Suzuki (Nagoya Inst. of Tech.) Justification of the Boltzmann relation  
 Emmanuel Grenier (ENS de Lyon)  
 Yan Guo (Brown Univ.)  
 Benoit Pausader (Brown Univ.)

概要 The classical Euler–Poisson system for electrons and ions, interacting through an electrostatic field, describes the motion of plasma. In a certain situation, physicists use a simpler system only for ions by assuming the Boltzmann relation. The simpler system can be derived by letting the mass ratio of an electron and an ion tend to zero. We justify this limit rigorously.

### 16:30～17:30 特別講演

- 瀬 片 純 市 (九 大 数 理) デルタポテンシャルをもつ非線形シュレディンガー方程式の解の長時間挙動  
 Jun-ichi Segata (Kyushu Univ.) Long time behavior of solution to the nonlinear Schrödinger equation with delta potential

概要 We summarize recent progress on long time behavior of solution to the initial-value problem for the one dimensional nonlinear Schrödinger equation with a delta potential. We first consider the case where potential is repulsive and prove that small global solutions decay in  $L^\infty$  and exhibit (modified) scattering. Next we mention the case where potential is attractive and prove that for sufficiently small initial data, the corresponding global solution decomposes into a small solitary wave plus a radiation term that decays and scatters as  $t \rightarrow \infty$ . In particular, we establish the asymptotic stability of the family of small solitary waves.

# 実函数論

3月18日(水) 第VIII会場

9:00~11:50

- 1 川崎敏治(日大工) The split feasibility problem with some projections in Banach spaces  
 眞中裕子(日大短大) ..... 15  
 Toshiharu Kawasaki (Nihon Univ.) The split feasibility problem with some projections in Banach spaces  
 Hiroko Manaka (Nihon Univ.)

概要 In this talk, we treat with the split feasibility problem in some Banach spaces. We show that a solution of this problem is a solution of the equivalent equation defined by using some kinds of projections. Then we show a strong convergence theorem using the method in mathematical programming, in order to find a solution of the split feasibility problem in some Banach spaces.

- 2 富澤佑季乃(新潟工大工) 距離空間における一様凸性 ..... 15  
 Yukino Tomizawa Uniform convexity in distance spaces  
 (Niigata Inst. of Tech.)

概要 It is thought that some distance space which are not linear spaces have properties such that the generalization of properties in linear spaces. However, it remains to be elucidated what geometrical properties exist in the spaces. Here we report the characterization of uniform convexity in Busemann spaces.

- 3 水口洋康(関西大システム理工) Radon plane におけるとある幾何学的定数と von Neumann–Jordan 定数  
 ..... 15  
 Hiroyasu Mizuguchi (Kansai Univ.) A certain geometric constant and von Neumann–Jordan constant in  
 Radon planes

概要 The orthogonality relation in inner product spaces is interesting and investigated by a lot of mathematicians. Moving general normed spaces, there exist many generalized notions of orthogonality. Among these, we treat Birkhoff orthogonality. The orthogonality in inner product spaces is symmetric. However Birkhoff orthogonality in normed spaces is not so in general. If Birkhoff orthogonality is symmetric in a normed space whose dimension is at least three, then the space has a inner product. A two dimensional space in which Birkhoff orthogonality is symmetric is called Radon plane. We study the upper bound of a geometric constant in Radon plane. In addition, using this result, we estimate the von Neumann–Jordan constant in Radon plane.

- 4 斎藤吉助(新潟大理) ラドン空間の構造についての一考察 ..... 15  
 小室直人(北教大旭川)  
 田中亮太郎(東京理大基礎工)  
 Kichi-Suke Saito (Niigata Univ.) A note on the structure of Radon spaces  
 Naoto Komuro  
 (Hokkaido Univ. of Edu.)  
 Ryotaro Tanaka (Tokyo Univ. of Sci.)

概要 Radon spaces are two-dimensional real normed spaces in which Birkhoff orthogonality is symmetric. We present a new characterization of Radon spaces, that is, every Radon space is isometrically isomorphic to special Day–James spaces generated by a pair of an absolute norm and its dual norm.

- 5 川崎 敏治 (日大工・玉川大工) 拡張された積分に関して積分可能な関数 ..... 15  
 Toshiharu Kawasaki Integrable functions for extended integration  
 (Nihon Univ./Tamagawa Univ.)

概要 In this talk, we describe the properties of a family of integrable functions for extended integration.

- 6 福田 亮治 (大分大理工) 単調測度の  $k$ -加法性の 2 つの非離散化 ..... 15  
 本田 あおい (九工大情報工)  
 岡崎 悦明 (フuzzyシステム研)  
 Ryoji Fukuda (Oita Univ.) Two non-discretizations for  $k$ -additivity of a monotone measure  
 Aoi Honda (Kyushu Inst. of Tech.)  
 Yoshiaki Okazaki  
 (Fuzzy Logic Systems Inst.)

概要 The concept of the  $k$ -additive measure is originally defined for a non-additive measure on a finite set. We propose two types of non-discretization for the definition of  $k$ -additive measure: “Constructive  $k$ -additive measure” and “Formulaic  $k$ -additive measure”. We found some their properties and relations.

- 7 中村 昌平 (首都大東京理) The tomography approach to the Fourier restriction theory ..... 15  
 J. Bennett  
 (Univ. of Birmingham)  
 Shohei Nakamura (Tokyo Metro. Univ.) The tomography approach to the Fourier restriction theory  
 Jonathan Bennett  
 (Univ. of Birmingham)

概要 This talk is based on the joint work with Prof. Jonathan Bennett in University of Birmingham. In this talk, we will explain our new approach to the Fourier restriction theory by using the idea of the tomography. This approach leads us to several problems including one sitting between the restriction conjecture and the Keakeya conjecture, and another one which unifies the restriction conjecture and the Mizohata–Takeuchi conjecture (when dimension 2). We also give some applications of our approach to the weight theory for the Fourier extension operator.

- 8 飯田 毅士 (福島工高専) Weighted norm inequalities on Morrey spaces for the Orlicz-fractional  
 maximal operators ..... 15  
 Takeshi Iida Weighted norm inequalities on Morrey spaces for the Orlicz-fractional  
 (Fukushima Nat. Coll. of Tech.) maximal operators

概要 In this talk, we discuss the weighted norm inequalities on Morrey spaces for the Orlicz-fractional maximal operators. We have investigated the boundedness of the weighted Lebesgue spaces and Morrey spaces for the Orlicz-fractional maximal operators and weighted estimates for the fractional integral and maximal operators in Morrey spaces. The main results which are based on these investigations give the estimates for the Orlicz-fractional maximal operators in Morrey spaces.

- 9 川澄 亮太 Generalized fractional integral operators on weak Orlicz spaces ..... 15  
 中井 英一 (茨城大理)  
 Ryota Kawasumi Generalized fractional integral operators on weak Orlicz spaces  
 Eiichi Nakai (Ibaraki Univ.)

概要 We consider the boundedness of the generalized fractional integral operator from a weak Orlicz space to another weak Orlicz space. We also consider some other operators.

- 10 石 明 磊 (茨城大理工) Commutators of Calderón–Zygmund and generalized fractional integral  
 新井龍太郎 (茨城大理工) operators with functions in generalized Campanato spaces on Orlicz–  
 中井英一 (茨城大理工) Morrey spaces ..... 15  
 Minglei Shi (Ibaraki Univ.) Commutators of Calderón–Zygmund and generalized fractional integral  
 Ryutaro Arai (Ibaraki Univ.) operators with functions in generalized Campanato spaces on Orlicz–  
 Eiichi Nakai (Ibaraki Univ.) Morrey spaces

概要 Let  $\mathbb{R}^n$  be the  $n$ -dimensional Euclidean space. Let  $b \in \text{BMO}(\mathbb{R}^n)$  and  $T$  be a Calderón–Zygmund singular integral operator. In 1976 Coifman, Rochberg and Weiss proved that the commutator  $[b, T] = bT - Tb$  is bounded on  $L^p(\mathbb{R}^n)$  ( $1 < p < \infty$ ), that is,  $\|[b, T]f\|_{L^p} = \|bTf - T(bf)\|_{L^p} \leq C\|b\|_{\text{BMO}}\|f\|_{L^p}$ , where  $C$  is a positive constant independent of  $b$  and  $f$ . For the fractional integral operator  $I_\alpha$ , Chanillo proved the boundedness of  $[b, I_\alpha]$  in 1982. These results were extended to Orlicz spaces by Fu, Yang and Yuan (2012, 2014). In this talk we discuss the boundedness of the commutator  $[b, I_\rho]$ ,  $[b, I_T]$  on Orlicz–Morrey spaces, where  $T$  is a Calderón–Zygmund operator,  $I_\rho$  is a generalized fractional integral operator and  $b$  is a function in generalized Campanato spaces.

#### 14:15~15:55

- 11 岩 瀨 司 (東北大理) Dirichlet および Neumann Laplacian に付随する Sobolev 空間における  
 積の評価式 ..... 15  
 Tsukasa Iwabuchi (Tohoku Univ.) Bilinear estimates in Sobolev spaces associated with Dirichlet and Neu-  
 mann Laplacian

概要 We study the bilinear estimates in the Sobolev spaces with the Dirichlet and the Neumann condition. The optimal regularity will be revealed to obtain such estimates in the half space case.

- 12 国 定 亮 一 (早大教育) Banach 極限の連続変数版について ..... 15  
 Ryoichi Kunisada (Waseda Univ.) On a continuous version of Banach limits

概要 Banach limits are normalized positive bounded linear functionals on  $l_\infty$  of the set of bounded functions on natural numbers. We consider a cotinuous analogue of Banach limits defined on  $L^\infty(\mathbb{R})$  and give a characterization of them in terms of invariance with respcet to convolution operators. We also discuss some applications.

- 13 野 ケ 山 徹 (首都大東京理) Local Muckenhoupt class for variable exponents ..... 15  
 澤 野 嘉 宏 (首都大東京理)  
 Toru Nogayama (Tokyo Metro. Univ.) Local Muckenhoupt class for variable exponents  
 Yoshihiro Sawano (Tokyo Metro. Univ.)

概要 In this paper, we define  $A_{p(\cdot)}^{loc}$  and show that the weighted inequality for local Hardy–Littlewood maximal operator on the Lebesgue spaces with variable exponent. This work will extend the theory of Rychkov, who developed the theory of  $A_p^{loc}$  weights. Due to the setting of variable exponents, a new method of extension of weights will be needed; the extension method is different from the one by Rychkov.

- 14 澤 野 嘉 宏 (首都大東京理)<sup>b</sup> Cantor functions associated with generalized expansions ..... 15  
 Yoshihiro Sawano (Tokyo Metro. Univ.) Cantor functions associated with generalized expansions

概要 The goal of this talk is to consider expansions generated by a sequence  $\{h_n\}_{n=1}^\infty$ , where  $h_n \in \mathbb{N} \cap [2, \infty)$  for each  $n$ . We consider the generalized Cantor function and discuss its non-differentiable points.

- 15 澤 野 嘉 宏 (首都大東京理)<sup>b</sup> Sparse non-smooth atomic decomposition of Morrey spaces ..... 15  
 Yoshihiro Sawano (Tokyo Metro. Univ.) Sparse non-smooth atomic decomposition of Morrey spaces

概要 The goal of this talk is to refine the existing result on the decomposition of functions in Morrey spaces. As an application, we revisit the equivalence obtained by Adams.

- 16 澤野嘉宏 (首都大東京理)<sup>b</sup> Modified Hardy–Littlewood maximal operator and modified fractional  
 下村哲 (広島大教育) integral operator on metric measure spaces ..... 15  
 Yoshihiro Sawano (Tokyo Metro. Univ.) Modified Hardy–Littlewood maximal operator and modified fractional  
 Tetsu Shimomura (Hiroshima Univ.) integral operator on metric measure spaces

概要 The goal of this talk is to simplify the example given by the first author in 2005. Recently Stempak constructed a simpler example showing that many results in a metric measure spaces fails unless they are modified suitably. Based on the idea of Stempak, we provide some further examples.

#### 16:15~17:15 特別講演

- 貞末岳 (大阪教育大数学教育) Martingale 空間と, martingale に対する分数べき積分  
 Gaku Sadasue (Osaka Kyoiku Univ.) Some martingale spaces and fractional integrals for martingale

概要 In recent years, various new martingale spaces are introduced and structures of these spaces are studied. In this talk, we give some recent results on several martingale spaces. We also give some results on the boundedness of martingale transforms on these spaces. We especially study a special class of martingale transforms called fractional integrals.

3月19日(木) 第VIII会場

#### 9:00~12:00

- 17 水上雅昭 (東京理大理) Uniform-in-time convergence of solutions for a chemotaxis-competition  
 model to those for the Lotka–Volterra competition model ..... 15  
 Masaaki Mizukami Uniform-in-time convergence of solutions for a chemotaxis-competition  
 (Tokyo Univ. of Sci.) model to those for the Lotka–Volterra competition model

概要 This work is concerned with the question that “how far does small chemotactic interaction perturb the Lotka–Volterra competition dynamics?”. A two-species chemotaxis-competition model was studied by e.g., Bai–Winkler (2016) and Lin–Mu–Wang (2015). However, there are still many open problems about the two-species chemotaxis-competition model. On the other hand, the Lotka–Volterra competition model has been studied extensively. Thus the development of this work will enable us to see new properties of solutions for the chemotaxis system. The main result of this talk gives uniform-in-time convergence of solutions for the two-species chemotaxis-competition system to those for the Lotka–Volterra competition model.

- 18 P. Colli (Univ. of Pavia) Existence for a phase separation system deduced from the entropy bal-  
 来間俊介 (東京理大理) ance ..... 15  
 Pierluigi Colli (Univ. of Pavia) Existence for a phase separation system deduced from the entropy bal-  
 Shunsuke Kurima (Tokyo Univ. of Sci.) ance

概要 This talk is concerned with a phase separation system deduced from the entropy balance. Both the viscous and the non-viscous cases are considered in the Cahn–Hilliard relations characterizing the phase dynamics. The entropy balance is written in terms of the absolute temperature and of its logarithm, appearing under time derivative. The initial and boundary value problem is considered for the system of partial differential equations. The existence of a global solution is proved via some approximations involving Yosida regularizations and a suitable time discretization.

- 19 来間俊介 (東京理大理) A Cahn–Hilliard approach to a nonlinear diffusion chemotaxis system ..... 15  
 Shunsuke Kurima (Tokyo Univ. of Sci.) A Cahn–Hilliard approach to a nonlinear diffusion chemotaxis system

概要 This talk deals with a nonlinear diffusion chemotaxis system. Colli–Fukao (2015) proved existence of solutions to a Cahn–Hilliard system as an approximation of a nonlinear diffusion equation by applying the abstract theory by Colli–Visintin (1990) for doubly nonlinear evolution inclusions with some bounded monotone operator and some proper lower semicontinuous convex function. Moreover, Colli–Fukao (2016) established existence of solutions to the nonlinear diffusion equation by passing to the limit in the Cahn–Hilliard equation. However, Cahn–Hilliard approaches to nonlinear diffusion chemotaxis systems seem not to be studied. This talk will try to derive existence of solutions to a nonlinear diffusion chemotaxis system by passing to the limit in a Cahn–Hilliard type chemotaxis system.

- 20 香川溪一郎 (早大理工) Asymptotic limits of the time-periodic problem for the viscous Cahn–  
 大谷光春 (早大理工) Hilliard equation ..... 10  
 Keiichiro Kagawa (Waseda Univ.) Asymptotic limits of the time-periodic problem for the viscous Cahn–  
 Mitsuharu Ôtani (Waseda Univ.) Hilliard equation

概要 We consider the asymptotic limits of the time-periodic problem for the viscous Cahn–Hilliard equation with the homogeneous Dirichlet boundary condition. We already reported about the asymptotic limits of the initial value problem for the viscous Cahn–Hilliard equation. It is shown that the Cahn–Hilliard equation and the Allen–Cahn equation are derived as the asymptotic limit of the viscous Cahn–Hilliard equation and hence we show the existence of the time-periodic solutions of their equations.

- 21 小杉千春 (日本女大理) 弾性体の収縮運動を記述する初期値境界値問題の弱解の存在について .. 15  
 愛木豊彦 (日本女大理)  
 Chiharu Kosugi (Japan Women’s Univ.) Existence of weak solutions to initial boundary value problems describ-  
 Toyohiko Aiki (Japan Women’s Univ.) ing shrinking motion of elastic materials

概要 In this talk we consider existence of weak solutions to an initial boundary value problem for beam equations. This problem describes a shrinking motion of elastic materials whose examples are rubber bands. The aims of this talk are to show an existence theorem of its weak solution and to present some numerical results for the approximation problems given as systems of ordinary differential equations. Also, we establish existence and unique of a solution to the system of ODE’s by the fixed point theorem and convergence of numerical solutions.

- 22 岸田崇裕 (名城大理工) 水分吸着過程を記述する 1 次元モデルの有限要素法による解析 ..... 15  
 村瀬勇介 (名城大理工)  
 Takahiro Kishida (Meijo Univ.) FEM analysis for mathematical model of adsorption phenomena in 1D  
 Yusuke Murase (Meijo Univ.) domain

概要 We study a mathematical model of adsorption phenomena in 1D domain. The model is free boundary problem which represent moisture adsorption phenomena in porous medium like a concrete. From recently researches, we got existence of unique time-local and global solution, existence of periodic solutions, large time behavior of solutions, and some numerical simulations. However, these numerical simulations are done with experimentally technics. So now we simulate and analyze this model by Free Element Method with Adaptive Moving Mesh Method. In this talk, we show some new numerical experiments and mathematical results.



- 23 紫村 一輝 (大分大工) 相分離を伴う弾性方程式の構造保存型差分法とその誤差評価 ..... 15  
吉川 周二 (大分大理工)  
Kazuki Shimura (Oita Univ.) Structure-preserving finite difference schemes for a Cahn–Hilliard system coupled with elasticity  
Shuji Yosikawa (Oita Univ.)

概要 This study was made to observe the behavior of the solution by numerical analysis on the Cahn–Hilliard system coupled with elasticity (CHE), which is one of nonlinear partial differential equations, and summarizes the results obtained by the solution and error estimate includes the case where the viscosity coefficient is zero. CHE is a system of 4th order evolution equations for two unknowns describing a phenomenon in which a substance having an elastic property such as a phase separation. Although mathematical proofs of the existence and uniqueness of the solution are given, it has not been clarified the behavior of solution yet. Therefore, in this study, we demonstrate numerical simulations and error estimate. For the purpose, we use structure-preserving numerical methods that is expected to be stable and accurate.

- 24 奥村 真善美 (阪大情報) 総質量保存則を持つ動的境界条件下の Cahn–Hilliard 方程式に対する構造保存スキーム ..... 15  
Makoto Okumura (Osaka Univ.) A structure-preserving scheme for the Cahn–Hilliard equation with dynamic boundary conditions which has the total mass conservation

概要 We propose a structure-preserving scheme for the GMS model by using the discrete variational derivative method (DVDM). In the model, two characteristic properties hold. One is the total mass conservation, which means that the sum of the mass in bulk and on the boundary is conserved. The other is the total energy dissipation, which represents the sum of energy in bulk and on the boundary decrease. In this study, we design a finite difference scheme for the GMS model so that the scheme inherits the properties from the original problem in a discrete sense. In this talk, we focus on the existence and uniqueness of the solution for the scheme.

- 25 佐々木 善雅 (新潟大自然) 空間的に不連続な流束をもつ保存則方程式の一意可解性について ..... 15  
應和 宏樹 (新潟大自然)  
Yoshimasa Sasaki (Niigata Univ.) Existence and uniqueness of solutions to conservation laws with spatially discontinuous flux  
Ohwa Hiroki (Niigata Univ.)

概要 We study the existence and uniqueness of solutions to conservation laws with spatially discontinuous flux functions.

- 26 渡邊 紘 (大分大理工) 放物型・双曲型単独保存則に対する進行波の構成とエントロピー解の漸近挙動 ..... 15  
Hiroshi Watanabe (Oita Univ.) Construction of traveling waves and asymptotic behavior of entropy solutions to scalar parabolic-hyperbolic conservation laws

概要 We consider one-dimensional Cauchy problems (CP) for scalar parabolic-hyperbolic conservation laws. The equation is regarded as a linear combination of the hyperbolic conservation laws and the porous medium type equations. Thus, this equation has both properties of hyperbolic equations and those of parabolic equations. Accordingly, it is difficult to investigate the behavior of solutions to (CP). In this talk, we construct discontinuous traveling waves and discuss the properties of them. Moreover, we show the asymptotic behavior of entropy solutions to (CP) using the constructed traveling waves.

- 27 都 築 寛 (広島修道大経済) Existence for Initial-boundary value problems for Vlasov–Poisson equations with angle error in magnetic field ..... 15  
 Yutaka Tsuzuki (Hiroshima Shudo Univ.) Existence for Initial-boundary value problems for Vlasov–Poisson equations with angle error in magnetic field

概要 We deal with initial-boundary problems for Vlasov–Poisson systems in a half-space. In 2013, Skubachevskii provides local-in-time solvability to the system. Furthermore, in 2017, existence result with weaker condition were also obtained where the magnetic force is horizontal to the wall. This talk provides another result for the equation where the magnetic force has angle error in the vertical direction and depending on the first element of the spatial variable.

#### 14:15~15:55

- 28 山崎 教昭 (神奈川大工) Control of parameter-dependent evolution equations governed by time-dependent subdifferentials ..... 15  
 剣持 信幸 (千葉大\*)  
 白川 健 (千葉大教育)  
 Noriaki Yamazaki (Kanagawa Univ.) Control of parameter-dependent evolution equations governed by time-dependent subdifferentials  
 Nobuyuki Kenmochi (Chiba Univ.\*)  
 Ken Shirakawa (Chiba Univ.)

概要 We consider nonlinear parameter-dependent evolution equations governed by double time-dependent subdifferentials in uniformly convex Banach spaces. In this talk, we investigate singular optimal control problems for doubly nonlinear parameter-dependent evolution state equations. Then, we show the existence of an optimal control for our problem.

- 29 久保田 翔大 (千葉大融合理工) 一般多次元での時間離散型 Kobayashi–Warren–Carter システムに対する最適制御問題 ..... 15  
 白川 健 (千葉大教育)  
 Shodai Kubota (Chiba Univ.) Optimal control problem for multidimensional semi-discrete system of Kobayashi–Warren–Carter type  
 Ken Shirakawa (Chiba Univ.)

概要 We consider a class of optimal control problems for state problems of multidimensional semi-discrete systems. Each state problem is denoted by  $(S)_\varepsilon$ , with  $\varepsilon > 0$ , and is based on the phase-field model of grain boundary motion. In this regard, each optimal control problem is denoted by  $(OP)_\varepsilon$ , with  $\varepsilon > 0$ , and it is prescribed as a minimization problem of a cost function. Additionally, the problems  $(S)_\varepsilon$  and  $(OP)_\varepsilon$  are supposed to admit limiting profiles as  $\varepsilon \downarrow 0$ , and then, the limiting problems are supposed to contain no little singularities. The main interest is in the case when  $\varepsilon > 0$  (regular case), and the mathematical results concerned with: (A) the existence of the optimal control when  $\varepsilon > 0$ ; (B) the necessary condition for the regular optimal control; (C) limiting observation as  $\varepsilon \downarrow 0$ ; will be reported as the main theorems of this talk.

- 30 白川 健 (千葉大教育) Sufficient condition for the existence of one-dimensional crystalline solution of the Kobayashi–Warren–Carter type system ..... 15  
 渡邊 紘 (大分大理工)  
 Ken Shirakawa (Chiba Univ.) Sufficient condition for the existence of one-dimensional crystalline solution of the Kobayashi–Warren–Carter type system  
 Hiroshi Watanabe (Oita Univ.)

概要 In this talk, we consider a one-dimensional Kobayashi–Warren–Carter type system, which is based on a phase-field model of grain boundary motion. This talk corresponds to a continuation of the last presentation at MSJ Spring Meeting 2019 (Tokyo), which was concerned with the uniqueness of special kind of solution, named “crystalline solution”. On this basis, we here focus on the existence issue of crystalline solution. Through the precise observations for solutions to a time-discretization scheme, a class of structural conditions for the initial data will be presented as the sufficient condition for the existence of crystalline solution.

- 31 熊崎 耕太 (長崎大教育) 氷膜形成に関連するある1次元自由境界問題について ..... 15  
 Kota Kumazaki (Nagasaki Univ.) A one-dimensional free boundary problem related to ice lenses formation

概要 In cold regions, buildings that are exposed to extremely low temperatures undergo freezing and build microscopic ice lenses that lead to the mechanical damage of the material. In this talk, we consider a mathematical model describing swelling of a pocket of water to understand the formation of ice lenses growing inside of porous materials. Our problem is posed on a halfline with a moving boundary at one of the ends, and the moving boundary conditions encode the swelling mechanism, while a diffusion equation is responsible for providing water content for the swelling to take place. In this talk, we discuss the existence and uniqueness, large time behavior of a solution to our problem.

- 32 深尾 武史 (京都教育大) 動的境界条件下でのCahn–Hilliard方程式系に対する境界拡散項の消滅  
 P. Colli ( Pavia Univ. ) について ..... 15  
 Takeshi Fukao (Kyoto Univ. of Edu.) Vanishing diffusion in a dynamic boundary condition for the Cahn–  
 Pierluigi Colli (Pavia Univ.) Hilliard equation

概要 In this talk, we will treat the Cahn–Hilliard equation with a dynamic boundary condition of Allen–Cahn type. We focus on the analysis of the surface diffusion on the dynamic boundary condition. By the asymptotic analysis, we can expect that the solution with the surface diffusion converges to the one of without the surface diffusion in a sense. The role of the surface diffusion will be expressed by means of the classes of the solutions.

- 33 伊藤 昭夫 Approach from the quasi-variational structure to tumor invasion with  
 non-smooth degenerate diffusion ..... 15  
 Akio Ito Approach from the quasi-variational structure to tumor invasion with  
 non-smooth degenerate diffusion

概要 We consider an initial-boundary value problem of a tumor invasion of Chaplain–Anderson type, in which the coefficient of random motility of tumor cells depends on the extracellular matrix and the diffusion flux of tumor cells is nonsmooth and degenerate. From these points of view, the Cauchy problem considered in this talk has a quasi-variational structure and this fact makes it more difficult and complicate to analyze this model mathematically. The aim of this talk, we give the existence of local-in-time solutions by applying the general theory, which was established in the paper, Evolution inclusion on a real Hilbert space with quasi-variational structure for inner products, Journal of Convex Analysis, 26 (2019), No. 4, 1185–1252.

#### 16:15～17:15 特別講演

- 高 棹 圭 介 <sup>b</sup> フェイズフィールド法による外力項付き平均曲率流方程式の弱解の存在  
 (京大理・京大白眉センター) について  
 Keisuke Takasao On the existence of the weak solution for the mean curvature flow with  
 (Kyoto Univ./Kyoto Univ.) forcing term via the phase field method

概要 We study the mean curvature flow with given non-smooth forcing term  $g$ . In 1993, Ilmanen proved the existence of the Brakke flow with  $g \equiv 0$ , by using the phase field method. Generally, the most difficult part of the proof of the existence theorem is the estimate of the positive part of the discrepancy measure. To solve the problem, Ilmanen showed the non-positivity of the discrepancy measure via the maximum principle. However, in the case of  $g \neq 0$ , the property does not hold for the usual phase field method of the problem. In this talk, we explain a new modified Allen–Cahn equation which satisfies the non-positivity of the discrepancy measure, and we prove the existence of the weak solution for the mean curvature flow with forcing term in suitable Sobolev spaces.

# 函数解析学

3月16日(月) 第VII会場

10:00~11:45

- 1 清瀬 周 (神戸大理) Floquet ハミルトニアンに対する Mourre 評価について ..... 15  
足立 匡義 (京大人間環境)  
Amane Kiyose (Kobe Univ.) On the Mourre estimates for Floquet Hamiltonians  
Tadayoshi Adachi (Kyoto Univ.)

概要 We introduce a new conjugate operator for a Floquet Hamiltonian associated with a Schrödinger operator with time-periodic potentials. Although Yokoyama (1998) already obtained a conjugate operator for the Floquet Hamiltonian, there was a difficulty that it won't afford extensions to the many-body systems. Here we introduce our new conjugate operator which suggests such extensions. Actually, combining our results and Yokoyama's, Adachi recently constructed a conjugate operator for three-body systems.

- 2 川越 大輔 (京大情報) Surface Riesz transforms and spectral property of the elastic Neumann–  
Hyeonbae Kang (Inha Univ.) Poincaré operator on less smooth domains in three dimensions ..... 15  
Daisuke Kawagoe (Kyoto Univ.) Surface Riesz transforms and spectral property of the elastic Neumann–  
Hyeonbae Kang (Inha Univ.) Poincaré operator on less smooth domains in three dimensions

概要 It was proved that the elastic Neumann–Poincaré operator, which is a boundary integral operator related to the Lamé system, is polynomially compact and, as a consequence, that its spectrum consists of three non-empty sequences of eigenvalues accumulating to certain numbers determined by Lamé constants, if the boundary of the domain where the operator is defined is  $C^\infty$ -smooth. We extend this result to less smooth boundaries, namely,  $C^{1,\alpha}$ -smooth boundaries for some  $\alpha > 0$ . The results are obtained by proving certain identities for surface Riesz transforms, which are singular integral operators of non-convolution type, defined by the metric tensor on a given surface.

- 3 川越 大輔 (京大情報) The essential spectrum of the elastic Neumann–Poincaré operator on a  
E. Bonnetier planar domain with a corner ..... 15  
(Univ. Grenoble-Alpes)  
C. Dapogny  
(Univ. Grenoble-Alpes)  
Hyeonbae Kang (Inha Univ.)  
Daisuke Kawagoe (Kyoto Univ.) The essential spectrum of the elastic Neumann–Poincaré operator on a  
Eric Bonnetier (Univ. Grenoble-Alpes) planar domain with a corner  
Charles Dapogny  
(Univ. Grenoble-Alpes)  
Hyeonbae Kang (Inha Univ.)

概要 Main results of this talk is a contribution to the study of the spectrum of the elastic Neumann–Poincaré operator, which is a boundary integral operator involved in the resolution of the Lamé system in a domain. Previous studies have shown that when the domain is smooth in two dimensions, the spectrum of the operator consists of two sequences of eigenvalues with accumulation points  $\pm\kappa_0$ , where the constant  $\kappa_0$  is explicitly determined by the Lamé constants. In this talk, we focus on the situation where the planar domain is smooth except at a corner; we then prove that the elastic Neumann–Poincaré operator has essential spectrum, which contains at least two intervals around the values  $\kappa_0$  and  $-\kappa_0$  respectively.

- 4 井上 寛 (第一薬大) 非自己共役ハミルトニアンから生成される量子力学系 ..... 15  
Hiroshi Inoue (Daiichi Univ. of Pharm.) Quantum dynamics based on non-self-adjoint hamiltonians

概要 This presentation introduces a study of quantum dynamics generated from non-self-adjoint hamiltonian based on biorthogonal sequences in Hilbert space. In this study, the notion of generalized Riesz systems plays an important role. We investigate under what conditions standard self-adjoint hamiltonian and non-self-adjoint hamiltonian defined by generalized Riesz systems construct quantum dynamics. Furthermore, we investigate under what conditions generalized Riesz systems define gibbs states on a  $*$ -algebra of unbounded operators.

- 5 岩田 順 敬 (関西大化学生命工) 作用素の対数表現を用いた抽象 Miura 変換 ..... 15  
Yoritaka Iwata (Kansai Univ.) Abstract Miura transform based on the logarithmic representation of operators

概要 Miura transform is known as the transformation between Korteweg–de Vries(KdV) and modified KdV (mKdV) equations. In this talk, based on the logarithmic representation of operators in Banach spaces, a structure common to both Miura and Cole–Hopf transforms is studied in an abstract manner. In conclusion, by means of the abstract version of the Miura transform in Banach spaces, an integrable class of nonlinear semigroup theory is clarified. It is the first step to pin down the correspondence of Backland transform in semigroup theory of operators.

- 6 渡辺 秀 司 (群馬大理工) 超伝導の BCS-Bogoliubov モデルにおける超伝導体の比熱の作用素論的研究 ..... 15  
Shuji Watanabe (Gunma Univ.) An operator-theoretical treatment of the specific heat of a superconductor in the BCS-Bogoliubov model of superconductivity

概要 In this talk we study the temperature dependence of the specific heat in the BCS-Bogoliubov model of superconductivity from the viewpoint of operator theory. We give the exact and explicit expression for  $\Delta C_V(T_c)/C_V^N(T_c)$ . Here,  $C_V^N(T_c)$  denotes the specific heat at constant volume at the transition temperature  $T_c$ , and  $\Delta C_V(T_c)$  its gap at  $T_c$ . We then show that  $\Delta C_V(T_c)/C_V^N(T_c)$  does not depend on superconductors and is a universal constant.

#### 14:15~15:15 2019年度(第18回)日本数学会解析学賞受賞特別講演

- 廣島 文 生 (九大数理) 汎関数積分によるくりこみ理論と基底状態の非摂動的解析について  
Fumio Hiroshima (Kyushu Univ.) Renormalization theory and non-perturbative analysis of ground states by functional integrations

#### 15:30~16:30 特別講演

- 中野 史 彦 (学習院大理) 1次元ランダムシュレーディンガー作用素の固有値・固有関数のスケール  
ング極限について  
Fumihiko Nakano (Gakushuin Univ.) Scaling limit of the eigenvalues and eigenfunctions of 1-dimensional random Schrödinger operators

概要 1-dimensional random Schrödinger operators have various spectral properties, and level statistics depending on the decay rate of the potential at infinity. Especially, the level statistics at the critical order is intimately related to the scaling limit of the Gaussian beta ensemble in the random matrix theory. In this talk, we discuss the joint scaling limit of the eigenvalues and eigenfunctions.

## 3月17日(火) 第VII会場

10:30~11:45

- 7 川村 晃 英 (京大 理) 球表現の分解と多変数超幾何型多項式の加法定理 ..... 15  
 Koei Kawamura (Kyoto Univ.) Decomposition of spherical representations and an addition theorem for multivariate hypergeometric polynomials

概要 Harmonic analysis on groups is useful to study properties of special functions which arise as zonal spherical functions. Dunkl (1976) induced an addition theorem for the Krawtchouk polynomials, which are discrete orthogonal polynomials of hypergeometric type, by decomposing spherical representations of wreath products of symmetric groups. We apply his method to multivariate case, that is, we use harmonic analysis on a non-Archimedean local field with a wreath products action. Then we have an addition theorem for multivariate Krawtchouk polynomials with coefficients in multivariate Hahn polynomials.

- 8 嵐 晃 一 (名大 多元数理) 有界等質領域上の holomorphic multiplier 表現 ..... 15  
 Koichi Arashi (Nagoya Univ.) Holomorphic multiplier representations for bounded homogeneous domains

概要 We consider the unitarizations in the spaces of holomorphic sections of equivariant holomorphic line bundles over a bounded homogeneous domain under the action of the identity component of an algebraic group acting transitively on the domain. We give a complete classification of unitary representations arising from such unitarizations.

- 9 田内 大 渡 (東大 数理) A generalization of the uniformly bounded multiplicity theorem ..... 15  
 Taito Tauchi (Univ. of Tokyo) A generalization of the uniformly bounded multiplicity theorem

概要 Let  $H$  be a closed subgroup of a real reductive Lie group  $G$  and  $G_{\mathbb{C}}, H_{\mathbb{C}}$  their complexifications. It was proved by T. Kobayashi and T. Oshima that the multiplicities of irreducible representations of  $G$  in the regular representation  $C^{\infty}(G/H)$  are uniformly bounded iff  $H_{\mathbb{C}}$  acts on the full flag variety  $G_{\mathbb{C}}/B$  with finitely many orbits. In this talk, we prove that the multiplicities of the irreducible representations of  $G$  induced from a parabolic subgroup  $Q$  in  $C^{\infty}(G/H)$  are uniformly bounded if  $H_{\mathbb{C}}$  acts on  $G_{\mathbb{C}}/Q_{\mathbb{C}}$  with finitely many orbits using the theory of holonomic  $\mathcal{D}_X$ -modules.

- 10 久保 利 久 (龍谷大 経済) The  $K$ -type formulas for Kable's differential operators of type  $A_3$  and  
 B. Ørsted (Aarhus Univ.) Heun polynomials ..... 15  
 Toshihisa Kubo (Ryukoku Univ.) The  $K$ -type formulas for Kable's differential operators of type  $A_3$  and  
 Bent Ørsted (Aarhus Univ.) Heun polynomials

概要 In 2012, Kable introduced a one-parameter family  $\mathcal{D}_n(s)$  with  $s \in \mathbb{C}$  of differential operators for  $\mathfrak{sl}(n, \mathbb{C})$ . The operators  $\mathcal{D}_n(s)$  can be thought of as  $\widetilde{SL}(n, \mathbb{R})$ -intertwining differential operators between parabolically induced representations; thus, the space  $Sol_n(s)$  of smooth solutions for  $\mathcal{D}_n(s)$  is naturally a representation space of  $\widetilde{SL}(n, \mathbb{R})$ . To date, the representations realized on  $Sol_n(s)$  are determined only for the case  $(n, s) = (3, 0)$ , for which the realized representations are precisely all minimal unitary representations of  $\widetilde{SL}(n, \mathbb{R})$  ([Kubo-Ørsted, 2019]). In this talk we then consider the case  $(n, s) = (3, \text{general})$ . We first classify the complex parameters  $s \in \mathbb{C}$  with  $Sol_3(s) \neq \{0\}$ . We then determine the  $K$ -type formulas for the space  $Sol_3(s)_K$  of  $K$ -finite solutions for  $\mathcal{D}_3(s)$  by using Heun polynomials.

## 13:15~14:15 特別講演

中濱良祐 (東大数理) 正則離散系列表現の制限に関する絡作用素の構成

Ryosuke Nakahama (Univ. of Tokyo) Construction of intertwining operators for restriction of holomorphic discrete series representations

概要 Let  $G$  be a Lie group,  $G' \subset G$  be a closed subgroup, and we consider a unitary representation  $\mathcal{H}$  of  $G$ . Then in general, the restriction  $\mathcal{H}|_{G'}$  decomposes into a direct integral of irreducible representations of  $G'$ . Especially, if  $\mathcal{H}$  is in a nice class of representations called “holomorphic discrete series representations” and  $(G, G')$  is a “symmetric pair of holomorphic type”, then  $\mathcal{H}|_{G'}$  decomposes discretely, and there exist  $G'$ -intertwining operators between  $\mathcal{H}|_{G'}$  and a representation  $\mathcal{H}'$  of the subgroup  $G'$  of both directions. The projection operator  $\mathcal{H}|_{G'} \rightarrow \mathcal{H}'$  (the symmetry breaking operator) is always given by a differential operator, and the embedding operator  $\mathcal{H}' \rightarrow \mathcal{H}|_{G'}$  (the holographic operator) is given by an infinite-order differential operator. In this talk the speaker gives some results on explicit construction of these intertwining operators in some examples.

3月18日(水) 第VII会場

## 9:45~12:00

11 伊佐浩史 (前橋工科大) The  $n$ -th Petz–Bregman divergence and the  $n$ -th residual relative operator entropy ..... 15  
 亀井栄三郎

遠山宏明 (前橋工科大)  
 渡邊雅之 (前橋工科大)

Hiroshi Isa (Maebashi Inst. of Tech.) The  $n$ -th Petz–Bregman divergence and the  $n$ -th residual relative operator entropy  
 Eizaburo Kamei

Hiroaki Tohyama  
 (Maebashi Inst. of Tech.)

Masayuki Watanabe  
 (Maebashi Inst. of Tech.)

概要 Let  $A$  and  $B$  be strictly positive operators on a Hilbert space,  $n \in \mathbb{N}$  and  $x, y \in \mathbb{R}$ . We have defined the  $n$ -th residual relative operator entropy  $\mathfrak{R}_{x,y}^{[n]}(A|B) \equiv A^{\frac{1}{2}} \Psi^{[n]}(A^{-\frac{1}{2}} B A^{-\frac{1}{2}}, x, y) A^{\frac{1}{2}}$ , where  $\Psi^{[n]}(a, x, y) = \frac{a^y (\log a)^n}{(n-1)!} \int_0^1 (1-t)^{n-1} a^{t(x-y)} dt$  ( $a > 0$ ) is the residual term of the Taylor expansion of  $a^x$  around  $y$  divided by  $(x-y)^n$ . By using  $\mathfrak{R}_{x,y}^{[n]}(A|B)$ , we can show the relations among the  $n$ -th relative operator entropies or more precise properties of them.

12 伊藤公智 (前橋工科大) Furuta type inequalities related to Ando–Hiai inequality with negative powers ..... 10  
 亀井栄三郎

Masatoshi Ito (Maebashi Inst. of Tech.) Furuta type inequalities related to Ando–Hiai inequality with negative powers  
 Eizaburo Kamei

概要 Recently, Kian and Seo had shown the Ando–Hiai type inequality with negative powers, and also Fujii and Nakamoto have given the Furuta type inequality with negative powers and its generalizations. In this talk, we show some improvement forms of their Furuta type inequalities based on properties of Furuta inequality.

- 13 東條玲央 (大阪教育大教育) Matrix Hölder–McCarthy inequality via matrix geometric means . . . . . 10  
瀬尾祐貴 (大阪教育大教育)  
中山棕介 (大阪教育大教育)

Reo Tojo (Osaka Kyoiku Univ.) Matrix Hölder–McCarthy inequality via matrix geometric means  
Yuki Seo (Osaka Kyoiku Univ.)  
Ryosuke Nakayama  
(Osaka Kyoiku Univ.)

**概要** In this talk, by virtue of an expression of matrix geometric means for positive semidefinite matrices via the Moore–Penrose inverse, we show matrix versions of the Hölder–McCarthy inequality and quasi-arithmetic power means via matrix geometric means.

- 14 中山棕介 (大阪教育大教育) Reverse matrix quasi-arithmetic power means via matrix geometric means  
瀬尾祐貴 (大阪教育大教育) . . . . . 10  
東條玲央 (大阪教育大教育)

Ryosuke Nakayama Reverse matrix quasi-arithmetic power means via matrix geometric means  
(Osaka Kyoiku Univ.)  
Yuki Seo (Osaka Kyoiku Univ.)  
Reo Tojo (Osaka Kyoiku Univ.)

**概要** In this talk, by virtue of an expression of matrix geometric means for positive semidefinite matrices via the Moore–Penrose inverse, we show matrix versions of the Hölder–McCarthy inequality, the Hölder inequality and quasi-arithmetic power means via the matrix geometric means, and their reverses for positive definite matrices via the generalized Kantorovich constant

- 15 瀬尾祐貴 (大阪教育大教育) Norm inequalities for deformed operator means . . . . . 10  
Yuki Seo (Osaka Kyoiku Univ.) Norm inequalities for deformed operator means

**概要** In this talk, for an  $n$ -tuple of positive invertible operators on a Hilbert space, we investigate the monotonicity of the power mean from the deformed operator mean from an  $n$ -variable operator mean by an operator mean in terms of the generalized Kantorovich constants under the operator order. Moreover, we improve the norm inequality for the operator power means related to the Log-Euclidean mean in terms of the Specht ratio.

- 16 藤井淳一 (大阪教育大教育) 同ランク正半定値の行列平均について . . . . . 15  
Junichi Fujii (Osaka Kyoiku Univ.) Matrix means for a fixed rank positive semi-definite matrices

**概要** The set of the positive operators on a Hilbert space forms a Riemannian (or Finsler) manifold where its geodesic is the geometric mean in the sense of Pusz–Woronowicz–Kubo–Ando. The Kubo–Ando operator mean for the non-invertible case is defined as the limit for the invertible case. But it is no longer within the geometric consideration. On the other hand, based on the geodesic of the Grassmannian, Bonnabel–Sepulchre (and Batzies–Huper–Machado–Leitesection) introduces a path of geometric means for positive semi-definite matrices with a fixed rank that coincides with the Kubo–Ando one for the invertible case (We call it BS-mean shortly). We extend any (differentiable) path of Kubo–Ando means to that of the corresponding BS means.



- 17 内山 充 Operator functions and operator means ..... 15  
 (島根大\*・立命館大総合科学技術研究機構)  
 Mitsuru Uchiyama Operator functions and operator means  
 (Shimane Univ.\* / Ritsumeikan Univ.)

概要 It is well-known that  $f(t)$  defined on a right half line is operator monotone if and only if  $f(t)$  is operator concave and  $f(\infty) > -\infty$ . This characterization does not hold if the domain of  $f(t)$  is a finite interval; for instance, an operator monotone function  $\tan t$  on  $(0, \pi/2)$  is not even numerically concave. We will show that  $f(t) > 0$  is operator monotone if and only if  $f(A!B) \leq f(A)!f(B)$  ( $\forall A, B$ ), where  $!$  represents the operator harmonic mean.

- 18 山崎 丈明 (東洋大理工) A generalization of the Aluthge transformation in the viewpoint of operator means ..... 15  
 Takeaki Yamazaki (Toyo Univ.) A generalization of the Aluthge transformation in the viewpoint of operator means

概要 The Aluthge transformation is a very famous operator transformation. It can be considered as a geometric mean of left and right multiplication operators in the finite dimensional case. In this talk, we will extend the Aluthge transformation by considering other operator means via double operator integrals.

#### 14:15~15:10

- 19 C. Bruce (Univ. of Victoria) Partition functions as  $C^*$ -dynamical invariants and actions of congruence monoids ..... 15  
 M. Laca (Univ. of Victoria)  
 武石 拓也 (京都工繊大基盤)  
 Chris Bruce (Univ. of Victoria) Partition functions as  $C^*$ -dynamical invariants and actions of congruence monoids  
 Marcelo Laca (Univ. of Victoria)  
 Takuya Takeishi (Kyoto Inst. Tech.)

概要 We study the phase transition of KMS states for the  $C^*$ -algebras of  $ax+b$ -semigroups of algebraic integers in which the multiplicative part is restricted to a congruence monoid, as in recent work of Bruce generalizing earlier work of Cuntz, Deninger, and Laca. Here we realize the extremal low-temperature KMS states as generalized Gibbs states by constructing concrete representations induced from extremal traces of certain group  $C^*$ -algebras. We use these representations to compute the Murray-von Neumann type of extremal KMS states and we determine explicit partition functions for the type I factor states. The collection of partition functions that arise this way is an invariant under  $R$ -equivariant isomorphism of  $C^*$ -dynamical systems, which produces further invariants through the analysis of the topological structure of the KMS state space. As an application we characterize several features of the underlying number field and congruence monoid in terms of these invariants.

- 20 梶原 毅 (岡山大環境) 高次元の分岐点集合を持つ自己相似写像に付随する  $C^*$ -環の次元群 ..... 15  
 綿谷 安男 (九大\*)  
 Tsuyoshi Kajiwara (Okayama Univ.) Dimension group of the  $C^*$ -algebras associated with self-similar maps  
 Yasuo Watatani (Kyushu Univ.\*) with higher dimensional branched points set

概要 In this talk, we present a method to represent the dimension group of the core of the  $C^*$ -algebra associated with self-similar maps with higher dimensional branched points set, and to get information of dimension groups of them.

- 21 濱田 裕 康 (佐世保工高専)  $C^*$ -algebras generated by multiplication operators and composition operators by functions with self-similar branches ..... 15  
 Hiroyasu Hamada  $C^*$ -algebras generated by multiplication operators and composition operators by functions with self-similar branches  
 (Sasebo Nat. Coll. of Tech.)

**概要** Let  $K$  be a compact metric space and let  $\varphi : K \rightarrow K$  be continuous. We study  $C^*$ -algebra  $\mathcal{MC}_\varphi$  generated by all multiplication operators by continuous functions on  $K$  and a composition operator  $C_\varphi$  induced by  $\varphi$  on a certain  $L^2$  space. Let  $\gamma = (\gamma_1, \dots, \gamma_n)$  be a system of proper contractions on  $K$ . Suppose that  $\gamma_1, \dots, \gamma_n$  are inverse branches of  $\varphi$  and  $K$  is self-similar. We consider the Hutchinson measure  $\mu^H$  of  $\gamma$  and the  $L^2$  space  $L^2(K, \mu^H)$ . Then we show that the  $C^*$ -algebra  $\mathcal{MC}_\varphi$  is isomorphic to the  $C^*$ -algebra  $\mathcal{O}_\gamma(K)$  associated with  $\gamma$  under some conditions.

### 15:20~16:20 特別講演

- 渡 邊 恵 一 (新潟 大理) ムビウスジャイロベクトル空間とその間の連続写像のあるクラスについて  
 Keiichi Watanabe (Niigata Univ.) On Möbius gyrovector spaces and a class of continuous mappings between them

**概要** We study some aspects of Möbius gyrovector spaces from viewpoints of basic theory of functional analysis. First of all, I introduce the notion of gyrocommutative gyrogroups, gyrovector spaces and Möbius gyrovector spaces due to A. A. Ungar. Then I present some fundamental results such as the structure of finitely generated gyrovector subspaces, orthogonal gyrodecomposition and gyroexpansion, Cauchy–Bunyakovsky–Schwarz type inequalities, continuous quasi gyrolinear functionals and a class of continuous mappings between Möbius gyrovector spaces corresponding Hilbert space operators.

## 統計数学

3月16日(月) 第VI会場

9:20~11:55

- 1 星野 浄生 (阪府大理) On a Riemann approximation of the stochastic integral ..... 15  
 Kiyoyuki Hoshino (Osaka Pref. Univ.) On a Riemann approximation of the stochastic integral

概要 We introduce a Riemann-type sum regarded as generalizations of the sums which define the Nualart–Pardoux–Stratonovich integral and the Ogawa integral with respect to the Haar system, and discuss the approximation of some stochastic integrals by this sum. We attempt to characterize some stochastic integrabilities by conditions on the Riemann sum, by which we also introduce a variation of the stochastic  $k$ -integral and give an application example of this related to the volatility estimation.

- 2 小川 重義 (立命館大理工) A noncausal counterpart of Girsanov’s theorem ..... 10  
 Shigeyoshi Ogawa (Ritsumeikan Univ.) A noncausal counterpart of Girsanov’s theorem

概要 We are concerned with an application of the theory of BPE (Brownian particle equation). More precisely, in the framework of the noncausal stochastic calculus, we intend to show a very simple derivation of noncausal counterpart of Girsanov’s theorem on the drift shift of Brownian motion.

- 3 植田 優基 (北大理) Free max-probability theory ..... 15  
 Yuki Ueda (Hokkaido Univ.) Free max-probability theory

概要 In 2006, Ben Arous and Voiculescu introduced free max-convolution and free extreme values. They are analogue of classical max-convolution and classical extreme values, respectively. We expect that there is a relation between free max-probability and asymptotically eigenvalue distributions of spectral maximum of random matrices. In additive case, Bercovici and Pata in 1999 proved the equivalence of limit theorems for classical and freely (additive) infinitely divisible distributions. Based on this result, we studied limit theorems for classical and freely max-infinitely divisible distributions. In this talk, we explain a relation between classical-max and free-max.

- 4 大坂 翔人 (横浜国大理工) 高木クラスの関数における収束の速さについて ..... 15  
 竹居 正登 (横浜国大工)  
 Shoto Osaka (Yokohama Nat. Univ.) On the rate of convergence for Takagi class functions  
 Masato Takei (Yokohama Nat. Univ.)

概要 We consider a generalized version of the Takagi function, which is one of the most famous example of nowhere differentiable continuous functions. We investigate a set of conditions to describe the rate of convergence of Takagi class functions from the probabilistic point of view: The law of large numbers, the central limit theorem, and the law of the iterated logarithm. On the other hand, we show that the Takagi function itself does not satisfy the law of large numbers in the usual sense.

- 5 中島 由人 (京大人間環境) Fractal  $n$ -gons の連結性/パラメータ集合とその部分集合の連結性 ..... 15  
 Yuto Nakajima (Kyoto Univ.) Connectedness of connectedness locus for fractal  $n$ -gons and the remarkable subset

概要 We consider the closure of the set of zeros of polynomials with complex coefficients randomly chosen from a “good” compact subset of complex plane. We prove that the closure of the set of zeros is connected. Furthermore, we apply this result to the study of connectedness locus ( $\mathcal{M}_n$ ) of fractal  $n$ -gons and the remarkable subset  $\mathcal{M}_n^0$ . Fractal  $n$ -gons and  $\mathcal{M}_n$  are introduced by C. Bandt and N. V. Hung (2008). It is already known that  $\mathcal{M}_2$ ,  $\mathcal{M}_2^0$  and  $\mathcal{M}_3$  are connected (T. Bousch (1988), Y. Himeki under supervision Y. Ishii (2018)). We prove that for each  $n = 2, 3, 4, 5, 6$ ,  $\mathcal{M}_n$  is connected and for all  $n$   $\mathcal{M}_n^0$  is connected.

- 6 伊藤 悠 (京都産大理) Integration with respect to Hölder rough paths of order greater than  $1/4$ : an approach via fractional calculus ..... 15  
 Yu Ito (Kyoto Sangyo Univ.) Integration with respect to Hölder rough paths of order greater than  $1/4$ : an approach via fractional calculus

概要 On the basis of fractional calculus, we introduce an integral of controlled paths with respect to Hölder rough paths of order  $\beta \in (1/4, 1/3]$ . Our definition of the integral is given explicitly in terms of Lebesgue integrals for fractional derivatives, without using any arguments from discrete approximation. We demonstrate that for suitable classes of  $\beta$ -Hölder rough paths and controlled paths, our definition of the integral is consistent with the usual definition given by the limit of the compensated Riemann–Stieltjes sum. This result also provides an approach to the integral of 1-forms against geometric  $\beta$ -Hölder rough paths.

- 7 河本陽介 (福岡歯大) Transitions of generalised Bessel kernels related to biorthogonal ensembles ..... 15  
 Yosuke Kawamoto (Fukuoka Dental Coll.) Transitions of generalised Bessel kernels related to biorthogonal ensembles

概要 Biorthogonal Laguerre ensembles are generalizations of classical Laguerre ensembles. Local fluctuation of these ensembles at the origin has been studied, and determinantal kernels in the limit are described in terms of the Wright generalised Bessel functions. The limit kernels are one parameter deformations of the Bessel kernel. I am going to talk about transitions from the generalised Bessel kernels to the sine kernel under appropriate scaling limits in common with the classical Bessel kernels.

- 8 長田翔太 (九大数理) Isomorphism between determinantal point processes and Poisson point processes ..... 15  
 Shota Osada (Kyushu Univ.) Isomorphism between determinantal point processes and Poisson point processes

概要 The determinant point process is a point process for which the determinants of its kernel function give its correlation functions. It describes a repulsive particle system, and some of them have remarkable properties such as rigidity and small variance property. On the other hand, Poisson point processes do not have these properties because the particles are regionally independent. As above, these two classes of point processes have different properties in correlations among particles. However, we prove that they are isomorphic to each other in the sense of ergodic theory. Translation invariant determinant point processes on integer lattices are isomorphic to Bernoulli shifts. This Bernoulli property is a sufficient condition for tail triviality in the measure-preserving dynamical systems. This result, the Poisson property, is the continuous version of the Bernoulli property.

- 9 新井裕太 (千葉大融合理工) The KPZ fixed point for discrete time TASEP ..... 15  
 Yuta Arai (Chiba Univ.) The KPZ fixed point for discrete time TASEP

概要 The totally asymmetric simple exclusion process (TASEP) is one of the simplest interacting stochastic particle system and can be interpreted as a stochastic growth model of an interface, which turns out to belong to the Kardar–Parisi–Zhang (KPZ) universality class. In this talk we consider a discrete time version of the TASEP with arbitrary right finite initial condition. We obtain a single Fredholm determinant representation for the joint distribution function of particle positions. Using this, we show that in the KPZ 1:2:3 scaling limit, the distribution function converges to the one describing the KPZ fixed point introduced by Matetski, Quastel and Remenik (2018).

**14:15~15:00**

- 10 松浦浩平 (京大理) Hölder continuity of Neumann heat kernels on a class of planar domains ..... 15  
 Kouhei Matsuura (Kyoto Univ.) Hölder continuity of Neumann heat kernels on a class of planar domains

概要 Gyrya and Saloff-Coste studied first-order Sobolev spaces on inner uniform domains with Neumann boundary condition. They proved the volume doubling property and the Poincaré inequality for the associated Dirichlet spaces. As a corollary, Neumann heat kernels on inner uniform domains are extended to continuous functions on appropriate completions of the domains and satisfy the two-sided Gaussian estimates. Moreover, the heat kernels are Hölder continuous on the complete metric spaces. However, in the result, it is difficult to calculate even the lower bounds of the indices of Hölder continuity of the heat kernels. In this talk, we give the lower bounds for Neumann heat kernels on a class of planar domains.

- 11 上村稔大 (関西システム理工) 特異な, または退化する係数をもつ対称安定過程の大域的性質について  
 岡村悠奈 (関西大理工) ..... 15  
 Toshihiro Uemura (Kansai Univ.) Global path properties of symmetric stable processes with singular/degenerate  
 Haruna Okamura (Kansai Univ.) coefficients

概要 We show some global path properties of symmetric stable processes with singular/degenerate coefficients in terms of Dirichlet form theory.

- 12 竹内敦司 (東京女大現代教養) Gradient formula for jump processes on manifolds ..... 15  
 Atsushi Takeuchi Gradient formula for jump processes on manifolds  
 (Tokyo Woman's Christian Univ.)

概要 Let  $M$  be a connected and compact Riemannian manifold of dimension  $m$ , and  $O(M)$  the bundle of orthonormal frames. Consider the  $O(M)$ -valued process  $R = \{R_t; 0 \leq t \leq T\}$  determined by the Marcus-type stochastic differential equation with jumps, associated with the canonical horizontal vector fields on  $O(M)$ . The main goal in this talk is to construct the integration by parts formula on the  $M$ -valued process  $X = \{X_t; 0 \leq t \leq T\}$  defined by  $X_t = \pi(R_t)$ , where  $\pi: O(M) \rightarrow M$  is the canonical projection.

**15:15~16:15 特別講演**

- 田口大 (岡山大異分野基礎研) 確率微分方程式の数値解析  
 Dai Taguchi (Okayama Univ.) Numerical analysis of stochastic differential equations

概要 The theory of stochastic calculus and stochastic differential equations (SDEs) introduced by Kiyosi Itô is used to model a random dynamical phenomena in many fields of applications, for example, mathematical finance, physics and biology. For instance, in the field of mathematical finance it has been actively studied from both sides of theory and practice. In particular, financial derivatives are priced by using expectations of a solution of stochastic differential equations, and it is required to accurately calculate their prices. However, since in general it is difficult to obtain explicit form of a solution of stochastic differential equation, it is necessary to approximate the solution by using “discretization”. In this talk, I will talk about recent developments and future issues of numerical analysis for a solution of stochastic differential equations.

**16:30~17:30 特別講演**

- コリンズブノワ (京大理)<sup>b</sup> On the operator norm of random matrices  
 Benoît Collins (Kyoto Univ.) On the operator norm of random matrices

概要 I will review recent results on the operator norm of some random matrices.

## 3月17日(火) 第VI会場

## 9:10~11:25

- 13 高橋 勇人 (Random Data Lab.) Martin-Löf random sets and consistent theorem of posterior distributions ..... 15

Hayato Takahashi (Random Data Lab.) Martin-Löf random sets and consistent theorem of posterior distributions

概要 By the theorem of Doob (1948), if there is a consistent estimator for all parameters, the posterior distribution is consistent for almost all parameters. By the theorem of Breiman et.al (1964), posterior distributions are consistent if conditional probabilities given disjoint cylinder sets are orthogonal, which is much weaker than the existence of consistent estimator for all parameters such as Wald (1949). Frequentist may concern about for which parameter the posterior distribution will converge (Diaconis and Freedman pp. 4). In this presentation, we show that posterior distributions converge weakly to Martin-Löf random set on parameter space with oracle  $P$  of joint distribution iff posterior distribution is consistent for almost all parameters. We discuss random sets that satisfy consistent theorems of Bayes models and propose a definition of random sets with respect to conditional probabilities.

- 14 市原 直幸 (青学大理工) 境界条件を持つ有限期間のマルコフ決定過程に対する値関数の収束について ..... 15

Naoyuki Ichihara (Aoyama Gakuin Univ.) Convergence of value functions for finite horizon Markov decision processes with boundary conditions

概要 We are concerned with the convergence of value functions for some finite horizon Markov decision processes on a countable state space with boundary condition. We observe three different limiting behaviors of value functions depending on the generalized principal eigenvalue of the associated ergodic problem.

- 15 土谷 正明 (金沢大\*)<sup>b</sup> Markov processes controlled by clocks with variable motion on a Euclidean space ..... 15

Masaaki Tsuchiya (Kanazawa Univ.\*) Markov processes controlled by clocks with variable motion on a Euclidean space

概要 Given a bundle of generating triplets of natural additive processes on a Euclidean space, we consider its cumulant and the pseudo-differential operator with the cumulant as symbol, which is an operator-valued measure on the time axis. By using the arguments on Markov selections, we study the existence of a solution with strong Markov property to the martingale problem for the pseudo-differential operator.

- 16 日比野 雄嗣 (佐賀大理工) 非標準表現を用いた標準表現の構成 ..... 10

Yuji Hibino (Saga Univ.) Construction of the canonical representation from a noncanonical representation

概要 A centered Gaussian process is determined by its covariance. However, the method to construct the canonical representation from the covariance has not been obtained. In this talk, we propose a new method to construct the canonical representation for a Gaussian process by using a noncanonical representation.

- 17 瀨口雄史(京大理) Time-inconsistent consumption-investment problems under general discount functions ..... 15  
 Yushi Hamaguchi (Kyoto Univ.) Time-inconsistent consumption-investment problems under general discount functions

概要 We study a time-inconsistent consumption-investment problem with random endowments in a possibly incomplete market under general discount functions. We provide a necessary condition and a verification theorem for an open-loop equilibrium consumption-investment pair in terms of a coupled forward-backward stochastic differential equation. Moreover, we prove the uniqueness of the open-loop equilibrium pair by showing that the original time-inconsistent problem is equivalent to an associated time-consistent one.

- 18 佐久間紀佳(愛知教育大教育) A Clark–Ocone–Häussmann type formula under change of measure for  
 鈴木良一(慶大理工)  $L^1$ -canonical additive processes and its applications ..... 10  
 Noriyoshi Sakuma (Aichi Univ. of Edu.) A Clark–Ocone–Häussmann type formula under change of measure for  
 Ryoichi Suzuki (Keio Univ.)  $L^1$ -canonical additive processes and its applications

概要 In this talk, we show a Clark–Ocone–Häussmann type formula under change of measure for  $L^1$ -canonical additive processes by using a Malliavin–Skorohod calculus in  $L^0$  and  $L^1$  for additive processes developed by Di Nunno–Vives (2017, Stochastics). This is a generalization of Suzuki (2013, COSA). We also show some applications of the main theorem to finance.

- 19 世良透(京大理) Resolution of sigma-fields for multiparticle finite-state action evolutions  
 矢野孝次(京大理) with infinite past ..... 15  
 伊藤悠(京都産大理)  
 Toru Sera (Kyoto Univ.) Resolution of sigma-fields for multiparticle finite-state action evolutions  
 Kouji Yano (Kyoto Univ.) with infinite past  
 Yu Ito (Kyoto Sangyo Univ.)

概要 For multiparticle finite-state action evolutions, we report that the observation sigma-fields are proven to have a resolution consisting of a reduced driving noise, the remote past noise and a third noise. The general theory of infinite convolutions on finite semigroups are utilized in our proofs.

- 20 井上友喜(愛媛大工) 中立的不動点をもつランダム力学系の不変測度 ..... 15  
 Tomoki Inoue (Ehime Univ.) Invariant measures of random dynamical systems with indifferent fixed points

概要 We consider a family of transformations and study a random dynamical system such that one transformation is randomly selected from the family and then applied on each iteration. Especially, we study random dynamical systems with indifferent fixed points and show the existence of absolutely continuous  $\sigma$ -finite invariant measures under some conditions. Further, we research whether the invariant measures are finite or not under some conditions.

11:30~12:00 統計数学分科会総会

3月18日(水) 第VI会場

9:00~12:00

- 21 後藤 佑一 (早大理工) Estimation of trigonometric moments for circular distribution of MA(p) type by using binary series ..... 10  
 Yuichi Goto (Waseda Univ.) Estimation of trigonometric moments for circular distribution of MA(p) type by using binary series

概要 Directional statistics have received a great deal of interest in recent years, and a variety of distributions on the circle have been proposed. In this paper, we propose circular distributions of a moving average model of order  $p$  type which includes the cardioid distribution, and discuss the problem of estimation of trigonometric moments based on binary series. We give an explicit form of the root  $n$  consistent estimator based on clipped series, which enables us to construct an efficient estimator by the Newton–Raphson iterative method. We also show a robustness of the proposed estimator when the probability density function is contaminated with a noise term.

- 22 木村 晃敏 (早大理工) The asymptotic variance estimators of the correlation estimator between latent processes and their asymptotic properties ..... 15  
 Akitoshi Kimura (Waseda Univ.) The asymptotic variance estimators of the correlation estimator between latent processes and their asymptotic properties

概要 In this talk, we treat a model in which the finite variation part of a two-dimensional semi-martingale is expressed by time-integration of latent processes. We propose a correlation estimator between the latent processes and show its consistency and asymptotic mixed normality. Moreover, we propose two types of estimators for asymptotic variance of the correlation estimator and show their asymptotic properties in a high frequency setting. We focus on the proof of the asymptotic properties. Our model includes doubly stochastic Poisson processes whose intensity processes are correlated Itô processes.

- 23 明石 郁哉 (東大経済) Robust regression on hyper-spheres with unspecified heteroscedastic errors ..... 15  
 H. Dette (Ruhr-Univ. Bochum) Robust regression on hyper-spheres with unspecified heteroscedastic errors  
 Fumiya Akashi (Univ. of Tokyo) Robust regression on hyper-spheres with unspecified heteroscedastic errors  
 Holger Dette (Ruhr-Univ. Bochum) Robust regression on hyper-spheres with unspecified heteroscedastic errors

概要 Statistical treatment for a random vector on a hyper-spheres attracts a lot attention recently, and has various applications such as seismic wave analysis, analysis for orientation of wild fire, etc. In this talk the nonlinear regression model whose predictor is a random vector on a hyper-sphere is considered. It is well known that the classical method in “linear statistic” does not work for spherical random vectors. To construct a robust estimator for the nonlinear regression function, this talk employees L1-regression method and kernel-type objective function. The proposed local-linear estimator has asymptotic normality even if the error process has infinite variance, dependent structure or heteroscedasticity. Some simulation experiments illustrate desired finite sample properties of the proposed method.



- 24 明石郁哉 (東大経済) Inference for heavy-tailed time varying processes by self-weighting . . . . 15  
蛭川潤一 (新潟大理)  
 K. Fokianos (Lancaster Univ.)  
Fumiya Akashi (Univ. of Tokyo) Inference for heavy-tailed time varying processes by self-weighting  
Junichi Hirukawa (Niigata Univ.)  
 Konstantinos Fokianos  
 (Lancaster Univ.)

概要 This talk considers a parameter estimation problem of time-varying autoregressive models under the presence of infinite variance. Although there is rich literature on locally stationary processes, the classical papers always assume the finite variance of the model. This talk constructs a robust estimator based on the self-weighting approach proposed by Ling (2005, Journal of Royal Statistical Society B) and least absolute deviations regression. The proposed estimator is show to be asymptotically normal regardless of whether the error term has infinite variance or not. Finite sample performance of the proposed method is also investigated by simulation experiments.

- 25 劉言 (早大理工) Persistence diagram for Granger causality . . . . . 15  
木村晃敏 (早大理工)  
谷口正信 (早大理工)  
 H. Ombao  
 (King Abdullah Univ. of Sci. Tech.)  
Yan Liu (Waseda Univ.) Persistence diagram for Granger causality  
Akitosho Kimura (Waseda Univ.)  
Masanobu Taniguchi (Waseda Univ.)  
 Hernando Ombao  
 (King Abdullah Univ. of Sci. Tech.)

概要 We propose a topological approach to statistically analyzing the Granger causality. Granger introduced his celebrated new measure of causality in the sense of prediction errors of multivariate time series 50 years ago. We localize his idea and construct a theory based on locally stationary processes for its alternative version, a natural refinement for stationary processes by Hosoya. To construct the theory, we provide a Gaussian approximation of the suprema of empirical spectral processes. Especially, the local extension of the theory serves for the statistical inference for the Granger causality curve. In addition, we provide a bootstrap procedure for the approximation to construct confidence bands. Finally, we discuss the persistence diagrams and persistence landscapes for the causality curves and numerically construct some examples of locally stationary processes for our simulations studies.

- 26 小池健一 (筑波大数理物質) ベイズ情報不等式の等号達成条件 . . . . . 10  
 Ken-ichi Koike (Univ. of Tsukuba) Attainment conditions for Bayesian information inequalities

概要 Necessary and sufficient conditions for the attainments of the Borovkov-Sakhanenko and the van Trees information bounds are given when the underlying distribution is an exponential family and the conjugate prior, and the estimand is a function of the parameter. Moreover, the attainments under the Jeffreys prior are also considered.

- 27 佃 康 司 (東大総合文化) Pitman–Yor 事前過程を用いた場合の事後過程の弱収束について …… 15  
 Koji Tsukuda (Univ. of Tokyo) A note on the weak convergence of the posterior process when the Pitman–Yor process prior is placed

概要 Under a problem setting of the Bayesian nonparametrics, the Pitman–Yor process, which has two parameters denoted by  $\alpha(\in [0, 1))$  and  $\theta(> -\alpha)$ , is a popular prior. In the literature, when the Pitman–Yor process is placed, the conditional weak convergence of the centered posterior process rescaled by the ordinary rate factor  $n^{1/2}$  is provided, where  $n$  is the number of observations. Introducing a zero-mean process rescaled by the different rate factor  $(\theta + n\alpha)^{1/2}$ , we present a corresponding result under the asymptotic regime that  $\theta$  and  $n$  simultaneously increase. This result reveals that the rate of convergence changes when  $\theta/n \rightarrow \infty$ .

- 28 柿 沢 佳 秀 (北 大 経 済) 非負データに対する密度比推定, 条件付き密度推定 …… 15  
 Yoshihide Kakizawa (Hokkaido Univ.) Density ratio/conditional density estimation for nonnegative data

概要 For the data supported on  $[0, \infty)$  or  $[0, 1]$  ( $[0, \infty)^d$  or  $[0, 1]^d$ ), asymmetric kernel density estimation has been well-studied in the recent literature. In this talk, we consider two problems, on the basis of the AK method: one is the estimation of density ratio in the two-sample setting, and the other is the estimation of conditional density for the bivariate dataset.

- 29 種 市 信 裕 (北 教 大 札 幌) 多次元分割表における多要因間独立性検定統計量の分布の漸近展開に基  
 関 谷 祐 里 (北 教 大 釧 路) づく近似 …… 15  
 外 山 淳 (数学利用研究所)

Nobuhiro Taneichi (Hokkaido Univ. of Edu.) Approximations of the distributions of test statistics for independence among groups of factors in a multi-way contingency table based on asymptotic expansion  
 Yuri Sekiya (Hokkaido Univ. of Edu.)  
 Jun Toyama (Inst. for Practical Appl. of Math.)

概要 We consider hypotheses of independence among groups of factors in a multi-way contingency table. Hypotheses include that of one factor independence from the other two in a 3-way contingency table and that of independence among three factors, one factor and one factor in a 5-way contingency table. In this report, we derive approximations of distribution of the test statistics for the hypotheses based on asymptotic expansion.

- 30 生 亀 清 貴 (日 大 経 済) 正方分割表における儉約な 2 変量 t 分布型対称モデル …… 15  
 Kiyotaka Iki (Nihon Univ.) Parsimonious bivariate t-distribution type symmetry models for square contingency tables

概要 For the analysis of square contingency tables with the same row and column ordinal classifications, this presentation proposes new models, which may be appropriate for a square ordinal table if it is reasonable to assume an underlying bivariate t-distribution having any degrees of freedom. The new models have stronger restriction than t-distribution type symmetry models (Iki et al., 2013; Iki et al., 2018). The simulation studies based on bivariate t-distribution are given.

- 31 弓場 弘 (国際学術交流センター) E\*-optimal balanced third-order designs of resolution  $R^*({10,01})$  with  $N < \nu(m)$  for  $3^m$  factorials ..... 15  
兵頭 義史 (岡山理大総合情報研)  
Hiromu Yumiba (Int. Center for Academic Exchange) E\*-optimal balanced third-order designs of resolution  $R^*({10,01})$  with  $N < \nu(m)$  for  $3^m$  factorials  
Yoshifumi Hyodo (Okayama Univ. of Sci.)

**概要** We consider a balanced third-order ( $3^m$ -BTO) design  $T$  of resolution  $R^*({10,01})$  derived from an SA( $m; \{\lambda_{x^m-x-yy}\}$ ) with  $N$  assemblies and  $m \geq 6$ . Let  $\sigma^2 \lambda_{\max}(T)$  be the maximum eigenvalue of the variance-covariance matrix of the estimators concerning with all the main effects based on  $T$ . If  $\lambda_{\max}(T_0) \leq \lambda_{\max}(T)$  for any  $T$ , then  $T_0$  is said to be E\*-optimal, where  $T_0$  is a  $3^m$ -BTO design of resolution  $R^*({10,01})$  derived from an SA with  $N$  assemblies. In this talk, we give E\*-optimal  $3^m$ -BTO designs of resolution  $R^*({10,01})$  derived from SA's for  $6 \leq m \leq 8$ , where  $N < \nu(m)$ . Here  $\nu(m) (= 1 + 2m + \frac{1}{6}m(m-1)(m+7))$  is the number of non-negligible factorial effects.

## 14:15~15:05

- 32 今野 良彦 (日本女大理) Shrinkage estimation of mean for complex multivariate normal distribution with unknown covariance when  $p > n$  ..... 15  
清田 聡美 (日本女大理)  
Yoshihiko Konno (Japan Women's Univ.) Shrinkage estimation of mean for complex multivariate normal distribution with unknown covariance when  $p > n$   
Satomi Seita (Japan Women's Univ.)

**概要** We consider the problem of estimating the mean vector of the multivariate complex normal distribution with unknown covariance matrix under an invariant loss function when  $n$ , the sample size, is smaller than  $p$ , the dimension of the mean vector. The problem of difficulty lies in the fact that the sample covariance matrix becomes singular. Following the approach of Chételet and Wells (2012, Ann. Statist. vol. 40 p. 3137–3160), we show that a modification of Baranchik-like-type estimators beats the MLE if it satisfies certain conditions. Based on this result, we propose the James–Stein-like shrinkage and its positive-part estimators.

- 33 矢田 和善 (筑波大数理物質) 高次元相互共分散行列の特異値推定について ..... 15  
青嶋 誠 (筑波大数理物質)  
Kazuyoshi Yata (Univ. of Tsukuba) Singular value estimation for high-dimensional cross-covariance matrix  
Makoto Aoshima (Univ. of Tsukuba)

**概要** In this talk, we consider singular value estimations for a high-dimensional cross-covariance matrix. We first consider asymptotic properties of the conventional estimator of the singular values. We show that the estimator is affected by high-dimensional noise structures directly, so that it becomes inconsistent. In order to overcome the difficulty, we proposed a new singular value estimation method via the extended cross-data-matrix methodology. We show that the new method can enjoy consistency properties for the singular values in high-dimensional settings.

- 34 石井 晶 (東京理大理工) 高次元固有ベクトルの検定について ..... 15  
 矢田 和善 (筑波大数理物質)  
 青嶋 誠 (筑波大数理物質)  
 Aki Ishii (Tokyo Univ. of Sci.) A test procedure for high-dimensional eigenvectors  
 Kazuyoshi Yata (Univ. of Tsukuba)  
 Makoto Aoshima (Univ. of Tsukuba)

**概要** In this talk, we consider testing high-dimensional eigenvectors. We produce a test statistic by using the extended cross-data-matrix (ECDM) methodology and show the unbiasedness of the ECDM test statistic even in a high-dimensional setting. We also show that the test statistic holds the asymptotic normality. We propose a new test procedure by using the asymptotic normality and evaluate its size and power asymptotically. We also give a real data analysis by using a microarray data set.

#### 15:20~16:20 特別講演

- 榎本 理恵 (成蹊大理工) 高次元成長曲線モデルにおける情報量規準の一致性  
 Rie Enomoto (Seikei Univ.) Consistency of some information criteria in high-dimensional growth curve models

**概要** In multivariate regression model, it is known that the AIC has no consistency and the BIC has consistency under large-sample framework. However, Fujikoshi et al. (2014) and Yanagihara et al. (2015) note that the AIC has consistency and the BIC has no consistency under a high-dimensional framework. The AIC and its modifications have been proposed for selecting the degree in the growth curve model under a large-sample framework and a high-dimensional framework by Satoh et al. (1997) and Fujikoshi et al. (2013), respectively. They note that the AIC and its modifications have no consistency property. The purpose of this paper to discuss high-dimensional asymptotic distributions of the estimators and consistency property of some information criteria under a high-dimensional framework. Our results are checked numerically by conducting a Monte Carlo simulation.

#### 16:35~17:35 特別講演

- 藤森 洸 (早大理工) 高次元・スパースな設定における確率過程の統計モデルに対する Dantzig selector  
 Kou Fujimori (Waseda Univ.) The Dantzig selector for statistical models of stochastic processes in high-dimensional and sparse settings

**概要** The Dantzig selector, which was proposed by Candès and Tao in 2007, is an estimation procedure for regression models in high-dimensional and sparse settings. In this talk, the Dantzig selectors for some statistical models of stochastic processes are discussed. We apply this procedure to Cox's proportional hazards model and some specific models of diffusion processes and prove the consistencies and the variable selection consistencies of the estimators. Based on partial likelihood and quasi-likelihood methods which were studied intensively in low-dimensional settings, we study these statistical models of stochastic processes in high-dimensional and sparse settings, which need some mathematically challenging tasks. The consistencies of the estimators are derived from the stochastic maximal inequalities which are derived from Bernstein's inequalities for martingales and conditions, which are known to be similar to the restricted eigenvalue conditions, on Hessian matrices of log-(partial or quasi)-likelihood functions. We prove that consistencies of the estimators imply the variable selection consistencies which enable us to reduce the dimension. Using the dimension reduction, asymptotically normal estimators can be constructed.

## 応用数学

3月16日(月) 第V会場

10:00~11:40

- 1 小林 雅人 (神奈川大工) Answer Henegphan–Petersen’s question on alternating permutations and Euler numbers ..... 15  
 Masato Kobayashi (Kanagawa Univ.) Answer Henegphan–Petersen’s question on alternating permutations and Euler numbers

概要 The following equality dates back to André in 1879:  $\sum_{n=0}^{\infty} E_n \frac{x^n}{n!} = \sec x + \tan x$  where  $E_n$  is half the number of alternating permutations in  $S_n$  (with  $E_0 = E_1 = 1$ ). Recently, Heneghan–Petersen in 2014 introduced the sequences  $(F_n)_{n \geq 2}$ ,  $(G_n)_{n \geq 2}$  as a refinement of  $E_n$ . In terms of formal power series, they proved  $F_{2n} - G_{2n} = E_{2n-2}$  saying at the end: Is there a bijective explanation for this? In this talk, I will show you an answer to this question with constructing some bijection explicitly.

- 2 稲葉 ゆき江 (日本女大理) Counting rooted spanning forests and Chebyshev polynomials ..... 15  
 藤田 玄 (日本女大理)  
 近藤 剛史 (鹿児島大理)  
 Yukie Inaba (Japan Women’s Univ.) Counting rooted spanning forests and Chebyshev polynomials  
 Hajime Fujita (Japan Women’s Univ.)  
 Takefumi Kondo (Kagoshima Univ.)

概要 In this talk we consider a counting of rooted spanning forests of weighted graphs with fixed set of nodes. This counting problem has its origin in Kenyon–Wilson’s formula, which gives a relation between counting of certain rooted spanning forests and minors of response matrix. We consider a family of weighted graphs which we call the  $n$ -gon graph. Our main result is an explicit formula of the counting for  $n$ -gon graph in terms of Chebyshev polynomials of the first kind.

- 3 佐藤 巖 (小山工高専) The partial differential coefficients for the second Bartholdi zeta function  
 松谷 茂樹 (金沢大理工) of a graph ..... 15  
 三橋 秀生 (法政大理工)  
 森田 英章 (室蘭工大工)  
 Iwao Sato (Oyama Nat. Coll. of Tech.) The partial differential coefficients for the second Bartholdi zeta function  
 Shigeki Matsutani (Kanazawa Univ.) of a graph  
 Hideo Mitsunashi (Hosei Univ.)  
 Hideaki Morita  
 (Muroran Inst. of Tech.)

概要 We present weighted versions for the results of Li and Hou’s on the partial derivatives of the determinant part in the determinant expression of the Bartholdi zeta function of a graph  $G$ . Furthermore, we give a formula for the weighted Kirchhoff index function of a regular covering of  $G$  in terms of that of  $G$ .

- 4 石川 彩香 (横浜国大理工) 有限有向グラフに対する四元数水野–佐藤ゼータ函数の伊原表示 ..... 15  
 Ayaka Ishikawa (Yokohama Nat. Univ.) The Ihara expressions of the quaternionic Mizuno–Sato zeta functions  
 for digraphs

概要 The Mizuno–Sato zeta functions are generalized Ihara zeta functions, and the weights are complex numbers. The definitions of the quaternionic Mizuno–Sato zeta functions for finite symmetric digraphs were given by Konno, Mitsunashi, and Sato. In this study, we re-define them for finite digraphs with loops and multi arcs and give the determinant expressions called Ihara expressions.

- 5 加田 修 (法政大理工) Characteristic polynomials and zeta functions of equitably partitioned graphs ..... 15  
Osamu Kada (Hosei Univ.) Characteristic polynomials and zeta functions of equitably partitioned graphs

概要 It is well known that when a graph is equitable, the characteristic polynomial of the quotient graph divides that of the original graph, but the remainder part is not well investigated. We define a deletion graph, and give a similarity transformation exchanging the adjacency matrix which is compatible with the equitable partition for a block triangular matrix whose diagonal blocks are the adjacency matrix of the quotient graph and the deletion graph, which gives the remainder part. We apply the result to the Ihara–Bartholdi zeta function.

- 6 森田 英章 (室蘭工大工) 行列式表示をもたないグラフゼータについて ..... 15  
Hideaki Morita (Muroran Inst. of Tech.) On the determinant expression for graph zeta functions

概要 Graph zeta functions naturally have the three expressions—the exponential expression, the Euler expression, and the determinant expression of Hashimoto type (the Hashimoto expression). In this talk, a new family of graph zetas is introduced, which does not have the Hashimoto expression.

#### 14:20~16:05

- 7 藤田 慎也 辺着色グラフの rainbow connectivity に関する最近の話題 ..... 10  
(横浜市大データサイエンス)  
Shinya Fujita (Yokohama City Univ.) Recent topics on rainbow connectivity in edge-colored graphs

概要 Some recent results on rainbow connectivity in edge-colored graphs will be reviewed.

- 8 古谷 倫貴 (北里大一般教育) Bounds on self domination number and an edge-deletion operation in trees ..... 15  
Michitaka Furuya (Kitasato Univ.) Bounds on self domination number and an edge-deletion operation in trees

概要 Let  $G$  be a graph, and let  $c \in \mathbb{R}^+ \cup \{\infty\}$ . A function  $f : V(G) \rightarrow \{0, 1, c\}$  is called a  $c$ -self dominating function of  $G$  if for  $x \in V(G)$  with  $f(x) < c$ ,  $\max\{f(y) : y \in N_G(x)\} \geq 1$ . The minimum weight  $w(f) := \sum_{x \in V(G)} f(x)$  of a  $c$ -self dominating function  $f$  of  $G$  is called the  $c$ -self domination number of  $G$ . In this talk, we give a sharp upper bound of a  $c$ -self domination number for all  $c \in \mathbb{R}^+ \cup \{\infty\}$ . To prove our main result, we focus on an edge-deletion operation for trees, and for  $\mathcal{P} \subseteq \{P_n : n \in \mathbb{N}\}$ , we characterize the trees  $T$  such that one of components of  $T - e$  belongs to  $\mathcal{P}$  for every  $e \in E(T)$ .

- 9 安藤 清 Contractible edges and contractible triangles in a 3-connected graph .. 15  
(国立情報学研・JST ERATO)  
江川 嘉美 (東京理大理)  
Kiyoshi Ando Contractible edges and contractible triangles in a 3-connected graph  
(Nat. Inst. of Information/JST ERATO)  
Yoshimi Egawa (Tokyo Univ. of Sci.)

概要 Let  $G$  be a 3-connected graph. An edge (a triangle) of  $G$  is said to be a 3-contractible edge (a 3-contractible triangle) if the contraction of it results in a 3-connected graph. We denote  $E_c(G)$  and  $\mathcal{T}_c(G)$  the set of 3-contractible edges of  $G$  and the set of 3-contractible triangles of  $G$ , respectively. We prove that if  $|V(G)| \geq 7$ , then  $|E_c(G)| + a|\mathcal{T}_c(G)| \geq \min\{\frac{1}{2} + \frac{a}{3}, \frac{6}{7}\}|V(G)|$  for every nonnegative real number  $a$ . As a corollary of this, if  $|V(G)| \geq 7$ , then  $|E_c(G)| + \frac{15}{14}|\mathcal{T}_c(G)| \geq \frac{6}{7}|V(G)|$ ; and if  $G$  has no contractible triangles, then  $|E_c(G)| \geq \frac{6}{7}|V(G)|$ . We also determine the extremal graphs of these results.

- 10 奈良知恵 (明大 MIMS) 高次元正軸多胞体の正三角形面からなる 2-スケルトンの連続平坦化 . . . . . 15  
 伊藤仁一 (嵯山女学園大教育)  
 Chie Nara (Meiji Univ.) Continuous flattening of the 2-skeletons of triangular faces in higher  
 Jin-ichi Itoh (Sugiyama Jogakuen Univ.) dimensional cross-polytopes

概要 We can continuously flatten the surface of a regular octahedron onto any of its faces without stretching and cutting. This is accomplished by moving creases to change the shapes of some faces successively, following Sabitov's volume preserving theorem. We extend this result to higher dimensional regular cross-polytopes by considering the 2-dimensional skeleton of a polytope, corresponding to the surface of a three dimensional polyhedron.

- 11 沼田泰英 (信州大理) ループ付き完全グラフから決まる行列の固有値について . . . . . 15  
 矢澤明喜子 (信州大総合医理工)  
 Yasuhide Numata (Shinshu Univ.) The eigenvalues of a matrix defined by the complete graph with selfloops  
 Akiko Yazawa (Shinshu Univ.)

概要 We consider a matrix whose indices are the edge of the complete graph with selfloops. Assume that entries are the same if the indices are isomorphic as graphs. We compute the eigenvalues of this matrix. We also apply the main theorem to the eigenvalues of the second Hessian matrix with respect to divided power operators of the elementary symmetric polynomial  $e_l(x_1^k, \dots, x_n^k)$ .

- 12 須田庄 (防衛大) 有向グラフの固有値の重複度 . . . . . 10  
 A. Gavrilyuk (Pusan Nat. Univ.)  
 Sho Suda (Nat. Defense Acad. of Japan) On the multiplicities of digraph eigenvalues  
 Alexander Gavrilyuk  
 (Pusan Nat. Univ.)

概要 We show an upper bound for the order of a digraph (or a mixed graph) whose Hermitian adjacency matrix has an eigenspace of prescribed codimension. This generalizes the so-called absolute bound for (simple) graphs first shown by Delsarte, Goethals, and Seidel (1977) and extended by Bell and Rowlinson (2003).

- 13 深作亮也 (九大数理) グラフのテンソル積の染色数とグレブナー基底 . . . . . 15  
 古谷倫貴 (北里大一般教育)  
 東谷章弘 (阪大情報)  
 Ryoya Fukasaku (Kyushu Univ.) Chromatic numbers of tensor products of graphs and Gröbner basis  
 Michitaka Furuya (Kitasato Univ.)  
 Akihiro Higashitani (Osaka Univ.)

概要 For a graph  $G$ , let  $\chi(G)$  denote the chromatic number. In graph theory, the following famous conjecture posed by Hedetniemi has been studied: For two graphs  $G$  and  $H$ ,  $\chi(G \times H) = \min\{\chi(G), \chi(H)\}$ , where  $G \times H$  is the tensor product of  $G$  and  $H$ . In this talk, we give a reduction of Hedetniemi's conjecture to an inclusion relation problem on ideals of polynomial rings, and we demonstrate computational experiments for partial solutions of Hedetniemi's conjecture along such a strategy using Gröbner basis.

**16:20~17:20 特別講演**

- 澤 正 憲 (神戸大システム情報) 古典直交多項式, 矩形求積公式, 幾何的デザインの有理性について  
 Masanori Sawa (Kobe Univ.) On the rationality of classical orthogonal polynomials, quadrature formulas and geometric designs

**概要** In this talk we first introduce a certain system of Diophantine equations, originally designed by Hausdorff (1909, Math. Ann.) as a simplification of Hilbert's solutions of Waring's problem, and then review a close relationship among classical orthogonal polynomials in special function theory, quadrature formulas in numerical analysis, and geometric designs in combinatorics, with particular emphasis on their 'rationality'. We also look at some recent results including, non-solvability results for our equations, an alternative proof of a theorem by Delsarte, Goethals and Seidel on tight spherical 6-designs (1977). This talk may moreover touch with a connection with optimal design problem in statistics. Some of the results given in this talk have been found as a joint work with Yukihiro Uchida.

3月17日(火) 第V会場

**9:50~11:15**

- 14 花岡 遼大 (横浜国大理工) 1次元2状態量子ウォークを用いた時系列解析 ..... 10  
 今野 紀雄 (横浜国大工)  
 小山 翔平 (横浜国大理工)  
Ryota Hanaoka (Yokohama Nat. Univ.) A time-series analysis based on two-state quantum walk in one dimension  
Norio Konno (Yokohama Nat. Univ.) sion  
Shohei Koyama (Yokohama Nat. Univ.)

**概要** Konno introduced a time-series model via the quantum walk in 2019. We show some properties of the model based on two-state quantum walk in one dimension. Moreover we present results on an application of BTC/USD price data.

- 15 浅野 雅裕 (横浜国大理工) 2次元正方格子上の Grover walk の時間平均極限測度について ..... 15  
 今野 紀雄 (横浜国大工)  
 成松 明廣 (横浜国大理工)  
Masahiro Asano (Yokohama Nat. Univ.) Long-time behavior of the Grover walk on the two-dimensional lattice  
Norio Konno (Yokohama Nat. Univ.)  
Akihiro Narimatsu  
 (Yokohama Nat. Univ.)

**概要** The Grover walk has been investigated for many applications, such as quantum searching algorithms. Inui et al. (2005) got both limit measure and weak limit theorem as a long-time behavior of the 3-state Grover walk on the one-dimensional lattice. However, the 4-state Grover walk on the two-dimensional lattice was studied only for time-averaged limit measure at the starting point and weak limit theorem. We obtained the time-averaged limit measure on coordinate axes and analyzed the asymptotic behavior for the Grover walk in two dimensions.



- 16 内藤 拓人 (横浜国大理工) ウォークによるレコメンデーションモデル (Part 1) ..... 10  
 黄海 仲星 (横浜国大理工)  
 今野 紀雄 (横浜国大工)  
 高橋 佐良人 (横浜国大理工)  
 Takuto Naito (Yokohama Nat. Univ.) Recommendation models based on walks (Part 1)  
 Chusei Kiumi (Yokohama Nat. Univ.)  
 Norio Konno (Yokohama Nat. Univ.)  
 Sarato Takahashi  
 (Yokohama Nat. Univ.)

概要 These days, we use online sites to see news, SNS, and so on. There are various services using recommendation models. Recently, the quality of recommendation models influences on the earnings of merchandises very much. In this situation, we propose new models based on user's selection process as walks. Moreover, we analyze user's preference using random, correlated, and quantum walks, respectively.

- 17 黄海 仲星 (横浜国大理工) ウォークによるレコメンデーションモデル (Part 2) ..... 10  
 今野 紀雄 (横浜国大工)  
 内藤 拓人 (横浜国大理工)  
 高橋 佐良人 (横浜国大理工)  
 Chusei Kiumi (Yokohama Nat. Univ.) Recommendation models based on walks (Part 2)  
 Norio Konno (Yokohama Nat. Univ.)  
 Takuto Naito (Yokohama Nat. Univ.)  
 Sarato Takahashi  
 (Yokohama Nat. Univ.)

概要 The recommendation models are used in many online sites. We consider recommendation models proposed in Part 1 based on random, correlated, and quantum walks, respectively. Then we show the behaviors of these models by numerical simulations. The quantum-walk version model has a peculiar feature compared with random- and correlated-walk version models.

- 18 小松 堯 (東大工) パスの数え上げによる 1 次元 2 状態量子ウォークに対する散乱行列の明  
 今野 紀雄 (横浜国大工) 示公式 ..... 15  
 森岡 悠 (愛媛大理工)  
 瀬川 悦生 (横浜国大環境情報)  
 Takashi Komatsu (Univ. of Tokyo) An explicit expression of scattering matrix of a two state quantum walk  
 Norio Konno (Yokohama Nat. Univ.) on one-dimensional lattice by path counting  
 Hisashi Morioka (Ehime Univ.)  
 Etsuo Segawa (Yokohama Nat. Univ.)

概要 We obtain an explicit expression of the scattering matrix of a free quantum walk on one-dimensional lattice with finite impurities by using a path counting method. We show that this expression is essentially obtained by counting the escaping amplitude from the impurity's region; in other word, an analysis on the principal submatrix of this unitary time evolution operator with respect to the impurities. As a byproduct, we clarify that the reflection and transmission rates of the quantum walk model introduced by Matsue et al (2018), which is connected to the stationary Schrödinger equation on the double well potentials, can be expressed by this scattering matrix.

- 19 井手 勇介 (金沢工大) 量子ウォークの極限分布に付随する直交多項式と対応するランダムウォークについて ..... 15  
 今野 紀雄 (横浜国大工) 今野 紀雄 (横浜国大工) 今野 紀雄 (横浜国大工) 今野 紀雄 (横浜国大工) 今野 紀雄 (横浜国大工)  
 Yusuke Ide (Kanazawa Inst. of Tech.) Relationships between orthonormal polynomial related to the limit distribution of quantum walk and corresponding random walk  
 Norio Konno (Yokohama Nat. Univ.) Relationships between orthonormal polynomial related to the limit distribution of quantum walk and corresponding random walk

概要 We consider relationships between orthonormal polynomial related to the limit distribution of discrete-time quantum walk on the line and corresponding random walk on the half line. In order to connect the orthonormal polynomial and the random walk, we use so-called Jacobi matrix for each objects. In this talk, we discuss about roles of the limit distribution in corresponding random walk and/or other induced processes.

11:30~11:50 2019年度日本数学会応用数学研究奨励賞授賞式

3月18日(水) 第V会場

9:30~12:00

- 20 中島 健 (静岡大理) On approximation of 2D persistence modules by interval-decomposables  
 浅芝 秀人 (静岡大理) ..... 15  
 E. G. Escolar  
 (理化学研AIP・京大高等研)  
 吉脇 理雄  
 (理化学研AIP・京大高等研・阪市大数学研)  
 Ken Nakashima (Shizuoka Univ.) On approximation of 2D persistence modules by interval-decomposables  
 Hideto Asashiba (Shizuoka Univ.)  
 Emerson Gaw Escolar  
 (RIKEN/Kyoto Univ.)  
 Michio Yoshiwaki  
 (RIKEN/Kyoto Univ./Osaka City Univ.)

概要 In this work, we propose a new invariant for 2D persistence modules called the compressed multiplicity and show that it generalizes the notions of the dimension vector and the rank invariant. In addition, we propose an “interval-decomposable approximation”  $\delta^*(M)$  of a 2D persistence module  $M$ . In the case that  $M$  is interval-decomposable, we show that  $\delta^*(M) = M$ . Furthermore, even for representations  $M$  not necessarily interval-decomposable,  $\delta^*(M)$  preserves the dimension vector and the rank invariant of  $M$ .

- 21 E. G. Escolar (理化学研 AIP・京大高等研) Every pair of  $\Lambda$ -interleavings is  $\tilde{\Lambda}$ -interleaved ..... 15  
 K. F. Meehan (京大高等研)  
 吉 脇 理 雄  
 (理化学研AIP・阪市大数学研・京大高等研)

Emerson Gaw Escolar (RIKEN/Kyoto Univ.) Every pair of  $\Lambda$ -interleavings is  $\tilde{\Lambda}$ -interleaved  
 Killian F. Meehan (Kyoto Univ.)  
 Michio Yoshiwaki (RIKEN/Osaka City Univ./Kyoto Univ.)

概要 There is an isometry theorem relating the interleaving distance between 1D persistence modules and the bottleneck distance of their corresponding barcodes. The bottleneck distance is defined by matchings between the barcodes, and can be seen as a “diagonal” interleaving of the persistence modules. We wish to study how far arbitrary interleavings are from diagonal interleavings. To that end, we work in a more general setting of prosets. We introduce the concept of a shoelace of a proset and show that the representation category of the shoelace is isomorphic to the category of interleavings. Through this, we can formulate interleavings between interleavings in this new setting. Finally, we show that any two  $\Lambda$ -interleavings are  $\tilde{\Lambda}$ -interleaved, where  $\tilde{\Lambda}$  is a “twisted” interleaving on the shoelace naturally induced from  $\Lambda$ .

- 22 大 林 一 平 (理化学研 AIP・東北大 AIMR) Field choice problem on persistent homology ..... 15  
 吉 脇 理 雄  
 (理化学研AIP・阪市大数学研・京大高等研)  
 Ippei Obayashi (RIKEN/Tohoku Univ.) Field choice problem on persistent homology  
 Michio Yoshiwaki (RIKEN/Osaka City Univ./Kyoto Univ.)

概要 In this presentation, I will talk about the problem of the choice of a coefficient field on persistent homology. When we compute a persistence diagram, we need to select a coefficient field before computation. We should understand the dependency of the diagram on the coefficient field for the better computation and interpretation of the diagram. We give some sufficient and necessary conditions for the independence of the diagram to the coefficient field. We also give an efficient algorithm to determine whether a given input satisfies the condition or not. The algorithm is already implemented in HomCloud, our data analysis software based on persistent homology. The software will be helpful for data analysis using persistent homology. This is the joint work with Michio Yoshiwaki (RIKEN).

- 23 見 上 達 哉 (京 大 理) 結晶格子における最速浸透問題 ..... 15  
 Tatsuya Mikami (Kyoto Univ.) First passage percolation on a crystal lattice

概要 The time evolution version of the percolation model is called the first passage percolation model: each edge of the cubical lattice is assigned a random time, and consider the growth of the percolation region  $B(t)$ , which consists of the vertices that can be arrived from the origin within time  $t > 0$ . One of the most basic theorem is called shape theorem, saying that the normalized region  $B(t)/t$  converges to some limit shape. In this study, a generalized FPP model, defined on a crystal lattice, is considered. In this setting, the generalized version of the shape theorem is obtained and the limit shape depends only on the graph structure and the period of the lattice.

- 24 清水超貴 (京大理) マルチパラメータパーシステントホモロジーの分解論における極限定理  
平岡裕章 (京大高等研) ..... 15  
Tatsuki Shimizu (Kyoto Univ.) Limit theorems in the decomposition theory of multi-parameter persis-  
Yasuaki Hiraoka (Kyoto Univ.) tent homology

**概要** Unlike the 1D case, there exists an algebraic difficulty in the decomposition theory of multi-parameter persistence modules. We try to solve this difficulty not purely algebraically, but stochastically by considering the appropriate scaling limit. We first show the existence of the scaling limit of some invariants: multiplicity of the interval representation and rank invariant.

- 25 E. G. Escolar Mapping firms' locations in technological space: A topological analysis  
 (理化学研AIP・京大高等研) of patent statistics ..... 15  
平岡裕章 (京大高等研)  
伊神満 (Yale Univ.)  
Y. Ozcan (MIT Sloan)  
Emerson Gaw Escolar Mapping firms' locations in technological space: A topological analysis  
 (RIKEN/Kyoto Univ.) of patent statistics  
Yasuaki Hiraoka (Kyoto Univ.)  
Mitsuru Igami (Yale Univ.)  
Yasin Ozcan (MIT Sloan)

**概要** Where do firms innovate? Locating and visualizing them in technological space is challenging, because it is high-dimensional and unstructured. We address this issue by using a method in topological data analysis called Mapper, which combines local clustering with global reconstruction. We apply this method to a panel of 333 major firms' patent portfolios in 1976–2005 in 430 technological areas and propose a definition of the characteristic “flares” that appear in the Mapper graph. Results suggest the Mapper graph captures salient patterns in firms' patenting histories, and the type and length of flares are correlated with firms' financial performances in a statistically and economically significant manner.

- 26 井元佑介 (京大ASHBi) 力学系モデルおよび統計的因果探索に基づく遺伝子制御ネットワークの  
平岡裕章 推定 ..... 15  
 (京大高等研・京大ASHBi・理化学研AIP)  
清水昌平  
 (滋賀大DS・理化学研AIP)  
前田高志ニコラス (理化学研AIP)  
小島洋児 (京大iPS細胞研)  
斎藤通紀  
 (京大ASHBi・京大医・京大iPS細胞研)  
Yusuke Imoto (Kyoto Univ.) Estimate of gene regulatory network based on dynamical system and  
Yasuaki Hiraoka statistical causal discovery  
 (Kyoto Univ./Kyoto Univ./RIKEN)  
Shohei Shimizu (Shiga Univ./RIKEN)  
Takashi Nicholas Maeda (RIKEN)  
Yoji Kojima (Kyoto Univ.)  
Mitinori Saitou  
 (Kyoto Univ./Kyoto Univ./Kyoto Univ.)

**概要** The gene regulatory network (GRN) is a directed graph model with annotated edges that shows the regulatory relationship of gene expressions. In this study, we develop a method that estimates GRN from gene expression data by utilizing a causal discovery method ‘LiNGAM’ and a dynamical system ‘switching model’. We derive a linear model that is applicable in LiNGAM by discretizing the switching model and applying the multi-regression to it. As a result, GRN is constructed by superposing the adjacency matrices estimated by LiNGAM.

## 14:30~15:50

- 27 後藤田 剛 (名大多元数理) Numerical study of initial configurations leading to collapse in the point-vortex system ..... 15  
 Takeshi Gotoda (Nagoya Univ.) Numerical study of initial configurations leading to collapse in the point-vortex system

概要 Self-similar collapse is a notable phenomena in the point vortex system. The structure of initial configurations leading to self-similar collapses has been made clear for the 3-point vortex problem and some examples of self-similar collapsing solutions for the 4 and 5-point vortex problem have been obtained. In this study, we investigate initial configurations leading to self-similar collapse for the N-point vortex problem by using numerical simulations and compare some properties of them by changing the number of N. We also show the existence of relative equilibria forming symmetrical configurations.

- 28 田内大渡 (東大数理) 軸対称な楕円体上の非圧縮 Euler 流に対する共役点の存在について ..... 15  
 米田 剛 (東大数理)  
 Taito Tauchi (Univ. of Tokyo) Existence of a conjugate point in the incompressible Euler flow on an ellipsoid  
 Tsuyoshi Yoneda (Univ. of Tokyo)

概要 Existence of a conjugate point in the incompressible Euler flow on a sphere and an ellipsoid is considered. Misiólek (1996) formulated a differential-geometric criterion (we call M-criterion) for the existence of a conjugate point in a fluid flow. In this talk, it is shown that no zonal flow (stationary Euler flow) satisfies M-criterion if the background manifold is a sphere, on the other hand, there are zonal flows satisfy M-criterion if the background manifold is an ellipsoid (even it is sufficiently close to the sphere). The conjugate point is created by the fully nonlinear effect of the inviscid fluid flow with differential geometric mechanism.

- 29 寺本 敬 (旭川医科大医) Pinned pulse solutions inside a bump type heterogeneity ..... 10  
 P. van Heijster  
 (Queensland Univ. of Tech.)  
 Takashi Teramoto Pinned pulse solutions inside a bump type heterogeneity  
 (Asahikawa Medical Univ.)  
 Peter van Heijster  
 (Queensland Univ. of Tech.)

概要 We analyse pinned front and pulse solutions in a singularly perturbed three-component FitzHugh-Nagumo model with a bump type heterogeneity. We derive explicit conditions for the existence and stability of pinned solutions by combining geometric singular perturbation techniques and an action functional approach. We show that the model in the 1 activator - 2 inhibitors framework only supports symmetric pinned pulse solutions that are pinned in the middle of the bump region.

- 30 久保隆徹 (お茶の水女大基幹) 時間遅れを考慮に入れた Burgers 方程式の時間大域解について ..... 15  
 上田好寛 (神戸大海事)  
 Takayuki Kubo (Ochanomizu Univ.) On global in time solution to Burgers equation with a time delay  
 Yoshihiro Ueda (Kobe Univ.)

概要 We consider the Burgers equation with a time delay, which is considered to be a model of traffic flow. We show the existence theorem for global in time solution to Burgers equation with a time delay. The key lemma of the proof of main theorem is the existence theorem for local in time solution to our problem and its a priori estimates.

- 31 渡辺 樹 (早大理工) Central limit theorem for data-diffusion with linear reactions ..... 15  
 豊泉 洋 (早大理工)  
 Itsuki Watanabe (Waseda Univ.) Central limit theorem for data-diffusion with linear reactions  
 Hiroshi Toyozumi (Waseda Univ.)

**概要** We compare two mathematical models of data-diffusion; (1) the deterministic model governed by a reaction-diffusion equation and (2) the stochastic model governed by a multi-dimensional jump Markov process, respectively. In this talk, by scaling the state and fluidizing data-pieces size, we show that the difference of two mathematical models weakly converges to the Ornstein-Uhlenbeck process on Skorokhod space, i.e., it is proved that the central limit theorem holds for these models.

### 16:00~17:00 特別講演

- 李 聖林 (広島大理) 反応拡散方程式, 生命のパターン形成におけるその無限の才能  
 Sungrim Seirin Lee (Hiroshima Univ.) Reaction-diffusion equation, its infinite talent in pattern formation of life science

**概要** Since A. Turing's landmark paper (1952), reaction-diffusion equations became a symbol of mathematical model for pattern formation in life science. A short-range self-activation and long-range inhibition have been considered as a fundamental mechanism by which a symmetry breaking is triggered and a pattern, namely, non-constant steady state, is formed in reaction-diffusion systems. And many mathematical models in pattern formation also have based on it. However, once we take a step back and look at the truly pure essence of reaction-diffusion equation(s), we can find more potential of reaction-diffusion equation(s) and see various patterning dynamics in a simple reaction-diffusion equation(s) without a basis of activation-inhibition chemical reactions. In this talk, I will introduce such an infinite potential of reaction-diffusion equation(s) as pattern forming system which has been done by myself during these 10 years. I also would like to discuss how mathematical modeling (or mathematical modeler) in life science and pure mathematics (or pure mathematician) can be collaborated and open a new trail of applied mathematics.

3月19日(木) 第V会場

### 9:15~10:50

- 32 堀口俊二 ニュートン法の2項展開と収束比較 ..... 10  
 Shunji Horiguchi Binomial expansions of Newton's method and comparison of convergence

**概要** We give the binomial expansion of Newton method from the first term to  $m$  (natural number of more than 2) term, and show that the convergences become the quadratic and linearly (Theorem 1.7). Next in case of the quadratic convergence, we give the convergence comparison of the binomial expansion of Newton method from the first term to  $m$  term and Newton method (Theorem 1.8 (1.13), (1.14)). Next we give the convergence comparison of the binomial expansion of the Newton method from the first term to the second term and from the first term to more than three term (Theorem 1.8 (1.15)). And we give convergence comparisons of the binomial expansion of Newton method from the first term to  $m$  term and Newton method using the curvatures, and concaves and convexes of curves (Theorem 2.3-2.9).

- 33 堀口俊二 ニュートン法の2項展開の数値計算の例 ..... 10  
 Shunji Horiguchi Examples of numerical calculations of the binomial expansions of Newton's method

概要 We give ranges of  $x$  satisfying the formulas (1.13), (1.14) and (1.15) of the convergence comparisons of Theorem 1.8 for  $f(x)=(x-1)(x-2)$ . Next in case of the quadratic convergence, we do numerical calculations of the binomial expansion of Newton method for  $f(x)$  from the first term to  $m$  ( $m=2,3,4$ ) term, and we give the comparisons of convergences of them. These are the confirmations of formulas (1.13), (1.14) and (1.15) of Theorem 1.8. We give ranges of  $x$  satisfying the formulas of Theorem 2.3, 2.4, 2.5 of the convergence comparisons by the curvatures of the binomial expansions of Newton method for  $f(x)$ . And we do numerical calculations of these formulas for confirmations.

- 34 坂口文則(福井大工) どこまで広がる? 微分方程式整数型解法の適用範囲—整数の四則演算だけでどこまでできるか— ..... 15  
 Fuminori Sakaguchi (Univ. of Fukui) A possibility of wider application of an algorithm for solving ODEs by means only of four arithmetical operations among integers

概要 A kind of generalization was proposed by the author for an integer-type algorithm for solving higher-order linear ODEs, which was proposed by the author and M. Hayashi several years ago, by means of algebraic extensions of the field of rational functions. By this generalization, for example, we can solve the higher-order linear ODEs whose coefficient functions are general algebraic functions, by means only of four arithmetical operations among integers. In this study, we show that the range of application of this algorithm can be expanded into a special class of the cases where the coefficient functions of ODEs involve exponential, trigonometric and hyperbolic functions. Moreover, some successful numerical examples are given for ODEs whose coefficient functions involve, for example, sigmoid and tangent functions.

- 35 穴田浩一(早大高等学院) 曲線短縮問題に現れる準線形放物型偏微分方程式に対する爆発解の漸近挙動に関する一考察 ..... 15  
 石渡哲哉  
 (芝浦工大システム理工)  
 牛島健夫(東京理大理工)  
 Koichi Anada A remark on asymptotic behavior of blow-up solutions to a quasi-linear parabolic equation for a curve shortening problem  
 (Waseda Univ. Senior High School)  
 Tetsuya Ishiwata  
 (Shibaura Inst. of Tech.)  
 Takeo Ushijima (Tokyo Univ. of Sci.)

概要 In this talk, we consider asymptotic behavior of blow-up solutions to a quasi-linear parabolic equation  $v_t = v^\delta(v_{\theta\theta} + v)$  for a curve shortening problem. It is known that solutions blow up regionally. Our purpose is to investigate a relation between behavior of solutions at the maximum point and ones on the boundary of the blow-up set.

- 36 木下武彦 有界作用素のレゾルベントに対するある近似作用素の強収束性について ..... 15  
 (九大情報基盤研究開発センター)  
 渡部善隆  
 (九大情報基盤研究開発センター)  
 中尾充宏(早大理工)  
 Takehiko Kinoshita (Kyushu Univ.) On the strong convergence of some approximate operators for resolvents of bounded operators  
 Watanabe Yoshitaka (Kyushu Univ.)  
 Mitsuhiro T. Nakao (Waseda Univ.)

概要 We consider the convergence of some approximate operators for resolvents of bounded linear operators. We report that this approximate operators strong converge to resolvents under reasonable assumptions. Moreover, the operator norm of this approximate operators also converge to the operator norm of resolvents.

- 37 高安亮紀 (筑波大システム情報) Rigorous numerics for nonlinear heat equations in the complex plane of  
 J.-P. Lessard ( McGill Univ. ) time ..... 15  
 J. Jaquette (Brandeis Univ.)  
 岡本久 (学習院大理)  
 Akitoshi Takayasu (Univ. of Tsukuba) Rigorous numerics for nonlinear heat equations in the complex plane of  
 Jean-Philippe Lessard (McGill Univ.) time  
 Jonathan Jaquette (Brandeis Univ.)  
 Hisashi Okamoto (Gakushuin Univ.)

概要 In this talk, we give two computer-assisted proofs for Cauchy problems of nonlinear heat equations for complex time values. The proofs are obtained by rigorous numerics, via a careful blend of functional analysis, semi-group theory, numerical analysis, fixed point theory, the Lyapunov–Perron method and interval arithmetic.

#### 11:00~12:00 特別講演

- 尾崎克久 行列積に対するエラーフリー変換: 基礎・応用・未来  
 (芝浦工大システム理工)  
 Katsuhisa Ozaki Error-free transformation for matrix multiplication: Basic, applications,  
 (Shibaura Inst. of Tech.) and future

概要 This talk concerns the numerical computations of matrix multiplication. Floating-point numbers and floating-point arithmetic defined in IEEE 754 are widely used in numerical computations. The performance of the numerical computation is very high. On the other hand, the problem of rounding errors is crucial. We proposed an error-free transformation of matrix multiplication. The matrix product is transformed into an unevaluated sum of floating-point matrices. This technique is useful for accurate numerical computations. Besides, the error-free transformation is applied to interval matrix multiplication and generation of test matrices in numerical linear algebra. Moreover, we can develop reproducible numerical algorithms and a fast algorithm for matrix multiplication using low precision arithmetic on GPGPU based on the error-free transformation.



## トポロジー

3月16日(月) 第IV会場

10:00~12:00

- 1 堀内 遼 (名大多元数理)<sup>b</sup> Verschiebung maps among  $K$ -groups of truncated polynomial algebras ..... 10  
 Ryo Horiuchi (Nagoya Univ.) Verschiebung maps among  $K$ -groups of truncated polynomial algebras

概要 Taking the sphere spectrum as base instead of integers gives Hochschild homology a structure called cyclotomic structure. Using the structure we evaluate certain maps between algebraic  $K$ -groups of some truncated polynomial algebras in terms of de Rham–Witt complex.

- 2 狩野隼輔 (東工大数理) Algebraic entropy of sign-stable mutation loops ..... 15  
 石橋 典 (東大数理)  
 Shunsuke Kano (Tokyo Tech) Algebraic entropy of sign-stable mutation loops  
 Tsukasa Ishibashi (Univ. of Tokyo)

概要 We introduce a property of mutation loops, called the sign stability, with a focus on an asymptotic behavior of the iteration of the tropical  $\mathcal{X}$ -transformation. A sign-stable mutation loop has a numerical invariant which we call the cluster stretch factor, in analogy with that of a pseudo-Anosov mapping class on a marked surface. We compute the algebraic entropies of the cluster  $\mathcal{A}$ - and  $\mathcal{X}$ -transformations induced by a sign-stable mutation loop, and conclude that these two coincide with the logarithm of the cluster stretch factor.

- 3 足助太郎 (東大数理)<sup>b</sup> 葉層構造の Fatou 集合について ..... 15  
 Taro Asume (Univ. of Tokyo) On Fatou sets of foliations

概要 We will discuss a criteria for finding Fatou sets of foliations.

- 4 森 淳秀 (大阪歯大歯) Bayes 推定の幾何 ..... 15  
 Atsuhide Mori (Osaka Dental Univ.) Geometry of Bayesian estimation

概要 In this talk, I would like to propose a purely geometric description of Bayesian estimation. We generalize the notion of relative entropy and Fisher information so that they become subjects of Lorentzian geometry and further those of the general relativity. In fact, we give an example of Ricci flat Lorentzian manifold appearing as an extended space of normal distributions.

- 5 北野晃朗 (創価大理工) Twisted Alexander polynomials of torus links ..... 10  
 森藤孝之 (慶大経済)  
 A. T. Tran  
 (Univ. Texas at Dallas)

Teruaki Kitano (Soka Univ.) Twisted Alexander polynomials of torus links  
 Takayuki Morifuji (Keio Univ.)  
 Anh T. Tran (Univ. Texas at Dallas)

概要 We give an explicit formula of the twisted Alexander polynomial of the torus link and show that it is a locally constant function on the  $SL(2; \mathbb{C})$ -character variety. We also discuss the similar things for higher dimensional twisted Alexander polynomials.

- 6 野崎雄太 (明大研究・知財) Abelian quotients of the  $Y$ -filtration on the homology cylinders via the  
佐藤正寿 (東京電機大未来) LMO functor ..... 15  
鈴木正明 (明大総合数理)  
Yuta Nozaki (Meiji Univ.) Abelian quotients of the  $Y$ -filtration on the homology cylinders via the  
Masatoshi Sato (Tokyo Denki Univ.) LMO functor  
Masaaki Suzuki (Meiji Univ.)

**概要** We construct a homomorphism on the graded quotient of the  $Y$ -filtration of the homology cylinders. The homomorphism restricted to the lower central series of the Torelli group is non-trivial and different from the Johnson homomorphisms. As an application, we give an abelian quotient of the Johnson kernel which is not described by the Casson invariant, the second and third Johnson homomorphisms. This is the joint work with Masatoshi Sato and Masaaki Suzuki.

- 7 丹下稜斗 (東京電機大数学) Twisted Alexander polynomials of hyperbolic twist knots and von Dyck  
groups ..... 10  
Ryoto Tange (Tokyo Denki Univ.) Twisted Alexander polynomials of hyperbolic twist knots and von Dyck  
groups

**概要** We study twisted Alexander polynomials of hyperbolic twist knots for non-abelian  $SL_2(\mathbb{C})$ -representations and discuss some relations with von Dyck groups.

- 8 野坂武史 (東工大理) 結び目の  $K_1$ -値振れ Alexander 多項式 ..... 15  
Takefumi Nosaka (Tokyo Tech)  $K_1$ -Alexander twisted polynomials of knots

**概要** Given a homomorphism from a knot group to a group, we introduce a  $K_1$ -class, which is a generalization of the 1-variable Alexander polynomial. We suggest several approaches to the  $K_1$ -class. As a corollary, we show a relation to Reidemeister torsions of finite cyclic covering spaces, and show reciprocity in some senses.

#### 14:15~15:15 特別講演

- 森藤孝之 (慶大経済) 双曲結び目・絡み目のねじれ Alexander 多項式  
Takayuki Morifuji (Keio Univ.) Twisted Alexander polynomials of hyperbolic knots and links

**概要** The twisted Alexander polynomial is defined for a pair of the group and its representation. It is a natural generalization of the Alexander polynomial and gives a powerful tool in the study of low-dimensional topology. Based on huge numerical calculations, Dunfield, Friedl and Jackson have proposed a conjecture that the twisted Alexander polynomial associated to the holonomy representation determines the genus and fiberedness of a hyperbolic knot. In this talk we will survey recent results on the conjecture and explain its generalization to hyperbolic links.

**15:30~16:30 特別講演**

- 藤田直樹 (東大数理) Newton–Okounkov 凸体およびクラスター構造から生じるトーリック退化  
 Naoki Fujita (Univ. of Tokyo) Toric degenerations arising from Newton–Okounkov bodies and cluster structures

**概要** The theory of toric varieties gives an elegant dictionary translating geometric and topological properties into combinatorial properties in terms of cones and polytopes. In order to apply this powerful dictionary to other projective varieties, we can use degenerations to toric varieties, called toric degenerations. In this talk, we study the following two systematic methods of constructing toric degenerations: Newton–Okounkov bodies and cluster algebras. In the first part of this talk, we survey the theory of Newton–Okounkov bodies and its applications in geometry, including Harada–Kaveh’s construction of completely integrable systems. In the second part, we study Newton–Okounkov bodies from the theory of cluster algebras. We can realize Gross–Hacking–Keel–Kontsevich’s toric degenerations of compactified cluster varieties as ones coming from Newton–Okounkov bodies. Throughout this talk, we especially focus on the case of flag varieties whose toric degenerations are closely related to representation theory.

3月17日(火) 第IV会場

**10:00~12:00**

- 9 石井 敦 (筑波大数理物質) 多重群ラックと有向空間曲面 ..... 10  
 松崎 尚作 (拓殖大工)  
 村尾 智 (筑波大数理物質)  
 Atsushi Ishii (Univ. of Tsukuba) A multiple group rack and oriented spatial surfaces  
 Shosaku Matsuzaki (Takushoku Univ.)  
 Tomo Murao (Univ. of Tsukuba)

**概要** In this talk, we define a multiple group rack, which is an algebraic structure consisting of the disjoint union of groups. We introduce a coloring invariant for oriented spatial surfaces by using multiple group racks, and give some examples.

- 10 石井 敦 (筑波大数理物質) The fundamental multiple conjugation quandle of a handlebody-knot ..... 10  
 Atsushi Ishii (Univ. of Tsukuba) The fundamental multiple conjugation quandle of a handlebody-knot

**概要** A multiple conjugation quandle is used to define a coloring invariant of a handlebody-knot. We introduce a presentation of a multiple conjugation quandle and define the fundamental multiple conjugation quandle of a handlebody-knot.

- 11 小林 竜馬 (石川工高専) 向き付け不可能曲面の写像類群とそのツイスト部分群の無限表示 ..... 15  
 大森 源城 (東京理大理工)  
 Ryoma Kobayashi Infinite presentations for the mapping class group and its twist subgroup  
 (Ishikawa Nat. Coll. of Tech.) of a compact non-orientable surface  
 Genki Omori (Tokyo Univ. of Sci.)

**概要** A finite presentation for the subgroup of the mapping class group of a compact non-orientable surface generated by all Dehn twists was given by Stukow. In this work, we give an infinite presentation for this group, mainly using the presentation given by Stukow and Birman exact sequences on mapping class groups of non-orientable surfaces. As an application, we give an infinite presentation for the mapping class group of a compact non-orientable surface.

- 12 大森源城 (東京理大理工) 種数 4 と 5 の場合の向き付け不可能曲面上の involution の Dehn twist-  
 阪田直樹 (埼玉大理工) crosscap slide 表示について ..... 15  
 Genki Omori (Tokyo Univ. of Sci.) Dehn twist-crosscap slide presentations for involutions on non-orientable  
 Naoki Sakata (Saitama Univ.) surfaces of genus 4 and 5

概要 Lickorish proved that any element of the mapping class group of a non-orientable is a product of Dehn twists and crosscap slides. We call the product Dehn twist-crosscap slide presentation for the element. In this talk, we give Dehn twist-crosscap slide presentations for all conjugacy classes of involutions on non-orientable surfaces of genus 4 and 5. The Dehn twist-crosscap slide presentations are constructed by products of Szepietowski's finite generating set.

- 13 浮田卓也 (東工大 理) Genus zero PALF structures on the Akbulut–Yasui plugs ..... 10  
 Takuya Ukida (Tokyo Tech) Genus zero PALF structures on the Akbulut–Yasui plugs

概要 We construct a genus zero PALF structure on each of plugs introduced by Akbulut and Yasui and describe the monodromy as a positive factorization in the mapping class group of a fiber. We also examine the monodromies of PALFs on a certain pair of compact Stein surfaces such that one is obtained by applying a plug twist to the other.

- 14 浅野喜敬 (東北大理) Vertical 3-manifolds in simplified genus 2 trisections of 4-manifolds ... 15  
 Nobutaka Asano (Tohoku Univ.) Vertical 3-manifolds in simplified genus 2 trisections of 4-manifolds

概要 A trisection is a decomposition of a closed 4-manifold by 3 tuple of 4-dimensional 1-handlebodies, which was introduced by Gay and Kirby. They proved the existence of a trisection for any closed 4-manifold by using a stable map (called a trisection map) from the 4-manifold to  $\mathbf{R}^2$ . After their work, a simplified trisection was introduced by Baykur and Saeki. They proved the existence of a simplified trisection from a simplified broken Lefschetz fibration. In this talk, we will give a classification of 3-manifolds that can be obtained as the preimages of arcs on  $\mathbf{R}^2$  by simplified genus 2 trisection maps, which we call vertical 3-manifolds.

- 15 谷口正樹 (東大数理) 2次元結び目の Seifert 超曲面と Chern–Simons 汎関数について ..... 15  
 Masaki Taniguchi (Univ. of Tokyo) Seifert hypersurfaces of 2-knots and Chern–Simons functional

概要 In this talk, we treat oriented 2-knots: smooth embeddings from  $S^2$  to  $S^4$  equipped with orientations. For a given oriented 2-knot, we introduce a functional from the  $SU(2)$ -representation space of its knot group to  $(0, 1]$ . The image of the functional gives an isotopy invariant of an oriented 2-knot. We show several properties of the functional, including a connected sum formula, a relation between the functional and  $SU(2)$ -representation, and an orientation reversing formula. Moreover, we calculate the images of the functionals for ribbon 2-knots and twisted spun knots of torus knots, rational knots, and Montesinos knots. Using a refinement of instanton Floer theory and several properties of the functional, we relate the existence of a certain class of Seifert hypersurface with the existence of an irreducible  $SU(2)$ -representation of its knot group.

- 16 今井 淳 (千葉大理) 単一曲線の自己インダクタンス ..... 15  
 Jun O'Hara (Chiba Univ.) Regularization of self-inductance

概要 We define the regularized self-inductances of a knot from Neumann formula and Weber formula of the mutual inductance of a pair of closed circuits. We show that they are different by twice the length. We also study the residues and related quantities.

**13:15~14:15 特別講演**

今野 北斗 (理化学研) Gauge theory and diffeomorphism and homeomorphism groups  
 Hokuto Konno (RIKEN) Gauge theory and diffeomorphism and homeomorphism groups

概要 I will survey recent development of applications of gauge theory to the diffeomorphism and homeomorphism groups of 4-manifolds.

It is known that, for an arbitrary closed manifold of dimension less than 4, the inclusion map from the diffeomorphism group to the homeomorphism group is a weak homotopy equivalence. Therefore it is interesting to ask if there is a homotopical difference between the diffeomorphism and homeomorphism groups in other dimensions. Thanks to one of Donaldson's results, it turned out that 4 is the first dimension in which such a difference appears. But until recently, no one had have any systematic way to extract such differences in dimension 4. This talk is aimed at explaining that gauge theory for families gives a systematic way as follows.

After Donaldson's celebrated diagonalization theorem, gauge theory has given strong constraints on the topology of smooth 4-manifolds, such as Furuta's 10/8-inequality. Combining such constraints with Freedman's theory, one may find many non-smoothable topological 4-manifolds. Recently, a family version of this argument was started by T. Kato, N. Nakamura and myself, and soon later it was developed also by D. Baraglia and his collaborating work with myself. More precisely, considering gauge theory for smooth fiber bundles of 4-manifolds, they obtained some constraints on the topology of smooth fiber bundles of 4-manifolds. Moreover, it turned out that one may detect non-smoothable topological fiber bundles of smooth 4-manifolds using such constraints. The existence of such bundles implies that there are homotopical differences between the diffeomorphism and homeomorphism groups of the 4-manifolds given as the fibers of the non-smoothable bundles.

3月18日(水) 第IV会場

**10:00~12:00**

17 石川 勝巳 (京大数理研) Minimal coloring numbers on minimal diagrams of torus links ..... 10  
 市原 一裕 (日大文理)  
 松土 恵理 (日大文理)

Katsumi Ishikawa (Kyoto Univ.) Minimal coloring numbers on minimal diagrams of torus links  
 Kazuhiro Ichihara (Nihon Univ.)  
 Eri Matsudo (Nihon Univ.)

概要 In this talk, we determine the minimal number of colors for non-trivial  $\mathbb{Z}$ -colorings on the standard minimal diagrams of  $\mathbb{Z}$ -colorable torus links. Also included are complete classifications of such  $\mathbb{Z}$ -colorings and of such  $\mathbb{Z}$ -colorings by only four colors.

18 大山口菜都美 (秀明大学校教師) Pallets of Dehn  $p$ -coloring for spatial graphs ..... 10  
 大城佳奈子 (上智大理工)

Natsumi Oyamaguchi (Shumei Univ.) Pallets of Dehn  $p$ -coloring for spatial graphs  
 Kanako Oshiro (Sophia Univ.)

概要 A Dehn  $p$ -coloring for a spatial graph diagram is an assignment of an element of  $\mathbb{Z}_p = \{0, \dots, p-1\}$  to each region. At each crossing, some coloring condition is satisfied. A region pallet of  $\mathbb{Z}_p$  is a subset of  $\bigcup_{n \in 2\mathbb{Z}_+} \mathbb{Z}_p^n$  which gives a coloring condition at each vertex for Dehn  $p$ -colorings. We classified the region pallets of  $\mathbb{Z}_p$ .

- 19 M. Eudave-Muñoz <sup>b</sup> The maximum and minimum genus of a multibranch surface ..... 15  
 (Univ. Nacional Autónoma de México)  
 小 沢 誠 (駒澤大総合)  
 Mario Eudave-Muñoz The maximum and minimum genus of a multibranch surface  
 (Univ. Nacional Autónoma de México)  
 Makoto Ozawa (Komazawa Univ.)

概要 In this talk, we give a lower bound for the maximum and minimum genus of a multibranch surface by the first Betti number and the minimum and maximum genus of the boundary of the neighborhood of it respectively. As its application, we show that the maximum and minimum genus of  $G \times S^1$  is equal to twice of the maximum and minimum genus of  $G$  for a graph  $G$  respectively. This is a joint work with Mario Eudave-Muñoz.

- 20 伊藤 昇 (東大数理) The tabulation of prime knot projections with their mirror images up  
 瀧村 祐介 (学習院中) to eight double points ..... 10  
 Noboru Ito (Univ. of Tokyo) The tabulation of prime knot projections with their mirror images up  
 Yusuke Takimura to eight double points  
 (Gakushuin Boys' Junior High School)

概要 We show how to tabulate knot projections by listing tangles. In particular, we obtain the complete table of prime knot projections with their mirror images up to eight double points systematically through a finite procedure.

- 21 市原 一裕 (日大文理) Two-bridge knots admit no purely cosmetic surgeries ..... 15  
 斎藤 敏夫 (上越教育大)  
 鄭 仁大 (近畿大理工)  
 T. W. Mattman  
 (California State Univ., Chico)  
 Kazuhiro Ichihara (Nihon Univ.) Two-bridge knots admit no purely cosmetic surgeries  
 Toshio Saito (Joetsu Univ. of Edu.)  
 In Dae Jong (Kindai Univ.)  
 Thomas W. Mattman  
 (California State Univ., Chico)

概要 We show that two-bridge knots and alternating fibered knots admit no purely cosmetic surgeries, i.e., no pair of distinct Dehn surgeries on such a knot produce 3-manifolds that are homeomorphic as oriented manifolds. Our argument, based on a recent result by Hanselman, uses several invariants of knots or 3-manifolds; for knots, we study the signature and some finite type invariants, and for 3-manifolds, we deploy the  $SL(2, \mathbb{C})$  Casson invariant.

- 22 中江 康晴 (秋田大理工) 種数1ファイバー結び目に沿ったデーン手術と基本群の左順序付け可能  
 市原 一裕 (日大文理) 性について ..... 15  
 Yasuharu Nakae (Akita Univ.) Dehn surgeries along genus one fibered knots and left-orderability of  
 Kazuhiro Ichihara (Nihon Univ.) fundamental groups

概要 We study Dehn surgeries along genus one fibered knots and which resultant manifolds have left-orderable fundamental groups. The 3-sphere has only two genus one fibered knots, the trefoil and the figure-eight knot. In contrast with the 3-sphere, some lens spaces have more genus one fibered knots. In 2014, Baker completely classified the number of genus one fibered knots in lens spaces. Along the classification, we show that, on some class of such knots in lens spaces, all integral surgeries yield 3-manifolds with left-orderable fundamental groups. In order to prove this theorem, we examine the existence of Anosov flow whose stable/unstable foliation is  $\mathbb{R}$ -covered in the resultant manifold.

- 23 田中利史 (岐阜大教育) 対称和表示をもつサテライト結び目について ..... 10  
 Toshifumi Tanaka (Gifu Univ.) On satellite knots with symmetric union presentations

概要 A symmetric union in the 3-space is a knot, obtained from a knot and its mirror image, which are symmetric with respect to a 2-plane, by taking the connected sum of them and moreover by connecting them with some vertical twists along the plane. A symmetric union is an example of a ribbon knot. In this talk, we show that if a knot is a composite symmetric union with minimal twisting number one, then the knot has a non-trivial knot and its mirror image as connected summands. We also show that a satellite symmetric union with minimal twisting number one such that the order of the pattern is an odd number  $\geq 3$  has at least two disjoint non-parallel essential tori in the complement.

- 24 安部哲哉 (立命館大理工) Table of annulus presentations of knots ..... 10  
 田神慶士 (水産大)  
 Tetsuya Abe (Ritsumeikan Univ.) Table of annulus presentations of knots  
 Keiji Tagami (Nat. Fisheries Univ.)

概要 We give a table of knots which have annulus presentations up to 8-crossings.

- 25 伊藤哲也 (京大理) 閉組みひも表示の無限性 ..... 10  
 Tetsuya Itoh (Kyoto Univ.) Infiniteness of closed braid representatives

概要 We show that if a link  $L$  has a closed  $n$ -braid representative admitting a non-degenerate exchange move, an exchange move that does not obviously preserve the conjugacy class,  $L$  has infinitely many non-conjugate closed  $n$ -braid representatives.

## 無限可積分系

3月18日(水) 第IX会場

## 14:15~16:15

- 1 平出 耕一 (愛媛大理) 複素エノン写像の力学系にあらわれる Stokes 的現象 ..... 15  
松岡 千博 (阪市大工)  
Koichi Hiraide (Ehime Univ.) Stokes-like phenomena which appear in dynamics of complex Henon  
Chihiro Matsuoka (Osaka City Univ.) maps

概要 We consider invariant curves for complex Henon maps associated with an eigenvalue, with absolute value not equal to one, of the derivative at a fixed point. By the method of Borel–Laplace transform, we can construct pairs  $(x_i(t), y_i(t))$  of functions, parameterizing the invariant curves. Such functions have forms of asymptotic expansions. In this talk, we give a relation with classical functions, due to Poincare, represented by Taylor series, and state a connection structure among functions' pairs  $(x_i(t), y_i(t))$ , which is like structures known as (nonlinear) Stokes phenomena in differential equations.

- 2 松浦 望 (久留米工大) 平面離散弾性曲線の明示公式 ..... 15  
Nozomu Matsuura Explicit formula for planar discrete elasticae  
(Kurume Inst. of Tech.)

概要 We give an explicit formula for the discrete elasticae in the Euclidean plane in terms of Jacobi's elliptic theta functions.

- 3 田所 勇樹 (木更津工高専) Nonlinear  $O(3)$  sigma model in discrete complex analysis ..... 15  
関口 昌由 (木更津工高専)  
鎌田 勝 (木更津工高専\*)  
Yuuki Tadokoro Nonlinear  $O(3)$  sigma model in discrete complex analysis  
(Nat. Inst. of Tech., Kisarazu Coll.)  
Masayoshi Sekiguchi  
(Nat. Inst. of Tech., Kisarazu Coll.)  
Masaru Kamata  
(Nat. Inst. of Tech., Kisarazu Coll.\*)

概要 We study a discrete version of the two-dimensional nonlinear  $O(3)$  sigma model derived from discrete complex analysis. We show the discrete version of an equality for the energy of this model.

- 4 桑野 泰宏 中心力場におけるルンゲ・レンツ様ベクトル ..... 15  
(鈴鹿医療科学大医用工)  
Yas-Hiro Quano Runge–Lenz like vectors for central force fields  
(Suzuka Univ. of Med. Sci.)

概要 In this talk, two-body problems for classical central force fields will be discussed. As is well known, there exists a conserved Runge–Lenz vector in addition to conserved quantities, the energy and the angular momentum.

I consider general central force problems, and show the existence of Runge–Lenz like vectors. In particular, I discuss a harmonic oscillator and a perturbed Kepler problem with a  $1/r^2$  potential in detail.

- 5 白石 潤一 (東大数理) 非定常 Ruijsenaars 関数 ..... 15  
Junichi Shiraishi (Univ. of Tokyo) Non-stationary Ruijsenaars function

概要 We introduce the formal power series  $f^{\hat{g}^N}(x, p|s, \kappa|q, t)$  which we call the non-stationary Ruijsenaars function. We present several theorems and conjectures concerning the non-stationary Ruijsenaars function.



- 6 白石潤一(東大数理) 非定常及び定常 Ruijsenaars 関数と Ruijsenaars 作用素の固有値問題 . . . 15  
Junichi Shiraishi (Univ. of Tokyo) Non-stationary and stationary Ruijsenaars functions and eigenvalue problem associated with Ruijsenaars operator

概要 Based on the non-stationary and the stationary Ruijsenaars functions, we present a conjecture concerning the eigenvalue problem associated with the Ruijsenaars operator.

- 7 太久保勇輔(東大数理) 非定常 Ruijsenaars 関数と Ding–Iohara–Miki 代数の intertwining 作用素  
白石潤一(東大数理) . . . . . 15  
福田真之(東大理)  
Yusuke Ohkubo (Univ. of Tokyo) Non-stationary Ruijsenaars functions and intertwining operators of the  
Jun'ichi Shiraishi (Univ. of Tokyo) Ding–Iohara–Miki algebra  
Masayuki Fukuda (Univ. of Tokyo)

概要 In this talk, we realize the affine analogue of the Macdonald functions, which are referred to as the non-stationary Ruijsenaars functions, by the Ding–Iohara–Miki algebra. Moreover, we present various combinatorial expressions of the non-stationary Ruijsenaars functions by using a duality formula for the intertwining operators of the Ding–Iohara–Miki algebra.

### 16:30~17:30 特別講演

- 渡邊英也(京大数理研)  $\imath$ quantizations  
Hideya Watanabe (Kyoto Univ.)  $\imath$ quantizations

概要  $\imath$ quantum groups are generalizations of quantum groups appearing as one-sided coideal subalgebras of quantum groups in the theory of quantum symmetric pairs. Recently, numerous results which we should call “ $\imath$ quantizations” have been reported.  $\imath$ quantizations have two meanings; (1) generalizing what is known for usual quantum groups to  $\imath$ quantum groups, (2) quantizing what is known for classical Lie algebras but cannot be quantized by usual quantum groups, by means of  $\imath$ quantum groups. In this talk, I introduce examples of  $\imath$ quantizations such as canonical basis, geometric construction, Hall algebraic construction, GT-basis of type B/D, Schur duality of type B/C/D, highest weight theory, and crystal basis.

3月19日(木) 第IX会場

### 9:45~11:30

- 8 森山翔文(阪市大理) 量子曲線とワイル群 . . . . . 15  
Sanefumi Moriyama (Osaka City Univ.) Quantum curves and Weyl groups

概要 Quantum curves are helpful for studying the Painlevé equations or many other mathematical physics equations such as the correspondence between spectral theories and topological strings. We provide explicit expressions for the quantum curves of genus one and the actions of Weyl group. We further comment on the applications of these curves to the analysis of mirror maps.

- 9 高崎金久(近畿大理工)  $CP^1$  の同変 Gromov–Witten 理論と同変戸田階層 . . . . . 15  
Kanehisa Takasaki (Kindai Univ.) Equivariant Gromov–Witten theory of  $CP^1$  and equivariant Toda hierarchy

概要 Okounkov and Pandharipande constructed what they call “dressing operators” in their study on the integrable structure of the Gromov–Witten theory of  $CP^1$ . These operators are used to convert a generating function of the equivariant Gromov–Witten invariants to a tau function of the equivariant Toda hierarchy. We redefine these operators in the perspective of the Lax formalism of the 2D Toda hierarchy. This leads to yet another proof of the fact that the generating function of the equivariant Gromov–Witten invariants yields a special solution of the equivariant Toda hierarchy.

- 10 山根 宏之 (富山大理) 一般化された量子群の典型的既約指標について ..... 15  
 Hiroyuki Yamane (Univ. of Toyama) On typical irreducible characters of generalized quantum groups

概要 In 1977, V. Kac gave the Weyl-type character formula of the typical finite dimensional irreducible modules of the basic classical Lie superalgebras. In the talk, we introduce an analogous character formula for the generalized quantum groups.

- 11 藤田 遼 (京大理)  $ADE$  型アフィン量子群の基本表現の間の正規化された  $R$  行列の特異性について ..... 15  
 Ryo Fujita (Kyoto Univ.) Singularities of normalized  $R$ -matrices between fundamental modules over the affine quantum groups of type  $ADE$

概要 We present a simple unified formula expressing the denominators of the normalized  $R$ -matrices between the fundamental modules over the affine quantum groups of type  $ADE$ . It has an interpretation in terms of representations of Dynkin quivers and can be proved in a unified way using the geometry of graded quiver varieties. As a by-product, we obtain a geometric interpretation of Kang–Kashiwara–Kim’s generalized quantum affine Schur–Weyl duality functor when it arises from a family of fundamental modules.

- 12 行田 康晃 (名大多元数理) 有限型団代数・階数 2 の団代数における  $d$  ベクトルと  $f$  ベクトルの関係式 ..... 15  
 Yasuaki Gyoda (Nagoya Univ.) Relation between  $f$ -vectors and  $d$ -vectors in cluster algebras of finite type or rank 2

概要 We study the  $f$ -vectors, which are the maximal degree vectors of  $F$ -polynomials in cluster algebra theory. When a cluster algebra is of finite type or rank 2, we find that the positive  $f$ -vectors correspond with the  $d$ -vectors, which are exponent vectors of denominators of cluster variables. Furthermore, using this correspondence and properties of  $d$ -vectors, we prove that cluster variables in a cluster are uniquely determined by their  $f$ -vectors when the cluster algebra is of finite type or rank 2.

- 13 大久保直人 (青学大理工) 団代数を用いた  $A_7^{(1)}$  型  $q$ -パンルヴェ系の高階化 ..... 15  
 増田 哲 (青学大理工)  
 津田 照久 (一橋大経済)  
 Naoto Okubo (Aoyama Gakuin Univ.) Cluster algebras and higher order  $q$ -Painlevé systems of type  $A_7^{(1)}$   
 Tetsu Masuda (Aoyama Gakuin Univ.)  
 Teruhisa Tsuda (Hitotsubashi Univ.)

概要 In this talk, we present higher order  $q$ -Painlevé systems of type  $A_7^{(1)}$  by using a cluster algebra.

#### 14:15~16:00

- 14 渋川 元樹 (神戸大理) Another proof of difference equations for interpolation Jack polynomials ..... 15  
 Genki Shibukawa (Kobe Univ.) Another proof of difference equations for interpolation Jack polynomials

概要 We give another proof of difference equations for interpolation Jack polynomials.

- 15 松 縄 竜 弥 (中 大 理 工) Variants of confluent  $q$ -hypergeometric equations ..... 15  
 佐 藤 智 輝 (中 大 理 工)  
 竹 村 剛 一 (お茶の水女大基幹)  
 Ryuya Matsunawa (Chuo Univ.) Variants of confluent  $q$ -hypergeometric equations  
 Tomoki Sato (Chuo Univ.)  
 Kouichi Takemura (Ochanomizu Univ.)

概要 Variants of the  $q$ -hypergeometric equation were introduced in our paper jointly with Hatano. We consider degenerations of the variants of the  $q$ -hypergeometric equation, which is the  $q$ -analogue of confluence of singularities in the setting of the differential equation. We also consider degenerations of solutions to the  $q$ -difference equations.

- 16 川 上 拓 志 (青 学 大 理 工) 4次元 Painlevé 型差分方程式について ..... 15  
 Hiroshi Kawakami On four-dimensional Painlevé-type difference equations  
 (Aoyama Gakuin Univ.)

概要 We focus on Fuchsian equations with four accessory parameters and three singular points. We see that the Fuchsian equations admit a “degeneration scheme” in some sense, which is expected to give rise to a degeneration scheme of discrete isomonodromic deformation equations with four-dimensional phase space. We compute an example of discrete isomonodromic deformation equations of a certain Fuchsian equation.

- 17 川 上 拓 志 (青 学 大 理 工) 行列第六 Painlevé 系の  $q$ -類似 ..... 15  
 Hiroshi Kawakami A  $q$ -analogue of the matrix sixth Painlevé system  
 (Aoyama Gakuin Univ.)

概要 We consider a connection-preserving deformation of a certain linear  $q$ -difference system. A system of non-linear  $q$ -difference equations thus obtained can be regarded as a  $q$ -analogue of the matrix sixth Painlevé system, or as a non-abelian analogue of the  $q$ - $P_{VI}$ .

- 18 伊 藤 雅 彦 (琉 球 大 理) 6 パラメータの  $G_2$  型楕円超幾何積分が満たす  $q$  差分方程式系について ..... 15  
 野 海 正 俊 (神 戸 大 理)  
 Masahiko Ito (Univ. of Ryukyus)  $q$ -Difference system for the elliptic hypergeometric integral of type  $G_2$   
 Masatoshi Noumi (Kobe Univ.) with six parameters

概要 We will present an explicit form of the  $q$ -difference system, which is satisfied by the elliptic hypergeometric integral of type  $G_2$  with six parameters.

- 19 長 尾 秀 人 (明 石 工 高 専) パデ法と  $q$ -quadratic 差分ガルニエ系たち ..... 15  
 山 田 泰 彦 (神 戸 大 理)  
 Hidehito Nagao (Akashi Coll. of Tech.) Padé method and  $q$ -quadratic Garnier systems  
 Yasuhiko Yamada (Kobe Univ.)

概要 We study a certain  $q$ -quadratic Garnier system. Choosing a suitable Padé problem on  $q$ -quadratic grid, we simultaneously construct the evolution equation, the Lax pair and determinant formulae of special solutions in terms of the very-well-poised balanced  $q$ -hypergeometric function. We discuss some specializations of the system.

**16:15~17:15 特別講演**

齊藤 義久 (立教大理) 楯円 Artin 群について

Yoshihisa Saito (Rikkyo Univ.) On elliptic Artin groups

概要 In the study of representation theory of Lie groups and Lie algebras, the regular Weyl group orbit spaces and their fundamental groups (called Artin groups or generalized braid groups) have quite important roles.

In the middle of 80's, motivated by the study of singularity theory, Kyoji Saito introduced the notion of elliptic root systems, and study their basic properties. Especially, he introduced an "elliptic analogue" of the regular Weyl group orbit spaces, so-called the elliptic regular orbit spaces, and study their detailed structure in algebraic and differential geometrical point of view.

In this talk, we study the fundamental groups of the regular elliptic Weyl group orbit spaces. These groups are presented by a generator system associated with the elliptic diagrams, and we call them the elliptic Artin groups. Furthermore, some basic properties of these groups will be also discussed. Especially, the elliptic regular orbit space is defined over the moduli space of elliptic curves. This fact leads us to the description of the elliptic modular group actions on elliptic Artin groups. This talk is based on a joint work with Kyoji Saito.