

🌸 日本数学会

2023年度年会

英文サマリ集

2023年3月

於 中央大学

2023 日本数学会 年会プログラム

期 日 2023年3月15日(水)～3月18日(土)

会 場 中央大学理工学部
東京都文京区春日1-13-27

連絡先 中央大学理工学部数学科
〒112-8551

東京都文京区春日1-13-27

E-mail chuo23mar@mathsoc.jp

一般社団法人 日本数学会

	第I会場 5号館5階 5533教室	第II会場 5号館3階 5333教室	第III会場 5号館3階 5335教室	第IV会場 5号館3階 5336教室	第V会場 5号館2階 5233教室	第VI会場 5号館2階 5234教室	第VII会場 5号館2階 5235教室	第VIII会場 5号館2階 5236教室	第IX会場 5号館1階 5138教室
15日 (水)	代 数 学 9:00～12:00 14:15～14:50	幾 何 学 9:30～11:40 14:15～16:00	トポロジー 9:30～12:00 15:40～17:30	関数方程式論 9:30～12:00 14:15～16:15	函 数 論 9:30～11:45	関数解析学 9:30～12:00 14:15～16:15	応用数学 9:50～11:50 14:15～15:25	統計数学 10:00～11:30	数学基礎論 および歴史 9:30～10:15 14:30～16:45
	企画特別講演 13:00～14:00								
	特別講演 15:00～16:00 16:15～17:15	特別講演 16:15～17:15	特別講演 14:20～15:20	特別講演 16:30～17:30	特別講演 14:15～15:15 15:30～16:30	特別講演 16:30～17:30	特別講演 15:45～16:45	特別講演 14:15～15:15 15:30～16:30	特別講演 10:30～11:30 17:00～18:00
16日 (木)	代 数 学 9:10～12:00 13:00～14:10	幾 何 学 9:30～12:00 特別講演 13:00～14:00	トポロジー 9:30～12:00 13:00～14:00	関数方程式論 9:15～12:00 特別講演 13:00～14:00	函 数 論 9:30～11:45 特別講演 13:00～14:00	関数解析学 9:30～12:00 特別講演 13:10～14:10	応用数学 10:00～11:45 13:00～14:10	統計数学 10:20～11:10	数学基礎論 および歴史 9:15～12:15
	日本数学会賞授賞式 (5号館 5階 5534教室) (14:30～15:05)								
	総 合 講 演 (") 日本数学会賞春季賞受賞者 (15:20～16:20) 日 比 孝 之 (阪 大*) (16:35～17:35)								
17日 (金)	代 数 学 9:00～12:00 14:15～14:55	幾 何 学 9:30～11:40 14:15～14:55	トポロジー 9:30～12:00 15:40～17:00	関数方程式論 9:30～12:00 14:15～16:15	実函数論 9:00～11:45 14:15～15:25	関数解析学 9:30～12:00 14:15～15:30	応用数学 9:50～11:50 14:45～16:30	統計数学 9:50～11:35	無限可積分系 9:45～10:45
	企画特別講演 13:00～14:00								
	特別講演 14:40～15:40 16:00～17:00	特別講演 15:10～16:10	特別講演 14:20～15:20	特別講演 16:30～17:30	特別講演 15:40～16:40 17:00～18:00	特別講演 15:45～16:45	特別講演 16:45～17:45	特別講演 14:25～15:25 15:40～16:40	特別講演 11:00～12:00
18日 (土)	代 数 学 9:15～12:00 14:15～16:45			関数方程式論 9:30～12:00 14:15～16:15	実函数論 9:00～12:00 14:15～16:30		応用数学 9:50～11:50 14:15～15:05	統計数学 10:00～11:10	無限可積分系 10:00～10:45
	企画特別講演 13:00～14:00								
				特別講演 16:30～17:30	特別講演 17:00～18:00		特別講演 15:20～16:20		特別講演 11:00～12:00

目次

総合講演	1
企画特別講演	2
数学基礎論および歴史	5
3月15日(水)	5
3月16日(木)	8
代数学	11
3月15日(水)	11
3月16日(木)	15
3月17日(金)	17
3月18日(土)	20
幾何学	25
3月15日(水)	25
3月16日(木)	28
3月17日(金)	30
函数論	33
3月15日(水)	33
3月16日(木)	35
函数方程式論	39
3月15日(水)	39
3月16日(木)	43
3月17日(金)	46
3月18日(土)	51
実函数論	57
3月17日(金)	57
3月18日(土)	60
函数解析学	66
3月15日(水)	66
3月16日(木)	70
3月17日(金)	72
統計数学	76
3月15日(水)	76
3月16日(木)	78
3月17日(金)	79
3月18日(土)	82

応用数学	84
3月15日(水)	84
3月16日(木)	87
3月17日(金)	89
3月18日(土)	94
トポロジー	98
3月15日(水)	98
3月16日(木)	101
3月17日(金)	104
無限可積分系	107
3月17日(金)	107
3月18日(土)	108

総 合 講 演

3月16日(木) 総合講演会場

2023年度日本数学会賞春季賞受賞講演

日本数学会賞春季賞受賞者 (15:20~16:20)

Spring Prize Winner

日 比 孝 之 (阪 大*) 多面体と単項式 (16:35~17:35)

Takayuki Hibi (Osaka Univ.*) Polytopes and Monomials

概要 The theory of convex polytopes, which has its origins in Euler's theorem (1752) and Pick's formula (1899), is one of the traditional areas of combinatorics. It underwent a dramatic change in trend around 1970. Backed by the trend from abstract to concrete, a fascinating contact of convex polytopes with commutative algebra was discovered, and a bridge was built from the abstract theory of commutative algebra to the concrete theory of convex polytopes. The bridge simultaneously fostered a movement to reimport combinatorial techniques back into commutative algebra, inviting the birth of the rich soil of so-called "commutative algebra and combinatorics" from the mid-1980s, which led to a historic development of commutative algebra for monomial ideals. Later, in the 1990s, with the rapid development of computers and the evolution of software, the theory of convex polytopes took on the color of computational algebra and benefited from Gröbner bases, which is a treasure trove of techniques. This plenary talk will overview the trends in convex polytopes and commutative algebra in the last half-century since 1970 and furthermore report on current of study for reflexive polytopes and Castelnuovo polytopes.

企 画 特 別 講 演

3月15日(水)

第I会場

特別招待講演 (日本応用数学会)

吉田真紀 (情報通信研究機構) NewSpace セキュリティ —民間事業者による宇宙開発のための暗号理論— (13:00~14:00)

Maki Yoshida NewSpace security —Cryptography for space development
(Nat. Inst. Information Comm. Tech.) by private companies—

概要 In the recent NewSpace era, many small satellites have been launched on various academic or commercial missions. A low cost yet highly secure wireless communication has been desired for small satellites and small launch vehicles, for protecting their critical commands and valuable data transmissions. I would like to first explain requirements that are indispensable in such context and whether the various types of cryptography used in existing network systems satisfy them or not. I will then show our approach and results that establish high security and low implementation cost. The latest flight test conducted on July 31, 2021 achieves successful downlink transmission of 34Mbytes at a practical effective bandwidth of 512kbps.

第VI会場

横山啓太 (東北大 理) 自然数論のモデルと逆数学 (13:00~14:00)

Keita Yokoyama (Tohoku Univ.) Models of arithmetic and reverse mathematics

概要 Reverse mathematics is an ongoing program in the field of mathematical logic whose aim is to classify mathematical theorems by their logical strengths. Then, how can we say that one theorem τ is logically different from another theorem σ ? One way to show this is to construct a mathematical structure which satisfies τ (together with some basic axioms) but not σ . In this talk, I will introduce several types of structures (models) of axioms for the natural number systems (arithmetic) and show some examples of classifications of mathematical theorems from the viewpoints of model theory.

3月17日(金)

第I会場

平岡裕章 (京大 高等研) パーシステントホモロジー: 理論と応用 (13:00~14:00)

Yasuaki Hiraoka (Kyoto Univ.) Persistent homology: theory and application

概要 Topological data analysis (TDA) has emerged in this century and shed new light on data science. A particularly important tool in TDA is persistent homology, which can provide useful information about "shape of data" in a multi-scale way. Much of the development of theoretical research on persistent homology has been motivated by applications. This talk will survey the progress of persistent homology from the perspective of both mathematical and applied research.

第III会場

石橋 典 (東北大理) クラスター代数と曲面のトポロジー (13:00~14:00)
Tsukasa Ishibashi (Tohoku Univ.) Cluster algebras and the topology of surfaces

概要 Since its foundation by Fomin–Zelevinsky (2001), the study of cluster algebras has been developed with fruitful connections with various areas of mathematics, such as topology, representation theory, integrable systems, mirror symmetry, and so on. It focuses on a common combinatorial structure, called cluster transformations, which typically arises as coordinate transformations on a certain space, such as the decorated Teichmüller space of Penner (1987). Its geometric aspect is formulated as cluster varieties by Fock–Goncharov (2006). One of the motivating examples is the moduli space of decorated (twisted) G -local systems on a marked surface, which includes the decorated Teichmüller space in the case $G=SL(2)$.

In this talk, I would like to talk on the interplay between the cluster algebras and the topology of surfaces, with a focus on the geometry and cluster structure of the moduli space mentioned above. I also mention recent developments on its quantization via the quantum cluster algebra and the relation to the skein theory.

第V会場

小川 卓克 (東北大理) 端点最大正則性と臨界型非線型問題への応用 (13:00~14:00)
Takayoshi Ogawa (Tohoku Univ.) End point maximal regularity and its application to a critical nonlinear problem

概要 Maximal regularity for the abstract Cauchy problem of evolution equations is well developed in the Lebesgue-Bochner space $L^p(I; X)$, where $1 < p < \infty$, I is a time interval and X is the UMD Banach space. If X is UMD then it is necessarily reflexive and maximal regularity in non-reflexive Banach spaces is depending on each setting. In this talk, we consider the end-point case $p = 1$ in the homogeneous Besov space $X = \dot{B}_{p,1}^s(\mathbb{R}_+^n)$ in the half Euclidean space. We then apply it to the initial boundary value problem of some nonlinear evolution equations such as the Navier-Stokes equations.

3月18日(土)

第III会場

山崎 雅人 (東大IPMU) 可積分系とチャーン=サイモンズ理論 (13:00~14:00)
Masahito Yamazaki (Univ. of Tokyo) Integrability and Chern-Simons Theory

概要 Integrable models, which are "exactly solvable", lie at the intersections of many different topics/areas in mathematics and physics. However, many of the existing results are gradually built up through decades of research, and often involve trials and errors. This makes it newcomers (including myself) to learn the subject. One might therefore hope to find a new conceptual framework for understanding integrable models from scratch. In this lecture, I will discuss recent developments (in collaboration with Kevin Costello and Edward Witten, among others) in understanding integrable models from the perspective of the "four-dimensional Chern-Simons theory." This is a precise analog to Witten's explanation of knot invariants from the three-dimensional Chern-Simons theory, but is based mostly on perturbative methods of quantum field theory, which can be formulated mathematically. If time allows, I will also discuss more recent developments concerning affine Yangians (and quantum toroidal algebras) and five-dimensional version of the Chern-Simons theory. The developments covered in this lecture give rise to many new mathematical structures, many of which are yet to be fully studied in mathematics.

第VI会場

浅 芝 秀 人

(静岡大*・京大高等研・阪公大数学研)

Hideto Asashiba

(Shizuoka Univ./Kyoto Univ./Osaka Metro. Univ.)

クイバー表現のパーステンス加群への応用: 区間加群によ

る近似と分解 (13:00~14:00)

Applications of quiver representations to persistence mod-

ules: approximations and resolutions by interval modules

概要 In topological data analysis, an input datum is given as a point cloud P , which is a set of points in \mathbb{R}^d for some positive integer d . Then for each r with $0 \leq r \leq d$ an r -th persistent homology $H^r(P)$ of P over a field \mathbb{k} is calculated, which is a representation of an equi-oriented type A_n quiver Q for some positive integer n over the field \mathbb{k} . By Gabriel's theorem, the "interval" representations V_I ($I := [a, b], 1 \leq a \leq b \leq n$) of Q form a complete list of indecomposable representations of Q . For each representation M of Q , let $d_M(I)$ be the multiplicity of V_I in the indecomposable decomposition of M . Then $d_M := (d_M(I))_I$ is a complete invariant up to isomorphisms of M by the Krull-Schmidt Theorem. The family $(H^r(P))_{0 \leq r \leq d}$ contains information on P valuable in applications, which is analyzed by the persistence diagrams $d_{H^r(P)}$ of $H^r(P)$ ($0 \leq r \leq d$). When P varies with another parameter, say, time, it yields a 2D persistence module. In general, persistence modules M are understood as modules over the path algebra (or category) A of an acyclic quiver Q with full commutativity relations. (Thus A is regarded as the incidence algebra of a poset.) We deal only with the case where Q is a finite quiver. Unlike the 1-parameter case, there are an infinite number of indecomposables in the general case, which makes it difficult to use d_M in real time. The interval representation above is generalized as an interval module V_I consisting of \mathbb{k} and the identity of \mathbb{k} as a representation of Q whose support I forms a connected and convex subquiver of Q . Instead of using d_M , we use approximations of M by (direct sums of) interval modules in two senses.

数学基礎論および歴史

3月15日(水) 第IX会場

9:30~10:15

- 1 只木孝太郎 (中部大工) アルゴリズム的ランダム列が満たす実効的大数の法則とその収束速度限界
 15
 Kohtaro Tadaki (Chubu Univ.) An effectivization of the law of large numbers for algorithmically random sequences and its absolute speed limit of convergence

概要 In this talk, we show that the law of large numbers can be effectivized for an arbitrary Schnorr random infinite sequence, with respect to an arbitrary computable Bernoulli measure. Actually, we show that an absolute speed limit of convergence in this effectivization exists and it equals 2 in a certain sense.

- 2 鈴木悠大 (東北大理) On the Weihrauch degrees between C_{ω^ω} and ATR 15
 横山啓太 (東北大理)
 Yudai Suzuki (Tohoku Univ.) On the Weihrauch degrees between C_{ω^ω} and ATR
 Keita Yokoyama (Tohoku Univ.)

概要 A binary relation over the Cantor space 2^ω is called a Weihrauch problem. In this talk, we introduce ω -model reflection for Weihrauch problems and show that the ω -model reflection of ATR_2 gives an upper bound for problems provable from ATR_0 in the context of reverse mathematics. We also consider the complexity of the existence of fixed point for monotone operators (FP) and its variations (LFP) from the viewpoint of Weihrauch degrees. We will show that LFPs form a linear hierarchy between the ω -model reflection of ATR_2 and C_{ω^ω} .

10:30~11:30 特別講演

- 藤原 誠 (東京理大理) 半直観主義算術の階層と保存拡大定理
 Makoto Fujiwara (Tokyo Univ. of Sci.) Hierarchy of semi-classical arithmetic and conservative extension theorems

概要 It is well-known that classical arithmetic PA is Π_2 -conservative over intuitionistic arithmetic HA. Using a generalized negative translation, we relativize this result with respect to theories of semi-classical arithmetic, which lie in-between PA and HA. In particular, it follows from our main result that PA is Π_{k+2} -conservative over $\text{HA} + \Sigma_k\text{-LEM}$ where $\Sigma_k\text{-LEM}$ is the low-of-excluded-middle scheme for formulas of Σ_k form. In addition, we show that this conservation theorem is optimal in the sense that for any semi-classical arithmetic T, if PA is Π_{k+2} -conservative over T, then T proves $\Sigma_k\text{-LEM}$. In the same manner, we also characterize conservation theorems for other well-studied classes of formulas by fragments of classical axioms or rules. This reveals the entire structure of conservation theorems with respect to the arithmetical hierarchy of classical principles. This is a joint work with Taishi Kurahashi.

14:15~14:30 歴史部門懇談会

14:30~16:45

- 3 田村 誠 『緝古算経』の方程式について 15
 (大阪産大大学院教育機構)
 張替俊夫
 (大阪産大大学院教育機構)
 Makoto Tamura (Osaka Sangyo Univ.) On the equations of “Continuation of Ancient Mathematics”
 Toshio Harikae (Osaka Sangyo Univ.)

概要 “Continuation of Ancient Mathematics” was written by early Tang dynasty calendarist and mathematician Wang Xiaotong some time after the year 626, and is known as the oldest extant book solving cubic equations. In this talk, we list up its equations and mention the features of them.

- 4 張替俊夫 『緝古算経』の逸文の復元について 15
 (大阪産大大学院教育機構)
 田村 誠
 (大阪産大大学院教育機構)
 Toshio Harikae (Osaka Sangyo Univ.) On restruction of lost chinese text in Continuation of Ancient Mathematics
 Makoto Tamura (Osaka Sangyo Univ.)

概要 “Continuation of Ancient Mathematics” was written by early Tang dynasty calendarist and mathematician Wang Xiaotong some time after the year 626, and was listed as one of the Ten Computational Canons compiled during Tang dynasty. The aim of this talk is to provide a restruction of lost chinese text in “Continuation of Ancient Mathematics.”

- 5 脇克志 (山形大理) 和算書内図形領域抽出の精度評価 15
 Katsushi Waki (Yamagata Univ.) Evaluation of the accuracy of extracting graphic regions in a WASAN book

概要 In this presentation, we first present a workflow for extracting graphic regions in a WASAN book, which is necessary for creating a WASAN database, using the Hough transform, and evaluate the accuracy of the extraction. In the latter half of the presentation, we will show how the character “now,” which is the starting point of the graphic problem, is extracted using artificial intelligence and used to determine the graphic region for the rectangular combined region, which is a candidate for the graphic region.

- 6 小川 東 岡之只『妙矩集』における数学思想の独立性に関する問題 15
 (四日市大関孝和数学研)
 Tsukane Ogawa (Yokkaichi Univ.) Problems related to the uniqueness of mathematical thought in Oka Yukitada’s *Collection of Exquisite Formulae*

概要 I present some arguments and summarize the current state of affairs regarding the originality of the mathematical thought of Oka Tadayuki (1791-?), that is, whether his mathematical methods and ideas were unique to him or not. In particular, I suggest that his use of the term “pole” is one clue, even though no new historical documents have been discovered at present.

- 7 真島秀行 (お茶の水女大*) “Seki Takakazu, his life and bibliography” 補遺 15
 Hideyuki Majima (Ochanomizu Univ.*) Supplement to the paper “Seki Takakazu, his life and bibliography”

概要 We will talk about some episodes concerning to the discovery of Seki Takakazu’s carrer in the Kofu clan.

- 8 真島 秀行 (お茶の水女大*) 関孝和, 建部賢弘の円周率の近似計算の理論的意味づけ 15
 Hideyuki Majima (Ochanomizu Univ.*) Theoretical meaning of the calculation of Pi by Seki Takakazu and
 Takebe Katahiro

概要 We will give some theoretical meaning on the calculation of the number Pi by Seki Takakazu and Takebe Katahiro. Seki found the so-called Aitken's delta-sequenced process in his theory, but there were some errors in his calculation and he only claimed that his approximate number of Pi was a little bit less than 3.14159265359. Takebe Katahiro followed Seki's method and discovered another method of calculation of Pi.

- 9 真島 秀行 (お茶の水女大*) 2022年, 関孝和没後314年, 「綴術算経」序から300年 15
 Hideyuki Majima (Ochanomizu Univ.*) The year 2022, the memorial 314th year of Seki Takakazu and the 300th
 anniversary of "Takebe Katahiro's Tetsujjutsu-Sankei"

概要 The author talked about the works of SEKI Takakazu (?-1708), and TAKEBE Katahiro (1664-1739) on Pi. In this time, we propose to build a stone monument near his tomb in Jorinji Temple, because of his significant work concerning to Pi.

- 10 田中 紀子 (愛知県立旭丘高) 不規則な運動に対する確率的表現についての一考察 15
 松原 望 (東大*)
 Noriko Tanaka A study on stochastic representations for random motion
 (Aichi Prefectural Asahigaoka High School)
 Nozomu Matsubara (Univ. of Tokyo*)

概要 Regarding the random motion, including Brownian motion, Kiyoshi Ito et al. constructed a probabilistic representation, based on P. Levy's point of view. D. Mumford wrote "The Dawning of the Age of Stochasticity", the basic object of study in probability is the random variable and he argues that it should be treated as a basic construct, like spaces, groups and functions, and it is artificial and unnatural to define it in terms of measure theory. This time, based on D. Mumford's article, we discuss stochasticity based on its historical background.

17:00~18:00 特別講演

小林 龍彦

(前橋工大*・四日市大関孝和数学研)

Tatsuhiko Kobayashi

(Maebashi Inst. of Tech.* / Yokkaichi Univ.)

新『関孝和全集』の編集を終えて

Finishing the editing of the new Seki Takakazu's Collected Works

概要 Seki Takakazu's life and his mathematical achievements were published in the Takakazu Seki's Collected Works (Osaka Kyouiku Tosho, 1974), edited by Akira Hirayama and others. In the compilation of historical sources, it is natural to check the content of the descriptions in the documents, but issues relating to their history and the authenticity of the sources must also be carefully considered. The collected works by Akira Hirayama and others was carefully compiled based on detailed research. However, a detailed examination of the documents included in this collected works reveals that the compilation was not based on sufficient historical criticism. In June 2023, a new edition of the Seki Takakazu's Collected Works, edited by Kenji Ueno, Tsukane Ogawa, Tatsuhiko Kobayashi and Kenichi Sato will be published by Iwanami Shoten, publishers. The new collected works consists of three volumes with approximately 4000 pages. Volume 1 introduces Seki's life and his mathematical achievements based on new historical documents discovered. Volume 2 is a collection of transliteration from books and manuscripts which written in classical Chinese relating to Seki's mathematical achievements. In preparing these, the editors conducted a comprehensive survey of relevant sources scattered throughout Japan, and then collated the different editions. Volume 3 is a collection of historical documents related to Seki's research on calendrical calculations and his life. In addition, the book contains the writings of his disciples, various records and statements by contemporary mathematicians. Many aspects of Seki Takakazu's life and his biography remain unclear. For this reason, genealogical records relating to the Seki family and him have been included. In this lecture we will introduce the editorial policy of the new Seki Takakazu's Collected Works, how the authenticity of Seki's writings was judged etc.

3月16日(木) 第IX会場

9:15~12:15

- 11 薄葉季路 (早大理工) Monotonicity of ultrafilter numbers 15
Toshimichi Usuba (Waseda Univ.) Monotonicity of ultrafilter numbers

概要 We consider monotonicity of ultrafilter numbers of infinite cardinals. We show that if there are cardinals $\lambda < \kappa$ with $u(\kappa) < u(\lambda)$, then there is an inner model of a large cardinal. Hence the failure of monotonicity of ultrafilter numbers has large cardinal strength.

- 12 竹内耕太 (筑波大数理物質) On the indivisibility of metric structures 15
Kota Takeuchi (Univ. of Tsukuba) On the indivisibility of metric structures

概要 An infinite L -structure M is said to be indivisible if for any partition $M = X_0 \sqcup X_1$ there is $i < 2$ such that X_i contains an isomorphic copy of M . Many important structures, such as infinite sets, random graphs, countable dense linear order, in model theory has this property. Recently, many structures with metrics are studied by model theorists using continuous model theory. So it seems that investigating the indivisibility of metric structures are interesting from the point of view of continuous logic. In this talk we discuss about the indivisibility of Urysohn universal spaces, which was proven by Sauer.

- 13 桔 梗 宏 孝 (神戸大システム情報) Hrushovski の擬平面の構造について 15
 Hirotaka Kikyo (Kobe Univ.) On the structure of Hrushovski's pseudoplanes

概要 We discuss the structure of Hrushovski's pseudoplanes. In the definition of the Hrushovski's pseudoplanes associated to an irrational number α , good approximations k_n/d_n of α from the above are introduced. For each n , a tree B_n with k_n nodes, d_n edges and $d_n + 1$ vertices is introduced. B_n is a biminimal zero extension of the set of its leaves. Let B_∞ be the union of all B_n . Let M be the Hrushovski's pseudoplane associated to α . Let $D_n < M$ be a copy of B_n in M . Let a be a node of D_n and F_n the set of leaves of D_n . The closure of the orbit of a over F_n will be whole M . Let $D_\infty < M$ be a copy of B_∞ and let be the set of leaves of D_∞ . Let a be a node of D_∞ . We discuss the non-superstability of M by considering the type of a over some finite subset F of F_∞ and also discuss on the orbit of a over F_∞ .

- 14 坪 井 明 人 (筑 波 大*) Ramsey's theorem and coheir sequences 10
 Akito Tsuboi (Univ. of Tsukuba*) Ramsey's theorem and coheir sequences

概要 Ramsey's theorem is very easy to prove. However, its relation to the notion of coheir sequences, which is quite important in model theory, has not been discussed much. Here we present a result that illustrates the relationship between the two.

- 15 L. Pacheco (東 北 大 理) Fixed-points in epistemic logic 15
 田 中 一 之 (東 北 大 理)
 Leonardo Pacheco (Tohoku Univ.) Fixed-points in epistemic logic
 Kazuyuki Tanaka (Tohoku Univ.)

概要 The μ -calculus is an extension of modal logic by least and greatest fixed-point operators. Its alternation hierarchy characterizes formulas by the entanglement of their fixed point operators. While the alternation hierarchy is usually strict, in many settings it may collapse. We study the consequences of the strictness or the collapse in epistemic logic. We show that different axioms for knowledge imply in different numbers of possible degrees of ignorance. We also show that if we consider two or more agents simultaneously, the alternation hierarchy is strict.

- 16 倉 橋 太 志 (神戸大システム情報) Friedman–Goldfarb–Harrington の定理の拡張 15
 Taishi Kurahashi (Kobe Univ.) Extensions of the Friedman–Goldfarb–Harrington theorem

概要 We investigate the Friedman–Goldfarb–Harrington theorem. In the frameworks of classical and modal propositional logics, we study the forms of sentences whose existence is guaranteed by the FGH theorem.

- 17 倉 橋 太 志 (神戸大システム情報) 単調性を満たす証明可能性述語の様相論理 15
 小 暮 晏 佳 (金 沢 大 理 工)
 Taishi Kurahashi (Kobe Univ.) Monotonic modal logics of provability predicates
 Haruka Kogure (Kanazawa Univ.)

概要 We investigate modal logical aspects of provability predicates $\text{Pr}_T(x)$ satisfying the following condition:
M: If $T \vdash \varphi \rightarrow \psi$, then $T \vdash \text{Pr}_T(\ulcorner \varphi \urcorner) \rightarrow \text{Pr}_T(\ulcorner \psi \urcorner)$.
 We proved the arithmetical completeness theorems for monotonic modal logics MN, MN4, MNP, MNP4, and MND with respect to provability predicates satisfying the condition **M**.

- 18 岩田 莊平 (神戸大システム情報) Sacchetti の論理のカット除去可能性について 15
 Sohei Iwata (Kobe Univ.) The cut-elimination theorem for Sacchetti's logics

概要 Sacchetti showed the modal logics \mathbf{wGL}_n (Here \mathbf{wGL}_n is obtained from the provability logic \mathbf{GL} by replacing the axiom $\Box(\Box A \rightarrow A) \rightarrow \Box A$ by an weaker one $\Box(\Box^n A \rightarrow A) \rightarrow \Box A$.) have the fixed-point property. Moreover, these logics also have desirable properties (for instance, the Lyndon interpolation property, arithmetical completeness, effective constructibility of fixed-points, etc.). We investigate them in the point of view of proof theory. We introduce sequent calculi for \mathbf{wGL}_n and give a purely syntactical proof of the cut-elimination property for these calculi.

- 19 大川 裕矢 (千葉大理) 様相論理 $\mathbf{IL}^-(\mathbf{P})$ について 15
 岩田 莊平 (神戸大システム情報)
 倉橋 太志 (神戸大システム情報)
 Yuya Okawa (Chiba Univ.) The logic $\mathbf{IL}^-(\mathbf{P})$
 Sohei Iwata (Kobe Univ.)
 Taishi Kurahashi (Kobe Univ.)

概要 The logic $\mathbf{IL}^-(\mathbf{P})$ is obtained by adding the persistence principle $\mathbf{P} : A \triangleright B \rightarrow \Box(A \triangleright B)$ to the sublogic \mathbf{IL}^- of the basic interpretability logic \mathbf{IL} . We investigate $\mathbf{IL}^-(\mathbf{P})$ from several perspectives. Firstly, we show that $\mathbf{IL}^-(\mathbf{P})$ has a weak version of fixed-point property. Secondly, we show that $\mathbf{IL}^-(\mathbf{P})$ is a natural basis for some relational semantics. Thirdly, we introduce a system of sequent calculus for $\mathbf{IL}^-(\mathbf{P})$, and prove the cut-elimination theorem for it. Finally, we show that $\mathbf{IL}^-(\mathbf{P})$ is sound and complete with respect to a suitable arithmetical semantics.

12:15~12:30 数学基礎論および歴史分科会総会

- 6 後藤悠一朗 (阪大 理) 擬遺伝構造の個数に関して 15
 Yuichiro Goto (Osaka Univ.) A remark on the number of quasi-hereditary structures

概要 Dlab and Ringel showed that algebras being quasi-hereditary in all total orders for indices of primitive idempotents becomes hereditary. So, we are interested in for which orders a given quasi-hereditary algebra is again quasi-hereditary. We say that two total orders are equivalent if they induce the same set of standard modules. As a matter of fact, if an equivalence class of orders make an algebra be quasi-hereditary, then we call this class a quasi-hereditary structure. Flores, Kimura and Rognerud gave the counting method of the number of quasi-hereditary structures on path algebras of Dynkin quivers. In this talk, I give a similar result for path algebras of the quivers Q . Here Q is a quiver whose underlying graph is gotten by adding a single vertex to type A_n and a single edge between the new vertex and a vertex originally in A_n .

- 7 宮本賢伍 (茨城大 工) 多項式増大型の対称代数の τ -傾有限性について 15
 王 起 (Tsinghua Univ.)
 Kengo Miyamoto (Ibaraki Univ.) On τ -tilting finiteness of symmetric algebras of polynomial growth
 Qi Wang (Tsinghua Univ.)

概要 In this talk, we report on the τ -tilting finiteness of finite dimensional symmetric algebras of polynomial growth over an algebraically closed field. It is known that τ -tilting finiteness is preserved under Morita equivalence, but usually not preserved under derived equivalence. However, we have not found any examples of symmetric algebras which do not preserve τ -tilting finiteness under derived equivalence. In this talk, any symmetric algebras of polynomial growth preserve τ -tilting finiteness under derived equivalence.

- 8 浅井聡太 (阪大 情報) 実 Grothendieck 群の元の rigid 部分 15
 伊山 修 (東大 数理)
 Sota Asai (Osaka Univ.) The rigid parts of the elements of the real Grothendieck groups
 Osamu Iyama (Univ. of Tokyo)

概要 Let A be a finite-dimensional algebra over a field K . For any complex X in the homotopy category $K^b(\text{proj}A)$ of the complexes over the category of finitely generated projective modules, we can naturally define the rigid part of X as the maximum presilting direct summand of X . In our study, we found that the rigid part of each element θ in the real Grothendieck group $K_0(\text{proj}A)_{\mathbb{R}}$, which can be identified with an Euclidean space, is also well-defined. In this talk, I will talk about some key properties to prove our result.

- 9 佐藤真久 中山 (東屋) の補題の拡張について 10
 (愛知大地域政策・山梨大*)
 Masahisa Sato On generalized Nakayama–Azumaya’s Lemma
 (Aichi Univ./Univ. of Yamanashi*)

概要 Let R be a ring and $J(R)$ its Jacobson radical. We prove the following lemma. Generalized Nakayama–Azumaya’s Lemma “Let M be a direct summand of a direct sum of finitely generated right R -modules. If M satisfies $MJ(R) = M$, then $M = 0$.”

This unifies the following known results; “Let M be a finitely generated module or a projective module. If $MJ(R)=M$, then $M=0$.”

As an application of this lemma, we show existence of a maximal submodule for some kind of modules.

- 10 中村 力 (三重大教育) Large tilting complexes and homomorphic images of Cohen–Macaulay
M. Hrbek (Czech Acad. of Sci.) rings 15
J. Šťovíček
(Charles Univ. in Prague)
- Tsutomu Nakamura (Mie Univ.) Large tilting complexes and homomorphic images of Cohen–Macaulay
Michal Hrbek (Czech Acad. of Sci.) rings
Jan Šťovíček (Charles Univ. in Prague)

概要 Tilting theory plays a central role in representation theory of algebras. Making mutation operations possible, the notion of tilting objects was extended to silting objects. While they are usually assumed to be compact objects in triangulated categories, the definitions have been naturally extended to possibly non-compact objects. In this talk, we give a concrete way to construct (possibly non-compact) silting objects and cosilting objects in the unbounded derived category of a commutative noetherian ring R by using strictly increasing functions from $\text{Spec } R$ to the integers. Then we discuss when they are tilting objects and cotilting objects. Finally, we relate our construction of (co)tilting objects with a question when R is a homomorphic image of a Cohen–Macaulay ring.

- 11 東谷 章弘 (阪大情報) 同変 Ehrhart 理論における効果性予想とそのある種の反例 15
Akihiro Higashitani (Osaka Univ.) Effectiveness conjecture on equivariant Ehrhart theory and a kind of its
counterexample

概要 Equivariant Ehrhart theory discusses lattice polytopes admitting a finite group action ρ . Note that the case where ρ is trivial corresponds to the usual Ehrhart theory. Since every lattice polytope admits a trivial group action, we can recover the usual Ehrhart theory. In the equivariant Ehrhart theory, the effectiveness conjecture is one of the central problems. This predicts the following: for any lattice polytope P , if the H^* -series of P , which is a formal power series appearing in the equivariant Ehrhart series, becomes just a polynomial, then the H^* -series becomes effective, i.e., the coefficients really become some representations. In this talk, we give a kind of a counterexample of this conjecture.

14:15~14:50

- 12 黒田 匡迪 (日本文理大工) 単項 GAPN 関数の分類について 15
三井 健太郎 (神戸大理)
Masamichi Kuroda On the classification of monomial GAPN functions
(Nippon Bunri Univ.)
Kentaro Mitsui (Kobe Univ.)

概要 Perfect nonlinear (PN) functions and almost PN (APN) functions over finite fields of characteristic p have high nonlinearity. When $p = 2$, they have been widely studied due to their applications to cryptography and coding theory. Recently, generalized APN (GAPN) functions were defined as an algebraic generalization of APN functions in characteristic 2 to odd characteristic $p > 2$ and they have been studied. In this talk, we introduce some results of the study for the classification of monomial GAPN functions.

- 13 山形 颯 (福岡大理) r -set の分類と超平面配置が non-very generic となるための十分条件について 15

So Yamagata (Fukuoka Univ.) On classification of r -sets and a sufficient condition for non-very genericity of arrangements

概要 Manin and Schechtman introduced a family of arrangements of hyperplanes generalizing classical braid arrangements, which they called the discriminantal arrangements. Athanasiadis proved a conjecture by Bayer and Brandt providing a full description of the combinatorics of discriminantal arrangements in the case of very generic arrangements. Libgober and Settepanella described a sufficient geometric condition for given arrangements to be non very generic in terms of the notion of dependency for a certain arrangement. Settepanella and the author generalized the notion of dependency introducing r -sets and $K_{\mathbb{T}}$ -vector sets, and provided a sufficient condition for non very genericity but still not convenient to verify by hand. In this talk we give a classification of the r -sets, and a more explicit and tractable condition for non very genericity.

15:00~16:00 特別講演

藤田 直樹 (熊本大先端) シューベルト・カルキュラスと凸多面体

Naoki Fujita (Kumamoto Univ.) Schubert calculus and convex polytopes

概要 A goal of Schubert calculus is to compute the structure constants of the cohomology ring of a flag variety with respect to the basis consisting of Schubert classes. One approach to such computation is to realize Schubert classes as concrete combinatorial objects such as Schubert polynomials. In type A case, monomials in a Schubert polynomial are naturally parametrized by specific faces of a Gelfand–Tsetlin polytope. These faces are called reduced Kogan faces and appear as the limit of a semi-toric degeneration of a Schubert variety. Since the Gelfand–Tsetlin polytope is a Newton–Okounkov body of the flag variety, Newton–Okounkov bodies and induced toric degenerations are expected to give a generalization of theories in Schubert calculus to more general projective varieties. Such generalization has partially developed in the case of flag varieties in different Lie types. In this talk, some fundamental objects in Schubert calculus will be explained from the viewpoints of convex polytopes and of Kashiwara crystal bases. We also discuss their generalization to more general projective varieties.

16:15~17:15 特別講演

松井 紘樹 (徳島大社会産業理工) 三角圏のスペクトラムとスキームの復元問題

Hiroki Matsui (Tokushima Univ.) Spectra of triangulated categories and reconstruction of schemes from their derived categories

概要 For a given noetherian scheme X , it is natural to ask whether we can reconstruct X from its perfect derived category $D^{\text{pf}}(X)$ or not. This reconstruction problem has been well studied so far in algebraic geometry. It is known that we cannot reconstruct X from the triangulated category structure of $D^{\text{pf}}(X)$. For example, Mukai proved that for an abelian variety A , the perfect derived categories of A and A^{\vee} are equivalent as triangulated categories. On the other hand, Balmer proved that we can reconstruct X from its perfect derived category using its tensor triangulated category structure. For this, Balmer defined the ringed space $\text{Spec}_{\otimes}(\mathcal{T})$ for a given tensor triangulated category \mathcal{T} , which is called the *Balmer spectrum* of \mathcal{T} . Using Balmer spectra, Balmer developed *tensor triangular geometry* and as an application, he proved that the Balmer spectrum of $D^{\text{pf}}(X)$ is isomorphic to X as ringed spaces. In this sense, X can be reconstructed from the tensor triangulated category structure of $D^{\text{pf}}(X)$.

In this talk, I will introduce another ringed space $\text{Spec}_{\Delta}(\mathcal{T})$ for a given triangulated category \mathcal{T} without using tensor structure to develop a “tensor-free” analog of tensor triangular geometry. We call $\text{Spec}_{\Delta}(\mathcal{T})$ the *spectrum* of \mathcal{T} . Using the spectra of derived categories, we will prove several reconstruction results. Among them, we will give an alternative and algebraic proof of Bondal–Orlov–Ballard reconstruction theorem.

3月16日(木) 第I会場

9:10~12:00

- 14 大竹 優也 (名大多元数理) n 球加群と n 振れ自由加群の安定圏 10
 Yuya Otake (Nagoya Univ.) Stable categories of n -spherical modules and n -torsionfree modules

概要 Auslander and Bridger introduced the notions of n -spherical modules and n -torsionfree modules. In this talk, we study the stable category of n -spherical modules. Moreover, we introduce the notion of n -G-spherical modules as a G-analogue and give similar results for the stable category of n -G-spherical modules. As an application, we prove that if R is a Gorenstein local ring of Krull dimension $d > 0$, then there exists a stable equivalence between the category of $(d - 1)$ -torsionfree R -modules and the category of d -spherical modules relative to the local cohomology functor.

- 15 石塚 侖 (東工大 理) 混標数における perfectoid almost Cohen–Macaulay 代数の明示的構成 15
 下元 数馬 (日大 文理)
 Ryo Ishizuka (Tokyo Tech) An explicit construction of perfectoid almost Cohen–Macaulay algebra
 Kazuma Shimomoto (Nihon Univ.) in mixed characteristic

概要 The existence and properties of non-Noetherian algebras over a given Noetherian ring play important roles in commutative algebra. Properties of such algebras are, for example, perfectoid and almost Cohen–Macaulay. In positive characteristic, the perfect closure has these properties. In this talk, we construct an analog of the perfect closure in mixed characteristic and show that it has the same properties.

- 16 伊城 慎之介 (日大総合基礎) The canonical module of a local log-regular ring 15
 Shinnosuke Ishiro (Nihon Univ.) The canonical module of a local log-regular ring

概要 Local log-regular rings are a certain class of Cohen–Macaulay rings that are treated in logarithmic geometry. This class has similar properties to those toric rings. From this perspective, I determined the form of the canonical module of a local-log regular ring. As an application, we gave the Gorenstein criterion of this class of rings. In this talk, we introduce these results.

- 17 松下 光虹 (阪大 情報) トーリック環の conic 因子的イデアルと安定集合環への応用 15
 Koji Matsushita (Osaka Univ.) Conic divisorial ideals of toric rings and applications to stable set rings

概要 Conic divisorial ideals of toric rings play important roles in the theory of non-commutative algebraic geometry as well as commutative algebra. In this talk, we introduce an idea to determine a region representing conic classes in the divisor class group of a toric ring and a description of the conic divisorial ideals of stable set rings of perfect graphs.

- 18 石原 侑樹 (東京理大理) パラメータ付きのイデアル操作の計算 15
 Yuki Ishihara (Tokyo Univ. of Sci.) Computation of ideal operations with parameters

概要 Ideal is one of the fundamental concepts in Commutative Algebra and Algebraic Geometry. For one or more ideals, we consider ideal operations that generate new ideals from them, e.g., the ideal intersection $I \cap J$, the ideal quotient $I : J$, the saturation $I : J^\infty$, the radical \sqrt{I} , etc., where I and J are ideals in the polynomial ring $\mathbb{C}[x_1, \dots, x_n]$. In this talk, we consider several algorithms to compute ideal operations with parameters by using Comprehensive Gröbner Systems. In addition, we present an algorithm for “feasible” parameterized primary decomposition by utilizing those operations.

- 19 服部京平 (新潟大自然) Rings of nilpotent elements of monomial derivations on polynomial rings
小島秀雄 (新潟大自然) 15
Kyohei Hattori (Niigata Univ.) Rings of nilpotent elements of monomial derivations on polynomial rings
Hideo Kojima (Niigata Univ.)

概要 In this talk, we give a classification of $\text{Nil } D$ of monomial derivations D on the polynomial ring $R[x, y]$ over a UFD R containing \mathbb{Q} . Furthermore, as its application, we show that the minimal number of generators of kernels of derivations on the polynomial ring $R[x, y, z]$ as a R -algebra is unbound.

- 20 木村海渡 (名大多元数理) 加群の局所化の漸近深度 10
Kaito Kimura (Nagoya Univ.) Asymptotic depths of localizations of modules

概要 Let R be a commutative Noetherian ring, I an ideal of R , and M a finitely generated R -module. The asymptotic behavior of the quotient modules $M/I^n M$ of M is an actively studied subject in commutative algebra. In this talk, we consider the existence of an integer k such that for all integers $n \geq k$ and all prime ideals \mathfrak{p} of R , the depth of $(M/I^n M)_{\mathfrak{p}}$ is equal to the depth of $(M/I^k M)_{\mathfrak{p}}$.

- 21 遠藤直樹 (明大政治経済) On the stratification of one-dimensional Cohen–Macaulay rings 15
Naoki Endo (Meiji Univ.) On the stratification of one-dimensional Cohen–Macaulay rings

概要 The aim of this talk is to discover a good candidate for natural generalizations both of almost Gorenstein and 2-almost Gorenstein rings from the point of view of Sally modules of canonical ideals. Even though our results are at this moment restricted within the case of dimension one, we expect that a higher dimensional notion might be possible after suitable modifications.

- 22 奥間智弘 (山形大理) The core of ideals in minimally elliptic singularities 15
吉田健一 (日大文理)
渡辺敬一
 (日大文理・明大研究・知財)
Tomohiro Okuma (Yamagata Univ.) The core of ideals in minimally elliptic singularities
Ken-ichi Yoshida (Nihon Univ.)
Kei-ichi Watanabe
 (Nihon Univ./Meiji Univ.)

概要 A two-dimensional normal local domain is called a minimally elliptic singularity if it is Gorenstein and its geometric genus is one. For an m -primary integrally closed ideal I in a minimally elliptic singularity (A, m) , we represent its core using a cycle on a resolution of the singularity.

- 23 吉田健一 (日大文理) Gorensteinness for normal tangent cones of the maximal ideals in Brieskorn
奥間智弘 (山形大理) hypersurfaces 15
渡辺敬一
 (日大文理・明大研究・知財)
Ken-ichi Yoshida (Nihon Univ.) Gorensteinness for normal tangent cones of the maximal ideals in Brieskorn
Tomohiro Okuma (Yamagata Univ.) hypersurfaces
Kei-ichi Watanabe
 (Nihon Univ./Meiji Univ.)

概要 Let (A, m) be a two-dimensional excellent normal local domain containing an algebraically closed field. The graded ring defined by all integral closures of powers of an ideal I of A is called the normal tangent cone of I . We classify all Brieskorn hypersurfaces for which the normal tangent cone of the maximal ideal m is to be Gorenstein.

13:00~14:10

- 24 是枝由統(広島大理) 標数2の D_4^1 型曲面特異点のジェットスキーム上のファイバー 13
Yoshimune Koreeda (Hiroshima Univ.) Jet scheme of a singular surface of type D_4^1 in characteristic 2

概要 Let k be an algebraically closed field, X a variety over k and m a nonnegative integer. There is a space X_m over X , called the jet scheme of X of order m , parameterizing m -th jets on X . The fiber over the singular locus of X is called the singular fiber. In this talk, we assume k is of characteristic 2 and X is a singular surface of type D_4^1 . For $m \gg 0$, we give the irreducible decomposition of the singular fiber of X and determine the maximal elements among the intersections of two different irreducible components.

- 25 赤池広都(阪大理) 楕円曲面上巡回被覆ファイバー曲面の自己同型群の位数の上界について 13
Hiroto Akaike (Osaka Univ.) Bounds for the order of automorphism groups of cyclic covering fibrations of an elliptic surface

概要 In this talk, we show upper bounds for the order of automorphism groups of primitive cyclic covering fibrations of an elliptic surface. The upper bound is an explicit function in the genus of the base curve, the covering degree and the square of relative canonical divisor.

- 26 舩谷亮祐(都立大理) 平面4次曲線の弱二重接線と有理楕円曲面について 13
Ryosuke Masuya (Tokyo Metro. Univ.) On weak-bitangents for plane quartics and rational elliptic surfaces

概要 It is well-known that a smooth plane quartic has twenty-eight bitangents. We consider weak-bitangents which are generalizations on bitangents for a reduced quartic. We study weak-bitangents from the viewpoint of rational elliptic surfaces attached to reduced quartics. In particular, we consider an analogy of syzygetic triads. As a result, we can give new proofs for some classical results on certain singular plane quartics and their bitangents.

- 27 山口樹(東大数理) Big Cohen–Macaulay test ideals in equal characteristic zero via ultra-products 13
Tatsuki Yamaguchi (Univ. of Tokyo) Big Cohen–Macaulay test ideals in equal characteristic zero via ultra-products

概要 Schoutens constructed big Cohen–Macaulay (BCM) algebras in equal characteristic zero using ultra-products. In this talk, we show that BCM test ideals with respect to Schoutens' BCM algebras coincide with multiplier ideals, which define the non-klt loci of normal varieties. As an application, we study singularities of pure subrings.

3月17日(金) 第I会場

9:00~12:00

- 28 桜井真(開智学園) カイラル共形場の次元と準連接層 13
Makoto Sakurai (Kaichi Gakuen) Dimensions of chiral conformal fields and quasi-coherent sheaves

概要 The so-called AGT correspondence between 4d gauge theories and 2d conformal field theories was under discussion. However, despite much effort by mathematicians (and some physicists), the scope of results is limited, although it is sophisticated. In this report, I will deliver a trial to synthesize the mathematical literature of Beilinson and Drinfeld and some of the more “natural” physics for formulae of “anomalies.” Several conventional definitions of “dimensions” will be briefly reviewed. Then I will seek a correct notion of dimensions for chiral fields in the context of S -duality and the geometrical dimension. Some more efforts will be on the (non-)quasi-coherent sheaves for chiral algebra theories and \mathcal{D} -modules.

- 29 岩見智宏 (九工大工) Majorana double solids for three-dimensional Miyaoka–Yau type inequality with the associated third Chern classes 13

Tomohiro Iwami (Kyushu Inst. of Tech.) Majorana double solids for three-dimensional Miyaoka–Yau type inequality with the associated third Chern classes

概要 Based on S. Mori’s proof for an three-dimensional extremal neighborhood $(X, C) \subset \mathbb{C}^4$ of type A in the existence of three-dimensional flips, for an extended extremal nbd (X, C_s) with C_s not necessary irreducible nor reduced ([I2019Mar]), the author found three-dimensional Miyaoka–Yau type inequality $(MY)_{3, c_3}$ with the associated c_3 ([I2018Mar, I2019Sep]), $(MY)_{3, c_3}$ with the differential operators ([I2020Sep]) and related ones, moreover introduced an intermediate Jacobians $\widetilde{IJ}(X, C_s)$ ([I2022Mar]) and found its strong approximation for its \mathbb{Z} -structure ([I2022Sep]). In this talk, as sequel to [I2022Sep], the author will report: a) to show precise form of strong approximation for $\widetilde{IJ}(X, C_s)$ via the differential operators, b) to give slightly extended diagonal decomposition ([Bloch–Srinivas 1983]) to introduce real/Majorana spinor by a), and c) to construct \mathbb{Z} -structure by gluing this underlying real structure via b) regarding to integral Hodge conjecture, in order to discuss alternative to [Artin–Mumford 1972].

- 30 高橋優太 (中大理工) 正標数におけるネフ接束をもつ4次元ファノ多様体の分類 13
渡邊 究 (中大理工)

Yuta Takahashi (Chuo Univ.) Fano 4-folds with nef tangent bundle in positive characteristic
Kiwamu Watanabe (Chuo Univ.)

概要 In characteristic 0, the Campana–Petersen conjecture claims that the only smooth Fano variety with nef tangent bundle should be homogeneous. In this talk, we present the positive characteristic version of the Campana–Petersen conjecture. In particular, we give a result for Fano 4-folds with nef tangent bundle and Picard number greater than one.

- 31 吉野太郎 (東大数理) On the degree of irrationality of complete intersections 13
Taro Yoshino (Univ. of Tokyo) On the degree of irrationality of complete intersections

概要 This is joint work with Lucas Braune. We obtain a lower bound of the degree of irrationality of very general complete intersections over the complex field from the recent results of Braune and Chen–Stapleton. For combining these results, we make a minor adjustment of Chen–Stapleton’s method using the trace map of differential modules.

- 32 水野雄貴 (早大理工) Some examples of noncommutative projective Calabi–Yau schemes ... 13
Yuki Mizuno (Waseda Univ.) Some examples of noncommutative projective Calabi–Yau schemes

概要 We construct some examples of noncommutative projective Calabi–Yau schemes by using noncommutative Segre products and quantum weighted hypersurfaces. We also compare them with commutative Calabi–Yau varieties and examples by Kanazawa.

- 33 星明考 (新潟大理) Birational classification for algebraic tori (I) 13
山崎愛一 (京大理)

Akinari Hoshi (Niigata Univ.) Birational classification for algebraic tori (I)
Aiichi Yamasaki (Kyoto Univ.)

概要 We give a stably birational classification for algebraic k -tori of dimensions 3 and 4 over a field k .

- 34 星明考 (新潟大理) Birational classification for algebraic tori (II) 13
山崎愛一 (京大理)

Akinari Hoshi (Niigata Univ.) Birational classification for algebraic tori (II)
Aiichi Yamasaki (Kyoto Univ.)

概要 We give a stably birational classification for algebraic k -tori of dimensions 3 and 4 over a field k .

- 35 助 永 真 之 (広島大先進理工) トロピカル平面曲線の交わりの実現問題 13
 Masayuki Sukenaga (Hiroshima Univ.) Tropical lifting problem for the intersection of plane curves

概要 Let D be a tropical divisor in the intersection of two tropical plane curves. We study when it can be realized as the tropicalization of the intersection of two algebraic curves. When a graph theoretic condition determined by these tropical curves is satisfied, we can algorithmically find algebraic curves such that the tropicalization of their intersection is D .

- 36 天 野 通 大 (明 星 大) Witt vectors のある準同型写像の核について II 10
 Michio Amano (Meisei Univ.) On kernels of certain homomorphisms of Witt vectors, II

概要 Let $W(A)$ be a Witt group over A and let $W^n(A)$ be the product of n of $W(A)$. T. Sekiguchi and N. Suwa have constructed a certain endomorphism U_n of $W^n(A)$ related to the deformation group scheme from the additive group scheme to the multiplicative group scheme. We study the equality of kernels of certain homomorphisms related to U_n .

- 37 川 辺 大 貴 (東 北 大 理) Chow motives of genus one fibrations 13
 Daiki Kawabe (Tohoku Univ.) Chow motives of genus one fibrations

概要 Genus 1 fibrations play a central role in the classification of surfaces. Let $f : X \rightarrow C$ be a genus 1 fibration from a surface to a curve. Then there is a Jacobian genus 1 fibration $j : J \rightarrow C$ that satisfies some good properties. In this talk, we prove that the motive of X is isomorphic to the motive of J . As an application, we prove motivic finite-dimensionality for surfaces not of general type with geometric genus 0. This can be regarded as a generalization of Bloch–Kas–Lieberman’s result to arbitrary characteristic.

- 38 安 藤 哲 哉 (千 葉 大 理) 4 変数 4 次対称 extremal 不等式と K3 曲面 13
 Tetsuya Ando (Chiba Univ.) Symmetric quartic extremal inequalities of four variables and K3 surfaces

概要 Let $P_{4,4}$ be the convex cone of quartic homogeneous polynomials of four variables which is positive semidefinite. We present a family of symmetric elements of $P_{4,4}$. Let f be a such polynomial. Then zero locus of f in the complex 3-dimensional projective space is a K3 surface which has 10 real A_1 type rational double points.

- 39 寄 崎 恵 美 子 (都 立 大 理) Ramified and Split models of rational elliptic surfaces and bitangent
 坂 内 真 三 (岡 山 理 大 理) lines for a quartic curve 10
 徳 永 浩 雄 (都 立 大 理)

Emiko Yorisaki (Tokyo Metro. Univ.) Ramified and Split models of rational elliptic surfaces and bitangent
 Shinzo Bannai (Okayama Univ. of Sci.) lines for a quartic curve
 Hiro-o Tokunaga (Tokyo Metro. Univ.)

概要 We introduce the notion of ramified and split models of elliptic surfaces. We consider how these two models are related. In particular, we apply our consideration to rational elliptic surfaces which arise as smooth models of double covers of the projective plane branched along quartics. As a result, we determine the number of concurrent bitangent lines for irreducible quartics with at most nodes.

14:15~14:30 2023年度(第26回)日本数学会代数学賞授与式

14:40~15:40 2023年度(第26回)日本数学会代数学賞受賞特別講演

権業善範(東大数理) 高次元極小モデル理論の構築とその応用について

Yoshinori Gongyo (Univ. of Tokyo) On higher dimensional minimal model theory and its applications

概要 I will report what I am going on projects related to the minimal model program. First, I will discuss some new invariants for studies of Fano varieties. We call it the “total index”, which is natural invariant mixing Shokurov’s complexities and the usual Fano index. In this study, the Birkar generalized pair naturally appears. Using this invariant, I propose some conjecture for the characterization of the product of projective spaces, which is some variant of Mukai’s type conjecture. The methods are to use the local-global method from singularities rather than studies of rational curves. To study this conjecture, the non-vanishing conjecture and characterization of toric varieties are involved. Second, I will explain what I consider recently on the mainstream of the minimal model theory.

16:00~17:00 2023年度(第26回)日本数学会代数学賞受賞特別講演

若槻 聡(金沢大理工) 跡公式とその応用

Satoshi Wakatsuki (Kanazawa Univ.) Trace formulas and their applications

概要 The trace formula is one of the main tools in the study of automorphic forms, and its theory has been greatly developed by many researchers (Selberg, Arthur, etc.). The trace formula plays an important role even today, such as the classification of automorphic representations by stable trace formulas and the study of automorphic periods by relative trace formulas. In this talk, I will give an overview of trace formulas and their applications. After that, I will introduce my recent research on asymptotic behaviors for dimensions of spaces of automorphic forms.

3月18日(土) 第I会場

9:15~12:00

40 飯高 茂(学習院大*) 第2種スーパー完全数と完全数 80 13
Shigeru Iitaka (Gakushuin Univ.*) Super perfect numbers of the second kind

概要 Let m denote translation parameter, and let h denote an odd prime.

Defining A to be $\sigma(a) + m$ if $h\sigma(A) = 2a(h+1) - h^2 + mh$ then a is said to be super perfect numbers of the second kind. A is called a partner of a .

41 小林 雅人(神奈川大工) 約数和関数のバランス 13
Masato Kobayashi (Kanagawa Univ.) Balance for the divisor sum function

概要 Let us say finite $A \subseteq \mathbb{N}$ is balanced if

$$\sum_{n \in A} \sum_{d|n, d \text{ odd}} d = \sum_{n \in A} \sum_{d|n, d \text{ even}} d.$$

For example, $\{1, 2\}, \{1, 3, 4\}, \{3, 6, 8, 9\}$ are balanced. For each natural number N , does there exist a balanced set A of the cardinality N ? I solved this problem affirmatively in June 2022 with Ramanujan’s theta function and Jacobi’s triple product identity. In this talk, I wish to show details of this.

- 42 金堂 優哉 (名大多元数理) The relation between a generalized Fibonacci sequence and the length of Cunningham chains 13

Yuya Kanado (Nagoya Univ.) The relation between a generalized Fibonacci sequence and the length of Cunningham chains

概要 Let p be a prime number and $\alpha \geq 3$ be an integer. In this talk, a generalized Fibonacci sequence $\mathcal{F}_\alpha = (F_n)_{n \in \mathbb{Z}}$ is defined as $F_0 = 0, F_1 = 1, F_{n+2} = \alpha F_{n+1} + F_n$ ($n \in \mathbb{Z}$), and $\mathcal{F}_\alpha \sigma(n) = \sum_{d|n, 0 < d \in \mathcal{F}_\alpha} d$ is called the divisor function on \mathcal{F}_α . Then we obtain that for two prime numbers p and q the fact $p = 2q + 1$ or $2q - 1$ is equivalent to $\mathcal{F}_\alpha \sigma(\mathcal{F}_\alpha \sigma(F_p)) = \mathcal{F}_\alpha \sigma(F_q)$ for some α .

- 43 栗野 俊一 (日大経済) The error term in the prime number theorem under the RH 10
中野 正俊 (気仙沼高等技術専)

Shun-ichi Kurino (Nihon Univ.) The error term in the prime number theorem under the RH
Masatoshi Nakano
(Kesennuma Coll. of Tech.)

概要 We propose the conjecture that the error term in the prime number theorem under the RH is $o(\sqrt{x})$.

- 44 杉山 真吾 (日大理工) Dirichlet L 関数の族の重み付き零点密度について 10
アデルマスリアジャヤ
(九大数理)

Shingo Sugiyama (Nihon Univ.) On a weighted density of zeros of Dirichlet L -functions in a family
Ade Irma Suriajaya (Kyushu Univ.)

概要 Katz and Sarnak suggested a relation between zeros of L -functions in a family and eigenvalues of random matrices in a compact matrix group. In this talk, we consider the analogous conjecture weighted by L -values and verify this conjecture in the case of the family of Dirichlet L -functions attached to nontrivial Dirichlet characters of a prime modulus. This is a joint work with Ade Irma Suriajaya (Kyushu University).

- 45 橋本 康史 (琉球大理) セルバークゼータ関数に関する普遍性定理 13

Yasufumi Hashimoto Universality theorem for the Selberg zeta function
(Univ. of Ryukyus)

概要 In 2013, Drungilas et al. proposed the universality theorem for the Selberg zeta function associated with the modular group, which is (probably) the first universality theorem for the zeta function of the order of more than one. Recently in 2021, Mishou extended it by proving the joint universality theorem for the tuple of Selberg zeta functions associated with principal congruence subgroups. In this talk, we present the universality theorem (and joint universality theorem) for the Selberg zeta function associated with (not necessarily congruence) subgroup of the modular group. We also note that our universality theorems are available in wider regions than the regions given in the previous works.

- 46 篠原 健 (名大多元数理) On shuffle product of desingularized multiple zeta functions 13

Takeshi Shinohara (Nagoya Univ.) On shuffle product of desingularized multiple zeta functions

概要 Furusho, Komori, Matsumoto, and Tsumura introduced the desingularized multiple zeta functions to resolve all singularities of multiple zeta functions (MZF for short). It is a natural question to ask how much desingularized MZFs retain the properties of MZFs. As one of the answers, we show that special values of desingularized MZFs at positive integer points satisfy the shuffle product that holds among special values of MZFs at positive integer points.

- 47 加藤 正輝 (富山 高専) ルート系のゼータ関数の q -類似について 13
 Masaki Kato On q -analogues of zeta functions of root systems
 (Toyama Nat. Coll. of Tech.)

概要 Komori, Matsumoto and Tsumura introduced zeta functions $\zeta_r(\mathbf{s}, \Delta)$ associated with root systems Δ . In this talk, we introduce q -analogues of functions $\zeta_r(\mathbf{s}, \Delta)$, denoted by $\zeta_r(\mathbf{s}, \mathbf{a}, \Delta; q)$. We establish q -extensions of the following properties satisfied by $\zeta_r(\mathbf{s}, \Delta)$: (a) Integral representations of sums of zeta functions which have Weyl group symmetries. (b) Explicit expressions of Witten volume formulas. (c) Functional relations including Witten volume formulas. We also introduce elliptic analogues of $\zeta_r(\mathbf{s}, \mathbf{a}, \Delta; q)$, denoted by $\zeta_r(\mathbf{s}, \mathbf{a}, \beta, \Delta; p, q)$, and investigate their basis properties. To extend the properties (a) and (b) to $\zeta_r(\mathbf{s}, \mathbf{a}, \beta, \Delta; p, q)$, the factorization formulas of Poincaré polynomials, established by Macdonald, plays important roles.

- 48 峰 正博 (上智大理工) 2次無理数をパラメータに持つ Hurwitz ゼータ関数に対する弱い稠密性定理 13
 Masahiro Mine (Sophia Univ.) Weak denseness theorem for the Hurwitz zeta-function with quadratic irrational parameter

概要 In this talk, we consider the value-distribution of the Hurwitz zeta-function with algebraic irrational parameter. The main result is that any complex number can be approximated by values of the Hurwitz zeta-function for arbitrary quadratic irrational parameters but with finite exceptions.

- 49 中井 啓太 (名大多元数理) 対数をとった Selberg クラスの L 関数の反復積分の普遍性定理 10
 Keita Nakai (Nagoya Univ.) Universality for the iterated integrals for the logarithm of L-functions in the Selberg class

概要 In 1975, Voronin proved the Universality theorem for the Riemann zeta-function and in 2022, Endo generalized this theorem for the iterated integrals for the logarithm of the Riemann zeta-function. In this talk, we give a more generalization of this theorem for the iterated integrals for the logarithm of L-functions in the Selberg class.

- 50 杉山 和成 (千葉工大) Siegel ゼータ関数の保型性について 13
 Kazunari Sugiyama The modularity of Siegel's zeta functions
 (Chiba Inst. of Tech.)

概要 Siegel defined zeta functions associated with indefinite quadratic forms, and proved their analytic properties such as analytic continuations and functional equations. We discuss the modularity of Siegel's zeta functions on the basis of converse theorems.

14:15~16:45

- 51 武田 渉 (東京理大理) Quadratic relations for 9th variation Schur functions involving Plücker relations 13
 中筋 麻貴 (上智大理工・東北大理)
 山崎 義徳 (愛媛大理)
 Wataru Takeda (Tokyo Univ. of Sci.) Quadratic relations for 9th variation Schur functions involving Plücker relations
 Maki Nakasuji
 (Sophia Univ./Tohoku Univ.)
 Yoshinori Yamasaki (Ehime Univ.)

概要 The 9th variation of Schur polynomial is known to satisfy determinant formulas such as Jacobi–Trudi, Gambelli, and Hamel–Goulden identities. In this talk, we apply the Plücker relations to these identities. These relations are known to give algebraic dependencies among the determinants of the various minors of an arbitrary matrix. Applying them to the $n \times n$ minors of an $n \times 2n$ matrix, we give some quadratic relations between the 9th variation of Schur polynomials.

- 52 渋川元樹 (神戸大理) Some monotonic properties of special values of the bivariate complete
後藤良彰 (小樽商大) homogeneous symmetric polynomials 10
Genki Shibukawa (Kobe Univ.) Some monotonic properties of special values of the bivariate complete
Yoshiaki Goto homogeneous symmetric polynomials
(Otaru Univ. of Commerce)

概要 We prove some monotonic properties of special values of the bivariate complete homogeneous symmetric polynomials.

- 53 星明考 (新潟大理) Hasse norm principle for M_{11} extensions 13
金井和貴 (新潟大自然)
山崎愛一 (京大理)
Akinari Hoshi (Niigata Univ.) Hasse norm principle for M_{11} extensions
Kazuki Kanai (Niigata Univ.)
Aiichi Yamasaki (Kyoto Univ.)

概要 Let k be a field and T be an algebraic k -torus. In 1969, over a global field k , Voskresenskii proved that there exists an exact sequence $0 \rightarrow A(T) \rightarrow H^1(k, \text{Pic } \bar{X})^\vee \rightarrow Sha(T) \rightarrow 0$ where $A(T)$ is the kernel of the weak approximation of T , $Sha(T)$ is the Shafarevich–Tate group of T , X is a smooth k -compactification of T , $\text{Pic } \bar{X}$ is the Picard group of $\bar{X} = X \times_k \bar{k}$. On the other hand, in 1963, Ono proved that for the norm one torus $T = R_{K/k}^{(1)}(G_m)$ of K/k , $Sha(T) = 0$ if and only if the Hasse norm principle holds for K/k . We determine $H^1(k, \text{Pic } \bar{X})$ for norm one tori $T = R_{K/k}^{(1)}(G_m)$ when the Galois group $\text{Gal}(L/k)$ of the Galois closure L/k of K/k is isomorphic to the Mathieu group M_{11} of degree 11. We also give a necessary and sufficient condition for the Hasse norm principle for such extensions K/k with $\text{Gal}(L/k) \simeq M_{11}$.

- 54 伊吹山知義 (阪大*) 符号付きパラモジュラー形式の次元公式 10
Tomoyoshi Ibukiyama (Osaka Univ.*) Dimensions of paramodular forms with involutions

概要 Paramodular forms are a part of Siegel modular forms. For degree two case, they are important for Shimura–Taniyama conjecture for abelian surfaces and the new form theory of Roberts and Schmidt. Here we give explicit dimension formulas for paramodular forms of any vector valued weight $\det^3 \text{Sym}(j)$ with $k \geq 3$ of prime level p with fixed sign of the Atkin–Lehner involution. It is known that such forms have a good correspondence with certain algebraic modular forms up to endoscopic part and old forms. We calculate dimensions of the corresponding algebraic modular forms and apply it to paramodular case.

- 55 伊吹山知義 (阪大*) ジーゲル保型形式上の微分作用素とラプラス変換 13
Tomoyoshi Ibukiyama (Osaka Univ.*) Differential operators on Siegel modular forms and Laplace transforms

概要 The talk has two aims. One is to give a new special basis of the polynomial ring $\mathbb{C}[T]$ in components of $n \times n$ symmetric matrix T . The basis $P(T)$ is characterized by the action of $P(\partial_Z)$ on $\det(CZ + D)^s$, where ∂_Z is the matrix of partial derivatives of the variable of the Siegel upper half space and $\det(CZ + D)$ is the automorphy factor. Everything is described explicitly. For example, the basis is obtained as coefficients of certain explicitly described generating series. The second aim is to apply this to the so-called pullback formula of the Siegel Eisenstein series of degree $2n$ and determine all appearing constants.

- 56 境 優一 (久留米工大基幹教育センター) On modular-form solutions of a certain modular linear differential equations for cocompact groups 13
 Yuichi Sakai (Kurume Inst. of Tech.) On modular-form solutions of a certain modular linear differential equations for cocompact groups

概要 In this talk, we give a modular linear differential equation of second order for a cocompact group of discriminant 6, and we also give its solutions. Furthermore we prove a relation between supersingular polynomials and certain polynomials derived from solutions of the modular linear differential equation.

- 57 杉山真吾 (日大理工) Hilbert 保型形式と Siegel 保型形式の Hecke 固有値の代数的整数性 10
 佐久川憲児 (信州大教育) Shingo Sugiyama (Nihon Univ.) Integrality of Hecke eigenvalues of Hilbert and Siegel modular forms
 Kenji Sakugawa (Shinshu Univ.)

概要 In this talk, we prove the integrality of Hecke eigenvalues of Hilbert modular forms and Siegel modular forms in a general setting. As applications, we estimate the degrees of Hecke fields of cuspidal automorphic representations of $GL(2d)$ with a prime number d and $Sp(2n)$.

- 58 加藤裕基 (宇部工高専) Almost mathematics and its algebraic K -theory from non-unital algebras viewpoint 13
 Yuki Kato (Ube Nat. Coll. of Tech.) Almost mathematics and its algebraic K -theory from non-unital algebras viewpoint

概要 While almost mathematics has various applications to arithmetic geometry, Quillen mentioned linear algebra over non-unital rings which is the same as almost mathematics introduced by Faltings; and Gabber and Ramero. Quillen's work is more conceptual in the sense of using categorical language: in his work, almost mathematics is characterized as bilocalization of an abelian category of modules. By using the theory of bilocalization, we define the derived category (or stable ∞ -category) of almost perfect complexes, enabling us to obtain the K -theory of almost modules. Furthermore, by modifying K -theory of almost algebra, we prove that the modified K -groups decompose into the almost algebraic K -theory and the almost acyclic part of them. The main result of almost K -theory is that the almost K -theory of A is represented by the modified K -theory of the corresponding the idempotent non-unital algebra.

- 59 南 範彦 (名工大) 任意のモチビック一般コホモロジー論に対する SBNR ((stably birationalized) unramified sheaf) と Gersten 型分解 13
 Norihiko Minami SBNR (stably birationalized unramified sheaf) and Gersten type resolution for any motivic generalized cohomology theory
 (Nagoya Inst. of Tech.)

概要 Suppose the base field is perfect. Then, for any generalized motivic cohomology theory, I shall discuss its Zariski sheafification which turns out to be unramified in Morel's sense and a Nisnevich sheaf in this particular case.

Amongst of all, I shall emphasize the existence of its Gersten type resolution as Zariski sheaves and its SBNR - stably birationalized Nisnevich subsheaf of the unramified sheaf.

- 60 南 範彦 (名工大) Integral Hodge 予想反例の高次元非有理性 13
 Norihiko Minami Higher codimensional irrationality of counter examples of the integral Hodge conjecture
 (Nagoya Inst. of Tech.)

概要 Some necessary condition for a non-singular complex projective manifold X to be a counter-example for the codimension k integral Hodge conjecture $Z^{2k}(X) \neq 0$ is given.

Our necessary condition is stated as codimension c_k irrationality for some c_k , which grows as k increases. This may suggest that counter-examples to the integral Hodge conjecture become ubiquitous as k increases.

幾何学

3月15日(水) 第II会場

9:30~11:40

- 1 キヨヴスキアントニ (沖縄科学技術大) Asymptotically mean value harmonic functions 15
 Antoni Kijowski (Okinawa Inst. of Sci. and Tech. Grad. Univ.) Asymptotically mean value harmonic functions

概要 The Blaschke–Privaloff–Zaremba theorem gives rise to a recently investigated concept of the asymptotic mean value harmonic function on a metric measure space X . For a function $u \in L^1_{loc}(X)$ define the r -laplacian of u as:

$$\Delta_r u(x) := \frac{u_{B_r(x)} - u(x)}{r^2},$$

where $u_{B_r(x)}$ stands for the mean-value of u over a ball $B_r(x)$. We say that u is amv-harmonic if its r -laplacian converges to 0 in an appropriate sense as $r \rightarrow 0$. In my talk I will present known results focusing on regularity, blow ups and characterization of these functions in the following settings: weighted Euclidean domains, Carnot groups and RCD spaces.

- 2 横田 滋 亮 (東北大理) Geometric data set の幾何学 15
 Shigeaki Yokota (Tohoku Univ.) Geometry of geometric data set

概要 Gromov originated a geometric theory of metric measure spaces based on the phenomenon of concentration of measure and the theory of the collapsing of Riemannian manifolds; Pestov pointed out the connection of Gromov’s theory with the curse of dimensionality in statistics. To improve time complexity, Hanika, Schneider, Stumme proposed to generalize metric measure space to ‘geometric data set’. The total set of the metric measure spaces has two natural distance functions, the observable and box distance functions, and their topological properties are well known, but the properties of the total set of the geometric data set are almost unknown. In this talk, we extend these results of metric measure space to the total set of geometric data sets.

- 3 数川 大 輔 (九大数理) 測度距離空間全体の位相的性質について 15
 中島 啓 貴 (東北大IEHE)
 塩谷 隆 (東北大理)
 Daisuke Kazukawa (Kyushu Univ.) Topological aspects of the space of metric measure spaces
 Hiroki Nakajima (Tohoku Univ.)
 Takashi Shioya (Tohoku Univ.)

概要 Gromov introduced two distance functions, the box distance and the observable distance, on the space of isomorphism classes of metric measure spaces and developed the convergence theory of metric measure spaces. We investigate several topological properties on the space equipped with these distance functions toward a deep understanding of convergence theory.

- 4 数川大輔 (九大数理) 測度距離空間全体のスケーリング作用に関する主束構造について 10
 中島啓貴 (東北大IEHE)
 塩谷隆 (東大理)

Daisuke Kazukawa (Kyushu Univ.) Principal bundle structure of the space of metric measure spaces
 Hiroki Nakajima (Tohoku Univ.)
 Takashi Shioya (Tohoku Univ.)

概要 We study the topological structure of the space \mathcal{X} of isomorphism classes of metric measure spaces equipped with the box or concentration topologies. We consider the scale-change action of the multiplicative group \mathbb{R}_+ of positive real numbers on \mathcal{X} , which has a one-point metric measure space, say $*$, as only one fixed point. We prove that the \mathbb{R}_+ -action on $\mathcal{X} \setminus \{*\}$ admits the structure of nontrivial and locally trivial principal \mathbb{R}_+ -bundle over the quotient space. A similar statement is obtained for the pyramidal compactification of \mathcal{X} , where we completely determine the structure of the fixed-point set of the \mathbb{R}_+ -action on the compactification.

- 5 松家拓稔 (都立大理) 距離空間の自由積と粗バウム・コンヌ予想 15
 Takumi Matsuka (Tokyo Metro. Univ.) Free products of metric spaces and the coarse Baum–Connes conjecture

概要 We introduce the notion of free products of metric spaces. When X and Y are coarsely convex, the free product $X * Y$ is coarsely convex. In particular, the free product $X * Y$ satisfies the coarse Baum–Connes conjecture. This talk is based on an ongoing joint work with Tomohiro Fukaya (Tokyo Metropolitan University).

- 6 伊敷喜斗 (理化学研RAP) グロモフ・ハウスドルフ空間の中の連結空間 15
 Yoshito Ishiki (RIKEN) Continua in the Gromov–Hausdorff space

概要 We first prove that for all compact metrizable spaces, there exists a topological embedding of the compact metrizable space into each of the sets of compact metric spaces which are connected, path-connected, geodesic, or CAT(0), in the Gromov–Hausdorff space with finite prescribed values. As its application, we show that the sets prescribed above are path-connected and their non-empty open subsets have infinite topological dimension. By the same method, we also prove that the set of all proper CAT(0) spaces is path-connected and its non-empty open subsets have infinite topological dimension with respect to the pointed Gromov–Hausdorff distance.

- 7 伊敷喜斗 (理化学研RAP) グロモフ・ハウスドルフ空間の中の木 15
 Yoshito Ishiki (RIKEN) Metric trees in the Gromov–Hausdorff space

概要 Using the wedge sum of metric spaces, for all compact metrizable spaces, we construct a topological embedding of the compact metrizable space into the set of all metric trees in the Gromov–Hausdorff space with finite prescribed values. As its application, we show that the set of all metric trees is path-connected and its all non-empty open subsets have infinite topological dimension.

14:15~16:00

- 8 米村拳太郎 (九大数理) 多様体構造を持つカンドルの Lie 群への埋め込み 10
 Kentaro Yonemura (Kyushu Univ.) Embedding smooth quandles in Lie groups

概要 We deal with embedding smooth quandles in Lie groups in this talk.

- 9 佐々木 優 (東京工高専) 結合的グラスマン多様体の部分多様体 15
 Yuuki Sasaki (Tokyo Nat. Coll. of Tech.) Some submanifolds of the associative Grassmann manifold

概要 We construct a totally complex immersion from almost complex submanifolds of the 6-dimensional sphere into the associative Grassmann manifold. Also, as an analogy of almost complex immersions we construct a CR immersion from 3-dimensional CR submanifolds of the 6-dimensional sphere into the associative Grassmann manifold. Moreover, in the associative Grassmann manifold we show that many orbits of the action of the isotropy subgroup of the isometry group are the image of some CR immersions.

- 10 木村 太郎 (鶴岡工高専) Biharmonic Cartan embeddings 10
 間下 克哉 (法政大理工)
 Taro Kimura Biharmonic Cartan embeddings
 (Nat. Inst. of Tech., Tsuruoka Coll.)
 Katsuya Mashimo (Hosei Univ.)

概要 In this talk, we determine all the proper biharmonic submanifolds in compact Lie groups which are the image of Cartan embeddings defined by automorphism of order 4. Moreover we show that if the Cartan embedding defined by automorphism of order 3 is biharmonic, then it is harmonic.

- 11 井川 治 (京都工繊大工芸) $Spin(8)$ の triality automorphism から誘導される σ 作用の軌道の幾何学
 間下 克哉 (法政大理工) 15
 Osamu Ikawa (Kyoto Inst. Tech.) The geometry of orbits of σ -action induced by triality automorphism of
 Katsuya Mashimo (Hosei Univ.) $Spin(8)$

概要 The universal covering group $Spin(8)$ of the special orthogonal group $SO(8)$ has an outer automorphism σ of order 3 called the triality automorphism. From this σ , the action of $Spin(8)$ on $Spin(8)$ itself, called the σ -action, is determined. This action is a hyperpolar action of cohomogeneity (= codimension of maximum-dimensional orbit) 2. We report on the orbit space and properties of orbits of this action.

- 12 塚田 和美 (お茶の水女大*) Lie 球面幾何学の複素化と実グラスマン多様体の全複素部分多様体 15
 Kazumi Tsukada (Ochanomizu Univ.*) The complexification of Lie sphere geometry and totally complex submanifolds of the real Grassmann manifold

概要 The real Grassmann manifold $Gr_4(R^n)$ of all 4-dimensional subspaces in R^n is known to have the quaternionic Kähler structure. We consider the set $H_2(C^n)$ of all 2-dimensional complex subspaces in C^n that are isotropic with respect to the standard complex inner product. $H_2(C^n)$ is a complex manifold of dimension $2n - 7$ and has the holomorphic contact structure. The natural projection of $H_2(C^n)$ onto $Gr_4(R^n)$ is a twistor fibration. Through $H_2(C^n)$, we show the relation between the geometry regarded as the complexification of Lie sphere geometry and that of totally complex submanifolds of $Gr_4(R^n)$.

- 13 坂根 由昌 (阪大*) コンパクト等質空間上のアインシュタイン計量の存在・非存在について 15
 Yusuke Sakane (Osaka Univ.*) Existence and non-existence of Einstein metrics on compact homogeneous manifolds

概要 The study of existence and non-existence of Einstein metrics on G/H is started by Wang and Ziller in 1986. More general results for non-existence are obtained by Böhm in 2005. One of his results is that $G/H = SU(n + k_1 + \cdots + k_p)/S(SO(n)U(1) \times U(k_1) \times \cdots \times U(k_p))$ (where $SO(n)U(1) \subset U(n)$) does not admit G -invariant Einstein metrics, if $n > (k_1 + \cdots + k_p)^2 + 2$. He has obtained the results by considering “traceless” part of Ricci tensor. For $G/H = SO(2n + k_1 + \cdots + k_p)/SO(n)U(1) \times SO(k_1) \times \cdots \times SO(k_p)$ (where $SO(n)U(1) \subset U(n) \subset SO(2n)$), we can not apply his method directly. In this talk, we will show that, if $n \geq 5(k_1 + \cdots + k_p)$ ($k_i \geq 3$), G/H does not admit G -invariant Einstein metrics. We also discuss existence of Einstein metrics for compact homogeneous manifolds G/H above.

16:15~17:15 特別講演

多羅間大輔 (立命館大理工) 半単純 Lie 群上の可積分測地流について

Daisuke Tarama (Ritsumeikan Univ.) On integrable geodesic flows over a semi-simple Lie group

概要 This talk deals with the geodesic flow of a semi-simple Lie group equipped with a left-invariant metric. After a short review over the geometric settings for general left-invariant Hamiltonian systems on the cotangent bundle to a Lie group, a class of left-invariant metrics is considered on a semi-simple Lie group, following Mishchenko and Fomenko. These metrics have been known to give rise to completely integrable geodesic flows since late 1970's. It is however rather recent that their equilibrium points are studied mostly on the basis of techniques in symplectic/Poisson geometry. In the talk, these developments are briefly reviewed. Then, as a main result, the classification of Williamson types of relative equilibrium points for a large subclass of above integrable geodesic flows is explained in terms of root systems. Related studies and remaining problems are also mentioned.

3月16日(木) 第II会場

9:30~12:0014 川村昌也 (香川高専) コンパクト概 Hermitian 多様体上の k -th Gauduchon 計量について 15

Masaya Kawamura

On a k -th Gauduchon metric on compact almost Hermitian manifolds

(Kagawa Nat. Coll. of Tech.)

概要 In this talk, we introduce some results about k -th Gauduchon metrics on compact almost Hermitian manifolds. We have shown that if both the conformally k -th Gauduchon condition and the conformally semi-Kähler condition are satisfied, then it becomes conformally quasi-Kähler. Also, we characterize the k -th Gauduchon condition and by applying its characterization, we have shown some geometric conditions for negative Kodaira dimension on compact almost Hermitian manifolds.

15 宮武夏雄 (九大IMI) Uniformization of compact Sasakian manifolds using basic Higgs bundles 15

糟谷久矢 (阪大理)

Natsuo Miyatake (Kyushu Univ.) Uniformization of compact Sasakian manifolds using basic Higgs bundles

Hisashi Kasuya (Osaka Univ.)

概要 Simpson introduced the notion of uniformizing variation of Hodge structure (VHS) and showed that the universal covering of a compact Kähler manifold is isomorphic to a bounded symmetric domain if and only if there exists a uniformizing VHS on the Kähler manifold. In this talk, on compact Sasakian manifolds, we consider an analogy of this. We can formulate an analogy of the uniformizing VHS also on compact Sasakian manifolds. However, even if there exists a uniformizing VHS on a compact Sasakian manifold, the constructed map collapses all the points on each leaf of the foliation of the universal covering into one point and can never be an isomorphism. I will explain how to properly take the lift of this map and construct an isomorphism from the universal covering to a homogeneous space.

- 16 宮武夏雄 (九大IMI) 正則ではない Higgs 場を持つ Higgs 束の Donaldson 汎関数の対角形の計量全体への制限 15

Natsuo Miyatake (Kyushu Univ.) Restriction of Donaldson's functional to diagonal metrics on Higgs bundles with not-holomorphic Higgs fields

概要 A Higgs bundle over a compact Kähler manifold X is the pair (E, θ) consisting of a holomorphic vector bundle $E \rightarrow X$ and a holomorphic section θ of $\text{End}E \otimes \wedge^{1,0}$ satisfying $\theta \wedge \theta = 0$. The holomorphic section θ is called a Higgs field. Usually, Higgs fields are assumed to be holomorphic. However, in this talk, I will consider Higgs bundles with not holomorphic Higgs fields. I will suppose that the holomorphic vector bundle E decomposes into a direct sum of holomorphic line bundles, and I will give some necessary and sufficient conditions for Donaldson's functional to attain a minimum over diagonal metrics concerning the decomposition. I will also slightly extend the theorem for not-holomorphic splittings of the vector bundle.

- 17 宮武夏雄 (九大IMI) 巡回 Higgs 束の Hermitian–Einstein 方程式の拡張とその放物型方程式と Dirichlet 問題 15

Natsuo Miyatake (Kyushu Univ.) Some generalizations of the Hermitian–Einstein equation of cyclic Higgs bundles and their parabolic equations and the Dirichlet problem

概要 In this talk, I will introduce some generalizations of the Hermitian–Einstein equation of cyclic Higgs bundles on Riemann surfaces. For such generalized equations, I will show the existence and the uniqueness of the time-global solution of the parabolic equations and I will also show the existence and the uniqueness of the solution to the Dirichlet problem.

- 18 松坂公暉 (北大教育推進機構) Recursion relation on moduli space of quasimaps in the case of Calabi–
秦泉寺雅夫 (岡山大自然) Yau hypersurface in CP^{N-1} 15

Koki Matsuzaka (Hokkaido Univ.) Recursion relation on moduli space of quasimaps in the case of Calabi–
Masao Jinzenji (Okayama Univ.) Yau hypersurface in CP^{N-1}

概要 In computing of B-model in classical mirror symmetry, Picard–Fuchs equation is very important. In the case of Calabi–Yau hypersurface in CP^{N-1} , its solutions are used in a mirror map, and it is known that a generating function of intersection numbers $w(\mathcal{O}_{h^{N-3}}\mathcal{O}_{h^0})_{0,d}$ of moduli space $\widetilde{M}p_{0,2}(N, d)$ of quasimaps from CP^1 with 2 marked points to CP^{N-1} coincides with the mirror map. In our previous work, we proved that the solutions are written by using a term $d \cdot w(\sigma_j(\mathcal{O}_{h^{N-2-j}}\mathcal{O}_{h^{-1}})_{0,d} + w(\sigma_{j-1}(\mathcal{O}_{h^{N-1-j}}\mathcal{O}_{h^{-1}})_{0,d}$, where σ_j corresponds to the Mumford–Morita–Miller class. In this talk, we see that the above term can be obtained from a recursion relation on moduli space $\widetilde{M}p_{0,2|1}(N, d)$ of quasimaps from CP^1 with $2+1$ marked points to CP^{N-1} .

- 19 小林和志 (阪大理) ある gerbe による複素トーラスの変形とホモロジー的ミラー対称性 15

Kazushi Kobayashi (Osaka Univ.) A gerby deformation of complex tori and the homological mirror symmetry

概要 Let (X, \check{X}) be a mirror pair of a complex torus X and its mirror partner \check{X} . This mirror pair is described as the trivial special Lagrangian torus fibrations $X \rightarrow B$ and $\check{X} \rightarrow B$ on the same base space B by SYZ construction. Then, we can associate a holomorphic line bundle $E(s, \mathcal{L}) \rightarrow X$ to a pair (s, \mathcal{L}) of a Lagrangian section s of $\check{X} \rightarrow B$ and a unitary local system \mathcal{L} along it. In this talk, we first construct the deformation $X_{\mathcal{G}}$ of X by a certain flat gerbe \mathcal{G} and its mirror partner $\check{X}_{\mathcal{G}}$ from the mirror pair (X, \check{X}) , and discuss deformations of objects $E(s, \mathcal{L})$ and (s, \mathcal{L}) over the deformed mirror pair $(X_{\mathcal{G}}, \check{X}_{\mathcal{G}})$.

- 20 中西 隼斗 (千葉大融合理工) モースホモトピーを用いたトーリックファノ曲面のホモロジー的ミラー対称性 15

Hayato Nakanishi (Chiba Univ.) Homological mirror symmetry of toric Fano surfaces via Morse homotopy

概要 Strominger–Yau–Zaslow (SYZ) proposed a way of constructing mirror pairs as pairs of torus fibrations. We apply this SYZ construction to toric Fano surfaces as complex manifolds, and discuss the homological mirror symmetry, where we consider Morse homotopy of the moment polytope instead of the Fukaya category.

- 21 山本 悠登 (IBS-CGP) 超曲面の周期積分とトロピカル幾何学 15

Yuto Yamamoto (IBS-CGP) Period integrals of hypersurfaces via tropical geometry

概要 Let $\{Z_t\}_t$ be a one-parameter family of complex toric hypersurfaces of dimension $d \geq 1$. We compute asymptotics of period integrals for $\{Z_t\}_t$ by applying the method of Abouzaid–Ganatra–Iritani–Sheridan, which uses tropical geometry. As integrands, we consider Poincaré residues of meromorphic $(d+1)$ -forms on the ambient toric variety, which have poles along the hypersurface Z_t . The cycles over which we integrate them are spheres and tori which correspond to tropical $(0, d)$ -cycles and $(d, 0)$ -cycles on the tropicalization of $\{Z_t\}_t$ respectively.

13:00~14:00 特別講演

橋本 義規 (阪公大理) 因子に沿って錐的特異点をもつ定スカラー曲率 Kähler 計量についての最近の進展

Yoshinori Hashimoto (Osaka Metro. Univ.) Recent developments on constant scalar curvature Kähler metrics with cone singularities along a divisor

概要 We present some recent results concerning constant scalar curvature Kähler metrics with cone singularities along a divisor, henceforth abbreviated as cscK cone metrics. One of the main results is that the existence of cscK cone metrics implies various stability conditions of the underlying pair of the manifold and the divisor, including G -uniform K -stability and K -polystability. We also prove that any Kähler manifold admits a cscK cone metric if the divisor is a generic member of the linear system defined by a sufficiently large multiple of the polarisation, and point out an analogy to the twisted constant scalar curvature Kähler metrics. This talk is based on a joint work with Takahiro Aoi and Kai Zheng.

3月17日(金) 第II会場

9:30~11:40

- 22 泉 宏明 (広島大) ゲーム理論の意義と概日リズムの定式化 15

Hiroaki Izumi (Hiroshima Univ.) Significance of game theory and formulization of in-phase and anti-phase circadian rhythms

概要 In this paper, we explain the significance of game theory, which frees a system of differential equations from the spell of integrability conditions. Also using it, we present the system of equations of in-phase and anti-phase circadian rhythms from the viewpoint of variational principle. In addition, the general formulization of the mathematical structure of phase transition and synchronization is carried out.

- 23 軸丸 芳揮 (九大IMI) 可積分幾何に基づく Michell トラス型構造と離散対数型美的曲線 15
 早川 健太郎 (京大工)
 林 和希 (京大工)
 梶原 健司 (九大IMI)
 横須賀 洋平 (鹿児島大理工)
 Yoshiki Jikumaru (Kyushu Univ.) On the Michell truss-like structure and discrete log-aesthetic curves
 Kentaro Hayakawa (Kyoto Univ.) based on integrable geometry
 Kazuki Hayashi (Kyoto Univ.)
 Kenji Kajiwara (Kyushu Univ.)
 Yohei Yokosuka (Kagoshima Univ.)

概要 The truss structure minimizing the product of axial forces and lengths for a given load and a support condition is called Michell truss, which is classically known in the field of structural mechanics. In this presentation, we have an analytical construction of Michell truss-like structure described by a discrete holomorphic function (discrete conformal map) in the context of discrete integrable systems. The validity of the structure is demonstrated by the result of structural analysis. Moreover, we show the truss structure is composed of so-called discrete log-aesthetic curves. This structure connects three objects (discrete holomorphic functions, discrete log-aesthetic curves, and classical truss structures) having completely different origins which were achieved through the collaboration of mathematics and architecture.

- 24 陶山 芳彦 (福岡大理) Generic で共形平坦な超曲面内の曲率曲面の拡張と近似 15
 松浦 望 (久留米工大工)
 Yoshihiko Suyama (Fukuoka Univ.) Extension and approximation of curvature surfaces in generic conformally flat hypersurfaces
 Nozomu Matsuura (Kurume Inst. of Tech.)

概要 We talk about curvature surfaces of generic conformally flat analytic-hypersurfaces in the Euclidean 4-space.

- 25 石渡 聡 (山形大理) リーマン多様体上の非対称拡散過程の離散近似 10
 河備 浩司 (慶大経済)
 Satoshi Ishiwata (Yamagata Univ.) A discretization of a non-symmetric diffusion on a Riemannian manifold
 Hiroshi Kawabi (Keio Univ.)

概要 In this talk, we give a graph discretized approximation of the diffusion for a Schrödinger operator with drift on a complete Riemannian manifold.

- 26 張 頁 (沖縄科学技術大) Gradient estimates for the heat semigroup on step-two Carnot groups 15
 Ye Zhang Gradient estimates for the heat semigroup on step-two Carnot groups
 (Okinawa Inst. of Sci. and Tech. Grad. Univ.)

概要 It is well-known that we can characterize the lower bound of the Ricci curvature by the bound for the gradient of the heat semigroup. However, on a nonabelian nilpotent Lie group endowed with a left-invariant Riemannian metric, there exists a direction of strictly negative Ricci curvature. This implies the gradient bound has exponential growth. In this talk, I will present how to improve the gradient bound to the constant in the setting of step-two Carnot groups on which the H.-Q. Li inequality holds.

- 27 武富雄一郎 (阪 公 大 理) リーマン計量の moduli 空間の極大元について 15
 Yuichiro Taketomi A maximal element of a moduli space of Riemannian metrics
 (Osaka Metro. Univ.)

概要 For a given smooth manifold X , we consider the moduli space of Riemannian metrics on X up to isometry and scaling. One can define a preorder on the moduli space by the size of isometry groups. We call a Riemannian metric that attains a maximal element with respect to the preorder a maximal metric. Maximal metrics give nice examples of self-similar solutions for various metric evolution equations such as the Ricci flow. In this talk, we construct many examples of maximal metrics on Euclidean spaces.

- 28 松本佳彦 (阪 大 理) 写像の繰り込みエネルギーと共形測地線 15
 Yoshihiko Matsumoto (Osaka Univ.) Renormalized energy of maps and conformal geodesics

概要 We introduce a certain renormalized energy of maps between asymptotically hyperbolic Einstein manifolds. Then, it is used to give a holographic description of conformal geodesics on the boundary at infinity, in a way deeply inspired by a work of Fine and Herfray on renormalized area minimization and conformal geodesics.

14:15~14:55

- 29 雪田友成 (早 大 教 育) On nerves and growth rates of 2-dimensional Coxeter systems 15
 Tomoshige Yukita (Waseda Univ.) On nerves and growth rates of 2-dimensional Coxeter systems

概要 For a Coxeter system (Γ, S) , the nerve $L(\Gamma, S)$ is a simplicial complex defined via the structure of finite parabolic subgroups of Γ . The dimension of (Γ, S) is defined by $\dim L(\Gamma, S) + 1$. In this talk, we consider the relationship between the combinatorial structure of $L(\Gamma, S)$ and the arithmetic properties of the growth rate of (Γ, S) . We show that the growth rate of a 2-dimensional Coxeter system (Γ, S) with vanishing (resp. positive) Euler characteristic $\chi(L(\Gamma, S))$ is a Salem (resp. Pisot) number. This is a joint work with Naomi Bredon.

- 30 牛島 颯 (兵庫県立国際商経) ケーラー・クライン幾何学としての双曲平面でのチェバの定理とメネラウスの定理 10
 Akira Ushijima (Univ. of Hyogo) Ceva's and Menelaus' theorems in the hyperbolic plane as Cayley–Klein geometry

概要 Using the construction of the hyperbolic plane as Cayley–Klein geometry, we will demonstrate how to obtain Ceva's and Menelaus' theorems in the hyperbolic plane from their corresponding theorems in the real projective plane.

15:10~16:10 特別講演

- H. Seshadri 正の双正則断面曲率を持つケーラー多様体の直径について
 (Indian Inst. of Sci., Bangalore)
 Harish Seshadri On the diameter of Kähler manifolds with positive bisectional curvature
 (Indian Inst. of Sci., Bangalore)

概要 I will discuss the Kähler analogues of the classical Bishop volume inequality and Myers–Bonnet diameter bound for Riemannian manifolds. In particular, I will outline the proof of diameter rigidity (an analogue of Cheng's theorem in the Riemannian case) for compact Kähler n -manifolds (M, ω) with bisectional curvature $BK \geq 1$. It is known from the work of Li–Wang that $\text{diam}(M) \leq \text{diam}(\mathbb{C}P^n)$ for such manifolds, where $\mathbb{C}P^n$ is endowed the Fubini–Study metric. In joint work with V. Datar, we proved that if $\text{diam}(M) = \text{diam}(\mathbb{C}P^n)$, then M is isometric to $\mathbb{C}P^n$.

函数論

3月15日(水) 第V会場

9:30~11:45

- 1 齋藤三郎 (群馬大*再生核研) Complex analysis and theory of reproducing kernels 10
 Saburoou Saitoh Complex analysis and theory of reproducing kernels
 (Gunma Univ.*/Inst. of Reproducing Kernels)

概要 The theory of reproducing kernels is very fundamental, beautiful and will have many applications in analysis, numerical analysis and data sciences. Based on a draft paper, I would like to give a global viewpoint for the relation of complex analysis and the theory of reproducing kernels. I listed the basic references on the topics.

- 2 齋藤三郎 (群馬大*再生核研) Complex analysis and division by zero calculus 15
 Saburoou Saitoh Complex analysis and division by zero calculus
 (Gunma Univ.*/Inst. of Reproducing Kernels)

概要 Based on a draft book manuscript, I would like to introduce the basic and essential relations of complex analysis and division by zero calculus. Basic references are cited in the references.

- 3 綾野孝則 (阪公大数学研) 種数3の超楕円関数と楕円関数の間の関係式 15
 V. M. Buchstaber
 (Steklov Inst. of Math.)
 Takanori Ayano (Osaka Metro. Univ.) A relationship between hyperelliptic functions of genus 3 and elliptic
 Victor M. Buchstaber functions
 (Steklov Inst. of Math.)

概要 In the nineteenth century, the reductions of hyperelliptic integrals to elliptic integrals were studied actively and many examples of the reductions were obtained. They are closely related to coverings of curves, split Jacobians, and automorphism groups of curves. This knowledge is applied to the isogeny-based cryptography. In our previous work, by using this knowledge, we derived the relationships between the hyperelliptic functions of genus 2 and the elliptic functions when the curve of genus 2 admits a morphism of degree 2 to an elliptic curve. In this talk, we will derive a relationship between the hyperelliptic functions of genus 3 and the elliptic functions when the hyperelliptic curve of genus 3 admits a morphism of degree 2 to an elliptic curve.

- 4 杉山真吾 (日大理工) 同変関数の存在と Schwarz 微分 10
 Shingo Sugiyama (Nihon Univ.) The existence of equivariant functions and the Schwarzian derivative

概要 Equivariant functions associated to a 2-dimensional representation of a Fuchsian group was introduced by Sabar and Sebbar. They are regarded as a generalization of automorphic functions. We prove the existence of equivariant functions associated to any generic 2-dimensional representation of any Fuchsian group. We prove such existence by using the Schwarzian derivative.

- 5 山岸 義和 (龍谷大先端理工) 回転対称性を加えた対数螺旋格子上の円充填とボロノイ分割 10
 上 蘭 拓 郎
 (マースネットワークス)
 須志田 隆道 (サレジオ工高専)
 Yoshikazu Yamagishi (Ryukoku Univ.) Voronoi tiling and circle packing on spiral lattices with rotational sym-
 Takuro Uezono (Mars Networks Corp.) metry
 Takamichi Sushida (Salesian Polytech.)

概要 It is shown that the bifurcation diagram of circle packings on logarithmic spiral lattices with rotational symmetry is graph-theoretically dual to the bifurcation diagram of Voronoi tessellations, by using the relative metric. If the rotation parameter (called divergence angle) is badly approximable, then the aspect ratio of the quadrilateral Voronoi cells is bounded. If the divergence angle is linearly equivalent to the golden section, then the shape of the quadrilateral cells tend to square as the plastochron ratio tends to 1.

- 6 熊谷 駿 (東北大情報) 折り紙のボロノイ分解について 15
 Shun Kumagai (Tohoku Univ.) Voronoi decomposition of origamis

概要 The Edwards–Sanderson–Schmidt method uses the Voronoi decomposition to present an excellent criterion for the equivalence class and the Veech group of a finite translation surface. The key argument of their method includes a complicated combinatorial structure. An origami is a typical example of a translation surface defined combinatorially. This talk discusses the Voronoi decomposition of a half-translation version of origami.

- 7 馬場 伸平 (阪大理) Bending maps of Teichmüller spaces and character varieties 15
 Shinpei Baba (Osaka Univ.) Bending maps of Teichmüller spaces and character varieties

概要 The space holomorphic quadratic differentials on a compact Riemann surface is a complex vector space. By the holonomy map of complex projective structures, this vector space properly embeds into the $\mathrm{PSL}(2, \mathbb{C})$ -character variety of the base surface. We construct certain analogues of such an embedding from the viewpoint of Thurston’s parametrization of complex projective structures.

- 8 小森 洋平 (早大教育) Applications of canonical polygons of Zieschang–Vogt–Coldewey 15
 Yohei Komori (Waseda Univ.) Applications of canonical polygons of Zieschang–Vogt–Coldewey

概要 As applications of canonical polygons of Zieschang–Vogt–Coldewey, we will show that (1) $6g-6$ angles between geodesics on Riemann surfaces of genus g globally parametrize Teichmüller space of genus g surfaces. (2) Mapping classes of genus 2 surface can be represented by the cut and paste argument of canonical polygons.

- 9 木坂 正史 (京大人間環境) Polynomial-like mappings and transcendental entire functions with ir-
 rationally indifferent fixed points 15
 Masashi Kisaka (Kyoto Univ.) Polynomial-like mappings and transcendental entire functions with ir-
 rationally indifferent fixed points

概要 In this talk, by using the theory of polynomial-like mappings, we construct various kinds of transcendental entire functions with irrationally indifferent fixed points. This talk is based on the joint work with Hiroto Naba.

14:15~15:15 特別講演

中 西 敏 浩 (島根大総理工) おもに種数 2 のタイヒミュラー空間とクライン群について

Toshihiro Nakanishi (Shimane Univ.) Teichmüller space of the genus two surface and Kleinian groups

概要 In this talk the Teichmüller space is a deformation space of marked Fuchsian groups of finite type, in most cases, genus two Fuchsian groups. We deal with (1) trace functions as parameters for a Teichmüller space, (2) matrix representation of the genus two surface group, (3) mapping classes represented by rational transformations. These materials are used to produce some examples of Kleinian groups related to surface groups.

15:30~16:30 2022年度(第21回)日本数学会解析学賞受賞特別講演

松 崎 克 彦 (早 大 教 育) 絶対連続タイヒミュラー空間の曲線と関数空間

Katsuhiko Matsuzaki (Waseda Univ.) Curves and function spaces of absolutely continuous Teichmüller spaces

概要 The universal Teichmüller space is the total space containing all quasiconformal Teichmüller spaces and it can be regarded as the space of normalized quasisymmetric functions f on the real line. According to the regularity of these functions f , there are two different directions of subspaces included in the universal Teichmüller space: Teichmüller spaces of Riemann surfaces are typically in the direction of totally singular f , and Teichmüller spaces of function spaces are absolutely continuous f .

Teichmüller spaces can be represented by quasisymmetric homeomorphisms mentioned above, and in other ways, they can be represented by the complex dilatations of quasiconformal mappings, the Schwarzian or pre-Schwarzian derivatives of conformal mappings, and the quasicircles given as the images of the circle. Fundamental studies of particular Teichmüller spaces are composed by the investigation of the relationship between these different representations and structures. In addition, if we deal with absolutely continuous Teichmüller spaces, the derivative f' or $\log f'$ defines the corresponding function spaces, and by using them, we can introduce simpler structures to these Teichmüller spaces.

In this talk, we exhibit several important examples of absolutely continuous Teichmüller spaces and explain how the identification of different representations works. The examples include Weil–Petersson Teichmüller spaces, BMO Teichmüller spaces and the space of chord-arc curves between them. One technical tool is the simultaneous uniformization imported from the theory of quasifuchsian spaces. We represent the space of particular quasicircles by the product of the corresponding Teichmüller spaces and consider biholomorphic correspondence between this parameter space and the complex function space. Furthermore, we apply this method to see (dis)continuous dependence of the Riemann mapping parametrizations of the family of Weil–Petersson curves and chord-arc curves to their arc-length parametrizations.

3月16日(木) 第V会場

9:30~11:45

10 上 野 康 平 (大 同 大) Dynamics of superattracting skew products 15

Kohei Ueno (Daido Univ.) Dynamics of superattracting skew products

概要 We consider the dynamics of a holomorphic skew product with a superattracting fixed point. We proved that it is generally conjugate to a monomial map on an invariant wedge whose closure contains the fixed point. In this talk we provide several kinds of plurisubharmonic functions on the complement of specific fibers in the attracting basin, which describe the vertical dynamics well.

- 11 Shaolin Chen (Hengyang Normal Univ.) Lipschitz constant of harmonic Bloch functions and composition operators on Bloch and Hardy type spaces 15
濱田英隆 (九州産大理工)
Jian-Feng Zhu (Huaqiao Univ.)
Shaolin Chen (Hengyang Normal Univ.) Lipschitz constant of harmonic Bloch functions and composition operators on Bloch and Hardy type spaces
Hidetaka Hamada (Kyushu Sangyo Univ.)
Jian-Feng Zhu (Huaqiao Univ.)

概要 The main purpose of this talk is to discuss Hardy type spaces, Bloch type spaces and the composition operators of complex-valued harmonic functions. We first establish a sharp estimate of the Lipschitz continuity of complex-valued harmonic functions in Bloch type spaces with respect to the pseudo-hyperbolic metric, which gives an answer to an open problem. Then some classes of composition operators on Bloch and Hardy type spaces will be investigated. The obtained results improve and extend some corresponding known results.

- 12 濱田英隆 (九州産大理工) Fekete–Szegő problem for univalent mappings in one and higher dimensions 15
G. Kohr (Babeş-Bolyai Univ.)
M. Kohr (Babeş-Bolyai Univ.)
Hidetaka Hamada (Kyushu Sangyo Univ.) Fekete–Szegő problem for univalent mappings in one and higher dimensions
Gabriela Kohr (Babeş-Bolyai Univ.)
Mirela Kohr (Babeş-Bolyai Univ.)

概要 In this talk, we will give the Fekete–Szegő inequality for the mappings f in various subclasses of normalized univalent mappings which are the first elements of g -Loewner chains on the unit disc \mathbb{U} in \mathbb{C} and also on the unit ball \mathbb{B} of a complex Banach space. As an application, we give the estimation of the third coefficient for f under the condition that the second coefficient of f is zero. This result gives a generalization of the estimation of the third coefficient for odd univalent functions on the unit disc \mathbb{U} . We also give the Fekete–Szegő inequality for the images of the first elements of g -Loewner chains on \mathbb{U} under the Roper–Suffridge type extension operators.

- 13 濱田英隆 (九州産大理工) Loewner PDE in infinite dimensions 15
G. Kohr (Babeş-Bolyai Univ.)
Hidetaka Hamada (Kyushu Sangyo Univ.) Loewner PDE in infinite dimensions
Gabriela Kohr (Babeş-Bolyai Univ.)

概要 In this talk, we generalize the existence and uniqueness result for the solutions of the Loewner partial differential equation (Loewner PDE) from \mathbb{C}^n to the case of separable reflexive complex Banach spaces.

- 14 Shaolin Chen (Hengyang Normal Univ.)
濱田英隆 (九州産大理工)
Shaolin Chen (Hengyang Normal Univ.)
Hidetaka Hamada (Kyushu Sangyo Univ.)
- Some sharp Schwarz–Pick type estimates and their applications of harmonic and pluriharmonic functions 15
- Some sharp Schwarz–Pick type estimates and their applications of harmonic and pluriharmonic functions

概要 The purpose of this talk is to study the Schwarz–Pick type inequalities for harmonic or pluriharmonic functions. By analogy with the generalized Khavinson conjecture, we first give some sharp estimates of the norm of harmonic functions from the Euclidean unit ball in \mathbb{R}^n into the unit ball of the real ℓ_p space. Next, we give several sharp Schwarz–Pick type inequalities for pluriharmonic functions from the Euclidean unit ball in \mathbb{C}^n or from the unit polydisc in \mathbb{C}^n into the unit ball of the ℓ_p space. Furthermore, we establish some sharp coefficient type Schwarz–Pick inequalities for pluriharmonic functions defined in the ℓ_p space. Finally, we use the obtained Schwarz–Pick type inequalities to discuss the Schwarz–Pick type lemmas of arbitrary order and the Bohr phenomenon of harmonic or pluriharmonic functions.

- 15 Shaolin Chen (Hengyang Normal Univ.)
濱田英隆 (九州産大理工)
S. Ponnusamy (Indian Inst. of Tech. Madras)
R. Vijayakumar (Indian Inst. of Tech. Madras)
Shaolin Chen (Hengyang Normal Univ.)
Hidetaka Hamada (Kyushu Sangyo Univ.)
Saminathan Ponnusamy (Indian Inst. of Tech. Madras)
Ramakrishnan Vijayakumar (Indian Inst. of Tech. Madras)
- Schwarz type lemmas and their applications in Banach spaces 15
- Schwarz type lemmas and their applications in Banach spaces

概要 The main purpose of this talk is to develop some methods to investigate the Schwarz type lemmas of holomorphic mappings and pluriharmonic mappings in Banach spaces. Initially, we extend the classical Schwarz lemmas of holomorphic mappings to Banach spaces, and then we apply to establish a sharp Bloch type theorem for pluriharmonic mappings on homogeneous unit balls of \mathbb{C}^n and to obtain some sharp boundary Schwarz type lemmas for holomorphic mappings in Banach spaces. Next, we improve and generalize the classical Schwarz lemmas of planar harmonic mappings into the sharp forms of Banach spaces and give some applications to sharp boundary Schwarz type lemmas for pluriharmonic mappings in Banach spaces. Finally, we give some sharp Schwarz–Pick type estimates of pluriharmonic mappings in JB^* -triples.

- 16 渡邊祐太 (東大数理) 乗数イデアル層で振じれた対数的微分形式の層に対する L^2 -型 Dolbeault 同型と消滅定理 15
- Yuta Watanabe (Univ. of Tokyo) L^2 -type Dolbeault isomorphisms and vanishing theorems for logarithmic sheaves twisted by multiplier ideal sheaves

概要 In this speech, we first establish an L^2 -type Dolbeault isomorphism for the sheaf of logarithmic differential forms twisted by the multiplier ideal sheaf. By using this isomorphism and L^2 -estimates equipped with a singular Hermitian metric, we obtain logarithmic vanishing theorems involving multiplier ideal sheaves on compact Kähler manifolds with simple normal crossing divisors.

- 17 大 沢 健 夫 (名大多元数理)b On the Levi problem on Kähler manifolds under the negativity of canonical bundles on the boundary 15
- Takeo Ohsawa (Nagoya Univ.) On the Levi problem on Kähler manifolds under the negativity of canonical bundles on the boundary

概要 It is proved that a bounded smooth pseudoconvex domain D in a Kähler manifold M can be mapped onto a locally closed analytic set in a complex number space holomorphically and properly with connected fibers if the canonical bundle of M is negative on a neighborhood of the boundary of D . A similar result holds for Zariski open domains of compact manifolds.

13:00~14:00 特別講演

- 阿 部 誠 (広島大先進理工) Stein 多様体上の岡・Grauert の原理をみたす領域について
- Makoto Abe (Hiroshima Univ.) Domains over a Stein manifold satisfying the Oka–Grauert principle

概要 We discuss the pseudoconvexity and the Steinness of a complex manifold or a complex space under some conditions. Among others, we announce the recent joint research with Sugiyama on the Steinness of a domain over a Stein manifold that satisfies the Oka–Grauert principle for a complex Lie group of positive dimension and some additional cohomological conditions.

関数方程式論

3月15日(水) 第IV会場

9:30~12:00

- 1 竹井優美子 (茨城工高専) (1, 1, 1, 2)-超幾何微分方程式の Voros 係数の位相的漸化式による表示とその応用 12
 Yumiko Takei On the expression of Voros coefficients for the hypergeometric differential equation of type (1,1,1,2) in terms of the topological recursion, and its applications
 (Ibaraki Nat. Coll. of Tech.)

概要 Voros coefficients are important objects in the exact WKB analysis for the global study of solutions of differential equations. In this talk I will report that the Voros coefficients for the hypergeometric differential equation of type (1,1,1,2) are given by the generating functions of free energies defined in terms of Eynard and Orantin's topological recursion.

- 2 松原幸栄 (熊本大先端) The signature of the invariant hermitian form for a regular holonomic
 後藤良彰 (小樽商大) GKZ system 12
 Saiei-Jaeyeong Matsubara-Heo The signature of the invariant hermitian form for a regular holonomic
 (Kumamoto Univ.) GKZ system
 Yoshiaki Goto
 (Otaru Univ. of Commerce)

概要 Frits Beukers and Carlos Verschoor conjectured that the signature of the monodromy invariant hermitian form for a regular holonomic GKZ system has an explicit formula. We prove that this conjecture is true, by computing the intersection matrix of the twisted homology group associated to Euler type integral representation of a solution to GKZ system.

- 3 伊藤秀一 (神奈川大工) シンプレクティック写像族のバーコフ標準化とハミルトン系の超可積分性
 12
 Hidekazu Ito (Kanagawa Univ.) Birkhoff normalization of a family of symplectic maps and superintegrability of Hamiltonian systems

概要 Integrability of a Hamiltonian system is defined by the existence of sufficiently many (Poisson-commuting) integrals. The Liouville–Arnold theorem gives description of the orbit structure near a compact regular level set of those integrals, however, is not valid near singularities of those integrals. The aim of this talk is to clarify the orbit structure of a super (noncommutatively) integrable system near singularities of its integrals. For this purpose, Birkhoff normalization for a family of symplectic maps plays a key role. We show that there exists a convergent Birkhoff normalization if a family of $2n$ -dimensional analytic symplectic maps has $n + q$ integrals near a fixed point of resonance degree q .

- 4 上村豊 (東京海洋大*) エネルギー依存逆散乱の2成分ソリトン 12
 Yutaka Kamimura Two-component solitons via an energy dependent inverse scattering
 (Tokyo Univ. of Marine Sci. and Tech.*)

概要 An extended KdV hierarchy is discussed. Based upon an inverse scattering method on an energy dependent Schroedinger operator, soliton solutions in the hierarchy are constructed in a unified manner.

- 5 畑 裕貴 (阪府大理) 2つの時間遅れをもつ線形微分方程式の漸近安定性 12
 松 永秀章 (阪公大理)
 Yuki Hata (Osaka Pref. Univ.) Asymptotic stability in a linear differential equation with two delays
 Hideaki Matsunaga
 (Osaka Metro. Univ.)

概要 This study is devoted to the investigation of the effect of delays on the asymptotic stability of a linear differential equation with two delays $x'(t) = -ax(t) - bx(t - \tau) - cx(t - 2\tau)$, $t \geq 0$, where a , b , and c are real numbers and $\tau > 0$. We establish some explicit conditions for the zero solution of the equation to be asymptotically stable.

- 6 鬼塚 政一 (岡山理大理) On Ulam stability of two-dimensional linear differential systems 10
 Masakazu Onitsuka On Ulam stability of two-dimensional linear differential systems
 (Okayama Univ. of Sci.)

概要 The main purpose of this talk is to classify the Ulam stability of the linear differential equation $\mathbf{u}' = A\mathbf{u}$, where A is a real-valued constant 2×2 matrix. The best Ulam constant is obtained for some special cases of A .

- 7 宇佐美 広介 (岐阜大工) 非線形項に単調性等を仮定しない常微分方程式の正值解の非存在定理について 10
 Hiroyuki Usami (Gifu Univ.) Study on nonexistence results of positive solutions of ODEs without assuming monotonicities on nonlinear terms

概要 Sufficient conditions for the existence of positive solutions of quasilinear ODEs are given. Usually, such results are proved under the monotonicity conditions of nonlinear terms. However, in this talk we will give such results without assuming monotonicities on nonlinear terms.

- 8 柴田 徹太郎 (広島大先進理工) Global and asymptotic behaviors of bifurcation curves of one-dimensional nonlocal elliptic equations 12
 Tetsutaro Shibata (Hiroshima Univ.) Global and asymptotic behaviors of bifurcation curves of one-dimensional nonlocal elliptic equations

概要 We consider the global behavior of the bifurcation curves and solution curves of one-dimensional nonlocal elliptic eigenvalue problems. We know that the bifurcation curve λ is a continuous function of the maximum norm $\alpha = \|u_\lambda\|_\infty$ such as $\lambda(\alpha)$. Here, u_λ is the positive solution associated with λ . Our purpose here is to establish the asymptotic formulas for $\lambda(\alpha)$ and u_λ as $\lambda \rightarrow \infty$.

- 9 濱本 直樹 (阪公大理) The Poincaré constant for curl-free vector fields on a ball 12
 Naoki Hamamoto (Osaka Metro. Univ.) The Poincaré constant for curl-free vector fields on a ball

概要 We investigate the best value of the constant in classical Poincaré inequality

$$\int_{B^N} |\nabla \mathbf{u}|^2 dx \geq C \int_{B^N} |\mathbf{u}|^2 dx$$

for a class of vector fields \mathbf{u} with $\mathbf{u}|_{\partial B^N} = \mathbf{0}$ on the unit ball $B^N \subset \mathbb{R}^N$ which are subject to the curl-free constraint. We find that the best value of C turns out to be the square of the first zero of Bessel function of the $(N/2)$ -th order.

- 10 齋藤三郎 (群馬大*・再生核研) Functional equations and theory of reproducing kernels 10
 Saburoou Saitoh Functional equations and theory of reproducing kernels
 (Gunma Univ.*/Inst. of Reproducing Kernels)

概要 By a survey paper, we shall introduce the basic relations of functional equations and the theory of reproducing kernels, globally, as a summary of the topics. In particular, I would like to introduce typical results with their principles in some concrete way based on the draft paper.

The general theory of reproducing kernels will give the basic tool of these widespread problems.

- 11 齋藤三郎 (群馬大*・再生核研) Representations of quotients $g/f = h$ in terms of $g = fh$ 12
 Saburoou Saitoh Representations of quotients $g/f = h$ in terms of $g = fh$
 (Gunma Univ.*/Inst. of Reproducing Kernels)

概要 We are considering the simple functional equation $hf = g$. By the paper (REPRESENTATIONS OF QUOTIENTS $g/f = h$ IN TERMS OF $g = fh$, International Journal of Reproducing Kernels Vol. 2 No. 1 (2023)), we shall introduce quotients g/f of general functions f and g in some deep and natural meanings, in a natural setting.

14:15~16:15

- 12 堀内利郎 (茨城大理) On general Caffarelli–Kohn–Nirenberg type inequalities involving non-doubling weights 12
 Toshio Horiuchi (Ibaraki Univ.) On general Caffarelli–Kohn–Nirenberg type inequalities involving non-doubling weights

概要 By introducing $W(R_+) = P(R_+) \cup Q(R_+) = \cup_{a \in [0, \infty]} W_a(R_+)$ as a class of weight functions, we will establish the Caffarelli–Kohn–Nirenberg type inequalities for $p \geq 1$ involving non-doubling weights in $W(R_+)$. The classical Caffarelli–Kohn–Nirenberg type inequalities are categorized into non-critical and critical cases, and it is known that there is some kind of mysterious relationship between them. Interestingly the new framework in this treatise allows them to be integrated and reveals the meaning of mysterious relationships. A function $w(t)$ on $(0, \infty)$ is said to be a doubling weight if there exists a positive number C such that we have $C^{-1}w(t) \leq w(2t) \leq Cw(t)$ ($0 < t < \infty$), where C is independent of each $t \in (0, \infty)$. When $w(t)$ does not possess this property, $w(t)$ is said to be a non-doubling weight in the talk.

- 13 梶木屋龍治 Boundedness of critical points in the symmetric mountain pass lemma 12
 (大阪電通大共通教育機構)
 Ryuji Kajikiya Boundedness of critical points in the symmetric mountain pass lemma
 (Osaka Electro-Comm. Univ.)

概要 We study the boundedness of critical points defined by the symmetric mountain pass lemma. We construct three types of functionals, which have an unbounded sequence of critical values. For the first one, the set of all critical points is bounded and the set of critical values is unbounded. The second one has both an unbounded sequence and a bounded sequence of critical points. For the last one, the set of critical points is countably infinite and unbounded.

- 14 赤木剛朗 (東北大理) Maximal regularity of distributional solutions to degenerate elliptic systems for locally integrable data 12
 宮川寛基 (東北大理)
 Goro Akagi (Tohoku Univ.) Maximal regularity of distributional solutions to degenerate elliptic systems for locally integrable data
 Hiroki Miyakawa (Tohoku Univ.)

概要 This talk is concerned with degenerate elliptic equations involving the p -Laplacian and absorption in unbounded domains. Main results consist of the existence of distributional solutions and their maximal regularity (estimates) for locally integrable forcing terms beyond the duality.

- 15 赤木 剛朗 (東北大理) On some quasistatic evolution equation arising from fracture mechanics
 佐藤光汰朗 (東北大理) 12
 Goro Akagi (Tohoku Univ.) On some quasistatic evolution equation arising from fracture mechanics
 Kotaro Sato (Tohoku Univ.)

概要 This talk deals with some quasistatic evolution equation, which relates fracture mechanics and is also written as an evolutionary variational inequality. The main result of this talk is concerned with the existence of strong solutions complying with three qualitative properties, that is, irreversibility, unilateral equilibrium and energy conservation, which were originally proposed in a phase-field model for brittle fracture.

- 16 岡 大 将 (東 大 工) Corrector results for space-time homogenization of fast diffusion equations without assumptions for smoothness of coefficients 12
 Tomoyuki Oka (Univ. of Tokyo) Corrector results for space-time homogenization of fast diffusion equations without assumptions for smoothness of coefficients

概要 In this talk, we shall discuss a space-time homogenization problem for fast diffusion equations with periodically oscillating (in space and time) coefficients. The main purpose of this talk is to present a corrector result, i.e., strong convergence of gradient of the solution with a corrector term. The proof is based on the unfolding method and the strong two-scale convergence theory.

- 17 鈴木 貴 (阪大 M M D S) ブラウン粒子気体モデルの数学解析 5
 Takashi Suzuki (Osaka Univ.) Mathematical analysis for Brownian particle gas

概要 We study Streater's model on Brownian particle gas in statistical mechanics on bounded domain in \mathbf{R}^n . If $n \leq 2$, there is a solution global in time, provided that the entropy production is bounded.

- 18 谷口 晃一 (東北大 AIMR) Hardy-Hénon 型半線型熱方程式の解の無条件一意性と非一意性 10
 千頭 昇 (名 工 大)
 池田 正弘
 (理化学研 AIP・慶大理工)

S. Tayachi
 (Univ. de Tunis El Manar)

- Koichi Taniguchi (Tohoku Univ.) Unconditional uniqueness and non-uniqueness of solutions for Hardy-Hénon parabolic equations
 Noboru Chikami (Nagoya Inst. of Tech.)
 Masahiro Ikeda (RIKEN/Keio Univ.)
 Slim Tayachi (Univ. de Tunis El Manar)

概要 In this talk we study the problems of uniqueness for Hardy-Hénon parabolic equations, which are semilinear heat equations with the singular potential (Hardy type) or the increasing potential (Hénon type) in the nonlinear term. To deal with the Hardy-Hénon type nonlinearities, we employ the weighted Lorentz spaces as the solution space. The purpose of this talk is to prove unconditional uniqueness and non-uniqueness for Hardy-Hénon parabolic equations in the weighted Lorentz spaces.

- 19 下條 昌彦 (都立大理) Spreading behavior of an SIR model with non-local dispersal 10
 Jong Shenq Guo (Tamkang Univ.)
 Amy Ai-Ling Poh (東 大 工)
 Masahiko Shimojo Spreading behavior of an SIR model with non-local dispersal
 (Tokyo Metro. Univ.)
 Jong Shenq Guo (Tamkang Univ.)
 Amy Ai-Ling Poh (Univ. of Tokyo)

概要 We explain a Liouville type of theorem on the positive bounded entire solution of a class of reaction-diffusion systems with nonlocal dispersal. Some application from epidemiology is also given.

- 20 下條昌彦 (都立大理) Spreading behavior of a predator-prey system with fractional Laplacian
 Jong Shenq Guo (Tamkang Univ.) 12
 Masahiko Shimojo Spreading behavior of a predator-prey system with fractional Laplacian
 (Tokyo Metro. Univ.)
 Jong Shenq Guo (Tamkang Univ.)

概要 We establish a Liouville theorem on the positive bounded entire solution of a class of reaction-diffusion systems with fractional diffusion. As an application, we investigate the spreading phenomena of a predator-prey system with fractional diffusion. More precisely, by a suitable limiting argument along with the Liouville-type theorem, we prove that the solution converges to the unique positive co-existence state behind the spreading front.

16:30~17:30 2022年度(第21回)日本数学会解析学賞受賞特別講演

- 池 畠 優 (広島大工) 時間領域における囲い込み法の展開
 Masaru Ikehata (Hiroshima Univ.) Development of the time domain enclosure method

概要 This talk is concerned with a reconstruction method in inverse problems formulated by using the solutions of various partial differential equations. Such equations are called the governing equations and the observation data are described by the solutions. The object to be reconstructed or extracted from the data is the discontinuity such as obstacles, inclusions, cracks etc.

The probe and enclosure methods are general ideas to extract information about unknown discontinuity embedded in a known background medium from the observation data. The data are given by the far field pattern of the scattered field, the Dirichlet-to-Neumann map or its partial knowledge at the boundary of the medium. The methods were introduced by the speaker and it has already been passed more than twenty years. The methods have been applied to several inverse problems, e.g., an inverse boundary value problem related to EIT (electrical impedance tomography), elasticity and inverse obstacle scattering problems of acoustic wave at a fixed wave number, whose governing equations are time-independent.

The next challenge and interest of the speaker were to extend the enclosure method to cover inverse problems governed by time-dependent PDEs. However, as a result of research over the last ten years, it has been found that the enclosure method is also effective for various time-dependent PDEs, such as the heat equation, the wave equation, a system of equations in thermoelasticity, the Maxwell system and quite recently the fractional diffusion equation.

This talk will give a brief history of probe and enclosure method and then describe some of recent developments of enclosure methods in the time domain.

3月16日(木) 第IV会場

9:15~12:00

- 21 水上雅昭 (京都教育大) Non-simultaneous blow-up in a two-species chemotaxis-competition model
 田中悠也 (東京理大理) with single production 12
 Masaaki Mizukami Non-simultaneous blow-up in a two-species chemotaxis-competition model
 (Kyoto Univ. of Edu.) with single production
 Yuya Tanaka (Tokyo Univ. of Sci.)

概要 In a two-species chemotaxis-competition model, blow-up of solutions was shown in a previous paper. In this model, the production term seems to promote blow-up phenomena; this implies that the lack of the production term makes the solution likely to be bounded. Thus, it is expected that the model without one of the production terms has a solution such that the species which does not produce the chemical substance remains bounded, whereas the other species blows up. The purpose of this paper is to prove that this conjecture is true.

- 22 田中悠也 (東京理大理) Finite-time blow-up in a degenerate parabolic–elliptic Keller–Segel system with logistic source 12

Yuya Tanaka (Tokyo Univ. of Sci.) Finite-time blow-up in a degenerate parabolic–elliptic Keller–Segel system with logistic source

概要 This talk deals with a degenerate parabolic–elliptic Keller–Segel system with logistic source. In the case of nondegenerate diffusion, Black, Fuest and Lankeit (Z. Angew. Math. Phys.; 2021; 72; no. 96, 23 pp.) obtained a condition such that finite-time blow-up occurs. The purpose of this talk is to show that finite-time blow-up occurs also in the case of degenerate diffusion.

- 23 千代祐太郎 (東京理大理) Boundedness of solutions to a chemotaxis system for tumor angiogenesis
水上雅昭 (京都教育大教育) 12

Yutaro Chiyo (Tokyo Univ. of Sci.) Boundedness of solutions to a chemotaxis system for tumor angiogenesis
Masaaki Mizukami
(Kyoto Univ. of Edu.)

概要 This talk deals with a chemotaxis system for tumor angiogenesis. When the third equation is simplified by the elliptic one and the sensitivity functions are constants, global existence and boundedness were proved by Tao and Winkler (Nonlinear Analysis; 2021; 112324; 16 pp). However, the case that the sensitivity functions are not constants and the third equation is parabolic seems not to have been studied yet. The purpose of this talk is to establish global existence and boundedness in a fully parabolic chemotaxis system for tumor angiogenesis with sensitivity functions.

- 24 勝呂剛志 (京大数理研) 放物-楕円型 Keller–Segel 系の初期値問題の一様局所可積分空間における適切性について 12

Takeshi Suguro (Kyoto Univ.) Well-posedness of the Cauchy problem of a parabolic-elliptic Keller–Segel system in uniformly local spaces

概要 We consider the Cauchy problem of the parabolic-elliptic Keller–Segel system on Euclidean space. This system is one of the diffusion equation involving a nonlocal term. It is interesting whether this problem is well-posed in function spaces containing the non-decaying function. We show that the Cauchy problem of a Keller–Segel system is well-posed in uniformly local Lebesgue spaces.

- 25 細野竜也 (東北大理) 4D attraction-repulsion chemotaxis system and applications of Brezis–Merle inequality 12
小川卓克 (東北大理)

Tatsuya Hosono (Tohoku Univ.) Global existence of solutions to the 4D attraction-repulsion chemotaxis system and applications of Brezis–Merle inequality
Takayoshi Ogawa (Tohoku Univ.)

概要 We consider the Cauchy problem for an attraction-repulsion chemotaxis system in the four space dimension. One of main topics in the study of such a system is the presence of L^1 threshold. In fact, the critical mass phenomenon called 8π -problem is well-known in the 2-dimensional setting. In this talk, using the Brezis–Merle type inequality and rearrangement arguments, we will show that the solution exists globally in time when the initial mass is less than 4-dimensional L^1 threshold value $(8\pi)^2$.

- 26 田代紀一 (東工大) Elliptic regularization による適切な一般化平均曲率流の構成について . 10

Kiichi Tashiro (Tokyo Tech) On the construction of canonical generalized mean curvature flow by elliptic regularization

概要 In this talk, we consider a generalized mean curvature flow constructed by elliptic regularization. Elliptic regularization is a method invented by Tom Ilmanen to construct an approximation of the generalized mean curvature flow using real analytic functional. We will show that the area change of the region moved by this flow is given by equality using the generalized mean curvature. The key to the proof is to establish approximate velocity and estimate it in L^2 .

- 27 三浦達哉 (東工大 理) Complete classification of planar p -elasticae 12
 吉澤研介 (九大 I M I)
 Tatsuya Miura (Tokyo Tech) Complete classification of planar p -elasticae
 Kensuke Yoshizawa (Kyushu Univ.)

概要 Euler's elastica is defined by a critical point of the total squared curvature under the fixed length constraint, and its L^p -counterpart is called p -elastica. In our talk, we completely classify all p -elasticae in the plane and obtain their explicit formulae as well as optimal regularity. To this end we introduce new types of p -elliptic functions which streamline the whole argument and result. As an application we also classify all closed planar p -elasticae.

- 28 長澤壯之 (埼玉大理工) 非局所曲率流の存在時間の下界評価と爆発レート 10
 Takeyuki Nagasawa (Saitama Univ.) A lower estimate of the life-span and a blow-up rate for non-local curvature flows for plane curves

概要 Several non-local curvature flows for plane curves with general rotation number are considered here. The flows include the area-preserving flow and the length-preserving flow. Previously, we showed sufficient conditions for finite-time blow-up, and gave an upper bound of the blow-up time. Here, we give estimates for solutions including at least two important applications. As the first one, it gives a lower bound of the life span, which is a lower bound of the blow-up time for blow-up solutions. Secondly, our estimate also improves the blow-up rate of blow-up solutions.

- 29 三浦達彦 (弘前大理工) Error estimate for classical solutions to the heat equation in a moving thin domain and its limit equation 12
 Tatsu-Hiko Miura (Hiroaki Univ.) Error estimate for classical solutions to the heat equation in a moving thin domain and its limit equation

概要 We consider the Neumann type problem of the heat equation in a moving thin domain around a given closed moving hypersurface. Our main result is an estimate in the sup-norm for the difference of a classical solution to the heat equation in the moving thin domain and a classical solution to a limit equation on the moving hypersurface which appears in the thin-film limit of the heat equation. To prove that error estimate, we construct a suitable approximate solution to the heat equation from a classical solution to the limit equation and then estimate the difference of a classical solution and the approximate one to the heat equation based on the maximum principle.

- 30 二宮広和 (明大総合数理) 等拡散によるパターン形成の例 12
 Hirokazu Ninomiya (Meiji Univ.) Example of pattern formation by equal diffusion

概要 In 1952, Turing proposed the mechanism of pattern formation in which a stable equilibrium of some kinetic system is destabilized by diffusion. In the case of two-component reaction-diffusion systems, however, the diffusion coefficients should be different. Conversely, there exist several chemical models that produce the stable patterns even though the diffusion coefficients are equal. To study the pattern formation by equal diffusion, this talk presents an example of a kinetic system with a asymptotically stable equilibrium, while the corresponding reaction-diffusion system has a family of unstable stationary solutions that is arbitrarily close to the homogeneous stationary solution.

- 31 二宮 広和 (明大総合数理) 非一様場における面積保存曲率流のダイナミクス 12
 Hirokazu Ninomiya (Meiji Univ.) Dynamics of area-preserving curvature flow of a convex plane curve in an inhomogeneous medium

概要 In 1986, Gage studied the area-preserving curvature flows in a two-dimensional homogeneous medium. He showed that an initially convex closed curve remains convex and converges to a circle as time goes to infinity. However, in many applications, the medium is not homogeneous. In this talk, we consider the area preserving flow in an inhomogeneous medium when the area enclosed by the interface is small. The reduced equation of its center will be explained.

13:00~14:00 特別講演

- 長澤 壯之 (埼玉大理工) Möbius エネルギーの分解と波動写像
 Takeyuki Nagasawa (Saitama Univ.) The decomposition of the Möbius energies and wave maps

概要 The Möbius energy for knots is known as a functional invariant under Möbius transformations. There is an energy for links having the same properties. For the energies for knots and two-component links, there exists a Möbius invariant decomposition, that is, each decomposed part is still invariant under Möbius transformations.

Recently, the author found two new expressions of decomposition using Gauss maps of knots and links. One is an “indirect expression”, the other is a “direct expression.”

Using the indirect expression, we can regard the decomposition as the *parallelogram law* of the Möbius energy. Furthermore, the cosine formula of the Möbius energy for knots is generalized to decomposed energies.

It has been known that the first decomposed energy relates to an energy of fractional harmonic maps. We can see from the direct expression that the second one does to an energy of wave maps. As an application, we show the existence or non-existence of knots and links which Gauss maps are wave maps.

3月17日(金) 第IV会場

9:30~12:00

- 32 佐藤 洋平 (埼玉大理工) 斥力的連立シュレディンガー方程式の偶対称基底状態 12
 Yohei Sato (Saitama Univ.) Even ground state for nonlinear Schrödinger systems with repulsive interaction

概要 It is known that the nonlinear Schrödinger systems with repulsive interaction has no ground states. Here ground state means a positive solution that attains a minimizing problem on the Nehari set with two constrains. On the other hand, when the minimizing problem is restricted to even functions, one calls a positive minimizer as an even ground state. We show the existence of even ground states of nonlinear Schrödinger systems with repulsive interaction. The keys of the proof are estimates of minimizing level and a classification of Palais-Smale sequences of even functions. In particular, the case $N = 1$ requires more detailed estimates than $N = 2, 3$.

- 33 平山浩之 (宮崎大教育) 微分型の非線形項を持つ非線形シュレディンガー方程式系の基底状態の
池田正弘 存在と安定性について 12
(理化学研 AIP・慶大理工)

Hiroyuki Hirayama (Univ. of Miyazaki) Existence and stability of ground states for the system of nonlinear
Masahiro Ikeda (RIKEN/Keio Univ.) Schrödinger equations with derivative nonlinearity

概要 We consider the Cauchy problem of the system of nonlinear Schrödinger equations with derivative nonlinearity. The well-posedness of this system in the Sobolev space H^1 is obtained by the previous works. But the long time behavior of the solution of this system is not well known. In this talk, we prove the existence of ground state solutions of this system. Furthermore, for 1 dimensional case, we also prove the stability of ground states by using the variational method.

- 34 深谷法良 (東京理大理) 1次元における二重べき型非線形シュレディンガー方程式の定常解の不安
林雅行 (ピサ大・早大理工) 定性 12
Noriyoshi Fukaya (Tokyo Univ. of Sci.) Instability of stationary solutions for double power nonlinear Schrödinger
Masayuki Hayashi equations in one dimension
(Univ. di Pisa/Waseda Univ.)

概要 We consider a double power nonlinear Schrödinger equation in one dimension and L^2 -subcritical setting. The equation has the algebraically decaying stationary solution ϕ_0 as well as exponentially decaying standing waves $e^{i\omega t}\phi_\omega(x)$ with positive frequency. It is known from the general theory that the standing wave with positive frequency is stable if $M'(\omega) > 0$ and unstable if $M'(\omega) < 0$, where $M(\omega) := \frac{1}{2}\|\phi_\omega\|_{L^2}^2$. In this talk we prove the instability of the stationary solution under the condition $M'(0) := \lim_{\omega \downarrow 0} M'(\omega) \in [-\infty, 0)$. The key in the proof is the construction of the one-sided derivative of $\omega \mapsto \phi_\omega$ at $\omega = 0$, which is effectively used to construct the unstable direction.

- 35 佐川侑司 (千葉工大工) 調和ポテンシャル付き非線形シュレディンガー方程式に対する小さな初
佐藤拓也 (東北大理) 期値の有限時間爆発解 12
川上翔汰 (埼玉大理工)
- Yuji Sagawa (Chiba Inst. of Tech.) Finite time blow up solutions to the nonlinear Schrödinger equation
Takuya Sato (Tohoku Univ.) with harmonic potential for arbitrary small initial data
Shota Kawakami (Saitama Univ.)

概要 We construct finite time blow up solutions to the power type nonlinear Schrödinger equation with harmonic potential for arbitrary small initial data in the positive time direction. We simultaneously observe that those solutions exist globally in time and its mass decays in the negative time direction. This part is an improvement of the previous work by Antonelli, Carles and Sparber.

- 36 平山浩之 (宮崎大教育) 2次の非線形項をもつ2次元非線形 Schrödinger 方程式の角度正則性を課
木下真也 (埼玉大理工) した Sobolev 空間での適切性 12
岡本葵 (阪大理)

Hiroyuki Hirayama (Univ. of Miyazaki) Well-posedness of the Cauchy problem for the two dimensional quadratic
Shinya Kinoshita (Saitama Univ.) nonlinear Schrödinger equation with angular regularity
Mamoru Okamoto (Osaka Univ.)

概要 We consider the Cauchy problem for the two dimensional nonlinear Schrödinger equation with the quadratic nonlinearity $|u|^2$. In the Sobolev space $H^s(\mathbb{R}^2)$, it is known that $s = -\frac{1}{4}$ is the threshold for the Cauchy problem to be well-posed. We prove the well-posedness in $H^s(\mathbb{R}^2)$ for $-\frac{1}{2} < s < -\frac{1}{4}$ by assuming some angular regularity on initial data.

- 37 津原 駿 (東北大理) 4次元 Sobolev 臨界の連立非線型 Schrödinger 方程式系の時間大域適切性
 小川 卓克 (東北大理) について 12
 Shun Tsuhara (Tohoku Univ.) Global well-posedness for the Sobolev critical nonlinear Schrödinger sys-
 Takayoshi Ogawa (Tohoku Univ.) tem in four space dimensions

概要 We consider the Cauchy problem of the system of nonlinear Schrödinger equations with cubic nonlinearity in four space dimensions. We prove the existence of the ground state solution and the global well-posedness for the system below the ground state solution under the mass resonance condition. We also show that the ground state solution is written by Aubin–Talenti function. The proof of the global well-posedness result is following the well-known results due to Kenig–Merle (2006).

- 38 佐藤 拓也 (東北大理) The initial boundary value problem of nonlinear Schrödinger equations
 林 仲夫 (東北大理) with a nonlinear Neumann boundary condition 12
 小川 卓克 (東北大理)
 Takuya Sato (Tohoku Univ.) The initial boundary value problem of nonlinear Schrödinger equations
 Nakao Hayashi (Tohoku Univ.) with a nonlinear Neumann boundary condition
 Takayoshi Ogawa (Tohoku Univ.)

概要 We consider the initial-boundary value problem of the nonlinear Schrödinger equation on the half line with a nonlinear Neumann boundary condition. After establishing the boundary Strichartz estimate in L^2 , we consider the time local well-posedness of the problem in the lower regularity spaces as L^2 .

- 39 長田 祐輝 (都立大理) 非対称ポテンシャルの下での3波相互作用をもつ非線形シュレディンガー
 方程式系の minimizer の存在 12
 Yuki Osada (Tokyo Metro. Univ.) Existence of a minimizer for a nonlinear Schrödinger system with three
 wave interaction under non-symmetric potentials

概要 In this talk, we show the existence of a minimizer for the L^2 -constrained minimization problem associated with a nonlinear Schrödinger system with three wave interaction without assuming symmetry for potentials by using interaction estimates.

- 40 池田 正弘 Global dynamics below a threshold for the nonlinear Schrödinger equa-
 (理化学研 AIP・慶大理工) tions with the Kirchhoff boundary and the repulsive Dirac delta bound-
 成 亥 隆 恭 ary on a star graph 12
 (阪大理・プリティッシュコロンビア大)
 浜野 大 (早大理工)
 清水 一慶 (阪大基礎工)
 Masahiro Ikeda (RIKEN/Keio Univ.) Global dynamics below a threshold for the nonlinear Schrödinger equa-
 Takahisa Inui tions with the Kirchhoff boundary and the repulsive Dirac delta bound-
 (Osaka Univ./Univ. of British Columbia) ary on a star graph
 Masaru Hamano (Waseda Univ.)
 Ikkei Shimizu (Osaka Univ.)

概要 We consider the nonlinear Schrödinger equations on the star graph with the Kirchhoff boundary and the repulsive Dirac delta boundary at the origin. In this talk, we show scattering-blowup dichotomy result below the mass-energy of the ground state on the real line. The proof of the scattering part is based on a concentration compactness and rigidity argument. Our main contribution is to give a linear profile decomposition on the star graph by using a new idea of decomposition of functions on the star graph.

- 41 川上隼平 (京大理) Averaging of strong magnetic nonlinear Schrödinger equations in energy space 12
 Jumpei Kawakami (Kyoto Univ.) Averaging of strong magnetic nonlinear Schrödinger equations in energy space

概要 We consider nonlinear Schrödinger equation with strong magnetic field in 3-dimension. To analyze this, R. L. Frank, F. Méhats, C. Sparber derived two nonlinear Schrödinger type models. One model is derived by spatial scaling and the other model is obtained by averaging the aforesaid spatial scaled model in time. We study these two models in energy space and obtain global solutions for some cases. We also prove that the time averaged model is energy-critical when this model has the nonic nonlinear power. Furthermore, we improve the previous result that the solution to the spatial scaled model converges to the solution to the time averaged model.

14:15~16:15

- 42 石塚健二郎 (京大数理研) ポテンシャルのついた非線形消散クライン・ゴールドン方程式の漸近挙動について 12
 Kenjiro Ishizuka (Kyoto Univ.) Long-time asymptotics of the one-dimensional damped nonlinear Klein–Gordon equation with a potential

概要 In this talk, we consider the one-dimensional nonlinear focusing damped Klein–Gordon equation with a potential. When a potential is good condition, it follows from standard arguments that the Cauchy problem for this equation is locally well-posed in the energy space. When a potential is zero, Cote–Martel–Yuan (2021) proved the full soliton resolution. Anyway we consider soliton resolution for this equation. Especially we will mainly talk the case that a potential is a repulsive delta potential.

- 43 眞崎 聡 (阪大基礎工) 3次非線形方程式系の分類について 12
 瀬片 純市 (九大数理)
 瓜屋 航太 (岡山理大理)
 Satoshi Masaki (Osaka Univ.) On classification of cubic nonlinear systems
 Jun-ichi Segata (Kyushu Univ.)
 Kota Uriya (Okayama Univ. of Sci.)

概要 In this talk, we introduce the classification of cubic nonlinear systems. The critical ingredient is to introduce a matrix representation of a system. It turns out that the rank of the matrix is an invariant quantity that describes the number of linearly independent conserved quantities. We give a classification of systems that have two distinct conserved quantities.

- 44 眞崎 聡 (阪大基礎工) 複素数値3次非線形方程式系の分類について 12
 Satoshi Masaki (Osaka Univ.) On classification of complex-valued cubic nonlinear systems

概要 In this talk, we introduce the classification of complex-valued gauge-invariant cubic systems. The critical ingredient is the matrix-vector representation of a system. This clarifies the effect of the change of variables and describes the systems' Hamiltonian structure. In particular, the rank of the matrix part denotes the number of the conserved quantities, and hence reflects the system's complexity. We show the classification of systems that have only one conserved quantity.

- 45 津田谷公利 (弘前大理工) 若杉勇太 (広島大先進理工) Blow up of solutions of semilinear wave equations with time-dependent damping 12
Kimitoshi Tsutaya (Hirosaki Univ.) Blow up of solutions of semilinear wave equations with time-dependent
Yuta Wakasugi (Hiroshima Univ.) damping

概要 We consider the Cauchy problem for nonlinear wave equations with time-dependent damping. We show that blow-up in a finite time occurs for the equation with arbitrary power nonlinearity as well as upper bounds of the lifespan of blow-up solutions. We also show the same results for the space derivative nonlinear term.

- 46 福田一貴 (信州大工) 平山浩之 (宮崎大教育) 一般化 KP–Burgers 方程式の初期値問題の解の長時間挙動と最適な減衰評価について 10
Ikki Fukuda (Shinshu Univ.) Large time behavior and optimal decay estimate for solutions to the
Hiroyuki Hirayama (Univ. of Miyazaki) Cauchy problem for the generalized KP–Burgers equation in 2D

概要 In this talk, we consider the Cauchy problem for the generalized KP–Burgers equation in 2D. This is one of the nonlinear dispersive-dissipative type equations, which has a spatial anisotropic dissipative term. Under some suitable regularity assumptions on the initial data u_0 , especially the condition $\partial_x^{-1}u_0 \in L^1(\mathbb{R}^2)$, it is known that the solution to this problem decays at the rate of $t^{-\frac{7}{4}}$ in the L^∞ -sense. In this study, we investigate the more detailed large time behavior of the solution and construct the approximate formula for the solution at $t \rightarrow \infty$. Moreover, we obtain a lower bound of the L^∞ -norm of the solution and prove that the decay rate $t^{-\frac{7}{4}}$ of the solution given in the previous work to be optimal.

- 47 S. Ibrahim (Univ. of Victoria) Phase transition threshold and stability of magnetic skyrmions 12
清水一慶 (阪大基礎工)
Slim Ibrahim (Univ. of Victoria) Phase transition threshold and stability of magnetic skyrmions
Ikkei Shimizu (Osaka Univ.)

概要 We examine the stability of vortex-like configuration of magnetization in magnetic materials, so-called the magnetic skyrmion. These correspond to critical points of the Landau–Lifshitz energy with the Dzyaloshinskii–Moriya interactions. In this talk, we prove that there is an explicit critical value of the parameter in the energy above which the skyrmion is unstable, while stable below this threshold. This mathematically explains the occurrence of phase transition observed in some experiments. We also show that in the unstable regime, the infimum of energy is not bounded below.

- 48 大縄将史 (東京海洋大海洋環境) 鈴木政尋 (名工大工) 周期境界条件下の浅水波系について 12
Masashi Ohnawa (Tokyo Univ. of Marine Sci. and Tech.) On the shallow water systems under periodic boundary conditions
Masahiro Suzuki (Nagoya Inst. of Tech.)

概要 We consider stationary flows over periodic topography modeled by one-dimensional shallow water equation. We first show the nonexistence of stationary solutions with discontinuities as well as the conditions on the existence of continuous subsonic/supersonic stationary solutions. We also claim that the subsonic stationary solutions are not asymptotically stable.

- 49 古屋 貴士 Inverse medium scattering problems with Kalman filter techniques ... 10
(島根大数理・データサイエンス教育研究センター)

Takashi Furuya (Shimane Univ.) Inverse medium scattering problems with Kalman filter techniques

概要 We study the inverse medium scattering problem to reconstruct the unknown inhomogeneous medium from the far-field patterns of scattered waves. The inverse scattering problem is generally ill-posed and nonlinear, and the iterative optimization method is often adapted. A natural iterative approach to this problem is to place all available measurements and mappings into one long vector and mapping, respectively, and to iteratively solve the linearized large system equation. However, this is computationally expensive because we must construct the larger system equations when the number of available measurements increases. In this talk, we propose the reconstruction algorithm based on the Kalman filter, which avoids the construction of a large system equation and retains the information of past updates.

16:30~17:30 特別講演

北 直 泰 (熊本大工) 複素係数をもつ非線形シュレーディンガー方程式の解の挙動

Naoyasu Kita (Kumamoto Univ.) On behaviors of solutions to nonlinear Schrödinger equations with complex coefficient in its nonlinearity

概要 We consider the Cauchy problem of nonlinear Schrödinger equations. Unlike the conservative system, the equation contains a complex coefficient in its nonlinearity. The physical background of such a non-conservative system is coming from the optical fiber engineering, where nonlinear Ohm's law and nonlinear amplification are applied. My talk consists of two parts. The first part treats a problem of global existence and asymptotic behavior of solutions without size restriction on the initial data, where Hayashi–Naumkin's theory basically contributes to the result. The second part treats a construction of blowing-up solutions. Combining the former result on asymptotic behavior of solutions, we can obtain blowing-up solutions even for small initial data. Recently, the result on the blowing-up solutions has been generalized by Cazenave–Martel–Zhao, Cazenave–Han–Martel and Kawakami–Machihara. If time permits, these recent progresses will be also presented.

3月18日(土) 第IV会場

9:30~12:00

- 50 岩田 直之 (名工大工) 多成分プラズマの Vlasov–Poisson 方程式の定常解 12
鈴木 政尋 (名工大工)

Naoyuki Iwata (Nagoya Inst. of Tech.) Stationary solutions of the Vlasov–Poisson system for a multicomponent
Masahiro Suzuki (Nagoya Inst. of Tech.) plasma

概要 In this talk, we mathematically investigate the formation of a plasma sheath, and analyze the Bohm criterion which is required for the formation. In 1995, Riemann derived the Bohm criterion from the stationary Vlasov–Poisson system which describes the motion of a multicomponent plasma. We study the solvability of boundary value problems of the system, and show that the Bohm criterion is a necessary condition but not a sufficient condition for the solvability.

- 51 鈴木政尋 (名工大) Vlasov–Poisson 方程式の定常解の安定性・不安定性 12
 高山正宏 (慶大理工)
 Katherine Zhiyuan Zhang
 (New York Univ.)
 Masahiro Suzuki (Nagoya Inst. of Tech.) Stability and instability of stationary solutions of the Vlasov–Poisson
 Masahiro Takayama (Keio Univ.) system
 Katherine Zhiyuan Zhang
 (New York Univ.)

概要 We investigate the formation of a plasma boundary layer (sheath) by considering the Vlasov–Poisson system on a half-line with the completely absorbing boundary condition. The solvability of the stationary problem has been established. In this talk, we study the nonlinear stability and instability of these stationary solutions of the Vlasov–Poisson system.

- 52 江口太一 (早大理工) Energy equality of MHD system under a weaker condition on magnetic
 field 12
 Taichi Eguchi (Waseda Univ.) Energy equality of MHD system under a weaker condition on magnetic
 field

概要 We prove the energy equality of MHD system in the space founded by Cheskidov–Constantin Friedlander–Shvydkoy (2008) and Berselli–Chiodaroli (2020). It is clarified that the energy equality is established for a larger class of the magnetic field than that of velocity field. Our result covers most of previous theorems on validity of the energy equality on the Navier–Stokes equations.

- 53 中里亮介 (早大理工) Fourier–Herz 空間上でのエネルギー法と Hall 効果を伴う準線形プラズマ
 モデルの漸近安定性への応用 12
 Ryosuke Nakasato (Waseda Univ.) Asymptotic stability for quasi-linear plasma models with Hall effects via
 energy methods on Fourier–Herz spaces

概要 We consider the initial-value problem for the compressible Hall-magnetohydrodynamic system with quantum effects in the 3-dimensional Euclidean space \mathbb{R}^3 . The aim of this talk is to obtain time-decay estimates for solutions as a perturbation from a constant equilibrium state $(1, 0, \bar{B})$ in a critical L^p framework. Here $\bar{B} \in \mathbb{R}^3$ is denoted by constant magnetic fields. To obtain such a result, we establish energy methods on Fourier–Herz spaces \hat{L}^p introduced by Grünrock (2004). By combining \hat{L}^p energy methods and a transformation of the momentum equation given by the effective velocity, we will derive the \hat{L}^p - $\hat{L}^{p/2}$ type time-decay estimate without the smallness requirement of low frequencies for initial data.

- 54 波多間備 (京大数理研) Stability of Fermi gas at zero temperature for the Hartree equation ... 12
 Sonae Hadama (Kyoto Univ.) Stability of Fermi gas at zero temperature for the Hartree equation

概要 We consider the Hartree equation describing the time evolution of wave functions of infinitely many fermions interacting with each other. The Hartree equation can be formulated in terms of the operators on $L^2(\mathbb{R}^d)$. It has infinitely many stationary solutions, and several authors have studied their stability, for example, Lewin–Sabin(2014), Chen–Hong–Pavloić(2018), and Collot–de Suzzoni(2020, 2022). However, the stability of Fermi gas at zero temperature, one of the most important stationary solutions from the physics point of view, was left open. In this talk, we report that it is stable; that is, any small perturbation from Fermi gas at zero temperature scatters.

- 55 古場 一 (阪大基礎工) 界面流を含む非粘性混相流の数値モデリング 10
 Hajime Koba (Osaka Univ.) Mathematical modeling of inviscid multiphase flow system with surface flow

概要 We consider the governing equations for the motion of the inviscid fluids in moving domains and surface from an energetic point of view. We employ an energetic variational approach to derive multiphase flow system with surface flow and tension. We make use of the feature of the barotropic fluid to derive the surface pressure and tension of the compressible fluid in the surface.

- 56 古場 一 (阪大基礎工) 相転移を含む非粘性混相流の数値モデリング 10
 Hajime Koba (Osaka Univ.) Mathematical modeling of inviscid multiphase flow system with phase transition

概要 We consider the governing equations for the motion of the fluid particles near air-water interface from an energetic point of view. We employ an energetic variational approach to derive inviscid multiphase flow system with phase transition, where a phase transition means a jump condition. We apply our energetic variational approach and jump condition to study the dominant equations for the densities of the fluids in moving domains.

- 57 古川 賢 (理化学研) 最大正則性のクラスでのプリミティブ方程式のデータ同化 10
 Ken Furukawa (RIKEN) Data assimilation of the primitive equations in maximal L^p - L^q settings

概要 We discuss about data assimilation to the primitive equations in maximal L^p - L^q settings in a mathematically rigorous way. The primitive equations describe the flow in a thin domain. The equations combined with some data assimilation methods are used in meteorology and ocean physics for predictions for phenomena. Our results are that the solution to data assimilation equations of the primitive equations converges to the solution to the primitive equations in maximal L^p - L^q spaces at $t \rightarrow \infty$.

- 58 野田 慶 (京大人間環境) 最大正則性に基づく多重線形作用素をもつ方程式の解の時空解析性について 10
 Kei Noda (Kyoto Univ.) Analyticity in space-time of solutions to evolution equations with multilinear operator based on maximal regularity

概要 In this presentation, we introduce a method to prove space-time analyticity for evolution equations with nonlinear terms represented by multilinear operators via the parameter trick. By extending the case of bilinear operators in the previous work, this method is expected to be applicable not only to the Navier–Stokes equations but also to the equations of Fujita type.

- 59 飯田 祥樹 (早大理工) Energy equality for the 3D inhomogeneous Navier–Stokes equations in Lorentz–Besov spaces 12
 Yoshiki Iida (Waseda Univ.) Energy equality for the 3D inhomogeneous Navier–Stokes equations in Lorentz–Besov spaces

概要 We establish the energy equality for Leray–Hopf weak solutions of the 3D incompressible *inhomogeneous* Navier–Stokes equations in Lorentz–Besov spaces. This result may be regarded as an extension of that by Cheskidov and Luo (2020).

14:15~16:15

- 60 青木基記 (東北大理) On the ill-posedness for the full system of the compressible Navier–
岩瀨司 (東北大理) Stokes equations 12
Motofumi Aoki (Tohoku Univ.) On the ill-posedness for the full system of the compressible Navier–
Tsukasa Iwabuchi (Tohoku Univ.) Stokes equations

概要 We consider the Cauchy problem for the equations of the ideal gas in the three space dimensions. In the critical Besov spaces, it is known that the Cauchy problem is uniquely solvable for all $1 < p < 3$ and ill-posed for all $p > 3$. We study the threshold case $p = 3$. We prove that the Cauchy problem is ill-posed by constructing a sequence of the initial data such that the solution map is discontinuous.

- 61 オーヘーダヒ (東北大理) Optimality of the decay estimate of solutions to the linearised curl-free
岩瀨司 (東北大理) compressible Navier–Stokes equations 12
Dáithí Ó hAodha (Tohoku Univ.) Optimality of the decay estimate of solutions to the linearised curl-free
Tsukasa Iwabuchi (Tohoku Univ.) compressible Navier–Stokes equations

概要 We discuss optimal estimates of solutions to the compressible Navier–Stokes equations in Besov norms. In particular, we consider the estimate of the curl-free part of the solution to the linearised equations, in the homogeneous case. We prove that our estimate is optimal in the L^∞ -norm by showing that the norm is bounded from below by the same decay rate. This is the first time such a bound from below has been obtained for the compressible Navier–Stokes equation.

- 62 岡部考宏 (阪大基礎工) Forced rapidly dissipative Navier–Stokes flows 12
L. Brandolese (Univ. Lyon 1)
Takahiro Okabe (Osaka Univ.) Forced rapidly dissipative Navier–Stokes flows
Lorenzo Brandolese (Univ. Lyon 1)

概要 We consider the asymptotic behavior of the solution to the incompressible Navier–Stokes equation on the whole space. By an action of a forcing term we control the nonlinear term of the flow and derive a rapid time decay. This talk is based on the joint work with Lorenzo Brandolese (Univ. Lyon 1).

- 63 中塚智之 (松山大経済) On solvability of the time-periodic problem for the Navier–Stokes equation 12
Tomoyuki Nakatsuka On solvability of the time-periodic problem for the Navier–Stokes equation
(Matsuyama Univ.) tion

概要 In this talk, we consider the existence of time-periodic solutions to the Navier–Stokes equation in the whole space. We decompose periodic solutions into steady and purely periodic parts, and we analyze the equations they should satisfy. Based on the analysis of the purely periodic solutions represented by the Fourier transform to the Stokes equation, their additional property can be obtained and we use it to construct a time-periodic solution of the Navier–Stokes equation.

- 64 小泉祐太 (早大理工) Convergence of approximating solutions of the Navier–Stokes equations
谷口透也 (早大理工) 12
Yuta Koizumi (Waseda Univ.) Convergence of approximating solutions of the Navier–Stokes equations
Toya Taniguchi (Waseda Univ.)

概要 We show that the approximating solutions $\{u_j\}_{j=0}^\infty$ of the Navier–Stokes equations constructed by Kato (1984) with the initial data $u(0) \in L_\sigma^n(\mathbb{R}^n)$ converge to the local strong solution u in the topology of $W^{2,n}(\mathbb{R}^n)$ provided the convergence in the scaling invariant norm in $L^p(\mathbb{R}^n)$ with the time weight holds.

- 65 大山 広樹 (九大数理) Fast rotation limit for the magnetohydrodynamics equations in a 3D layer 10
米田 慧司 (九大数理) layer 10
Hiroki Ohyama (Kyushu Univ.) Fast rotation limit for the magnetohydrodynamics equations in a 3D layer
Keiji Yoneda (Kyushu Univ.) layer

概要 We consider the initial value problem for the incompressible magnetohydrodynamics system with the Coriolis force in a three-dimensional infinite layer. We prove the unique existence of global solutions for initial data in the scaling invariant space when the speed of rotation is sufficiently high. Moreover, we show that the global solution converges to that of the coupled system of the 2D incompressible magnetohydrodynamics equations and the 3D incompressible Maxwell equations as the rotation speed tends to infinity.

- 66 江頭 貴成 (九大数理) Large time behavior of solutions to the 3D rotating Navier–Stokes equations 10
高田 了 (東大数理) tions 10
Takanari Egashira (Kyushu Univ.) Large time behavior of solutions to the 3D rotating Navier–Stokes equations
Ryo Takada (Univ. of Tokyo) tions

概要 We consider the large time behavior of the solutions for the initial value problem of the Navier–Stokes equations with the Coriolis force in the three-dimensional whole space. We show the temporal decay estimates with the dispersion effect of the Coriolis force for the global solutions. Moreover, we prove the large time asymptotic expansion of the solutions behaving like the first-order spatial derivatives of the integral kernel of the corresponding linear solution.

- 67 藤井 幹大 (九大数理) Global strong solutions to the compressible Navier–Stokes equation with the Coriolis force 12
Mikihiro Fujii (Kyushu Univ.) Global strong solutions to the compressible Navier–Stokes equation with the Coriolis force

概要 In this talk, we consider the global solvability for the compressible Navier–Stokes equation with the Coriolis force in the scaling critical Besov space. Despite many studies on the weak solutions, the existence of global strong solutions has been an open problem. Unlike the incompressible case, the Coriolis force causes the linear solutions to behave like a semigroup $\{e^{-t\Delta^2}\}_{t>0}$, which creates difficulties in nonlinear estimates in the low frequency part. In this research, we propose a new idea of the low frequency analysis and overcome these difficulties.

- 68 齋藤 平和 (電通大情報理工) On decay properties of solutions to the two-phase Stokes equations with surface tension and gravity 10
Hirokazu Saito On decay properties of solutions to the two-phase Stokes equations with surface tension and gravity
(Univ. of Electro-Comm.)

概要 In this talk, we consider the two-phase Stokes equations with surface tension in $\dot{\mathbf{R}}^N = \mathbf{R}_+^N \cup \mathbf{R}_-^N$ and in the presence of a uniform gravitational field acting vertically downward. Let ρ_{\pm} be positive constants describing the densities of fluids occupying \mathbf{R}_{\pm}^N . It is well-known that the Rayleigh–Taylor instability occurs when the upper fluid is heavier than the lower one, i.e., when $\rho_+ > \rho_-$. On the other hand, this talk aims to introduce time decay estimates of solutions to the two-phase Stokes equations under the condition that the lower fluid is heavier than the upper one, i.e., $\rho_- > \rho_+$.

16:30~17:30 特別講演

筒井容平(京大 理) **Convergence to the initial data and weighted estimates for the incompressible Navier–Stokes equations**

Youhei Tsutsui (Kyoto Univ.) **Convergence to the initial data and weighted estimates for the incompressible Navier–Stokes equations**

概要 We discuss conditions on data for which ensure the solution to the incompressible Navier–Stokes equations strongly converges to the initial data from weak L^n space on the whole space. We also consider weighted estimates for the solutions.

実 函 数 論

3月17日(金) 第V会場

9:00~11:45

- 1 豊田昌史(東邦大理) ボール空間における不動点定理と Caristi の不動点定理 15
Masashi Toyoda (Toho Univ.) Fixed point theorem in a ball space and Caristi fixed point theorem

概要 The purpose of this talk is to introduce a fixed point theorem in a ball space, which is proved without the Axiom of Choice. Moreover, using the fixed point theorem in a ball space, we prove Caristi's fixed point theorem. Therefore the proof of Caristi's fixed point theorem in this paper does not depend on the Axiom of Choice.

- 2 厚芝幸子(東京女大現代教養) Convergence theorems for families of monotone nonexpansive mappings
in ordered Banach spaces 15
Sachiko Atsushiba Convergence theorems for families of monotone nonexpansive mappings
(Tokyo Woman's Christian Univ.) in ordered Banach spaces

概要 In this talk, we prove weak and strong convergence theorems for monotone nonexpansive mappings and convergence theorems for monotone nonexpansive semigroups in Banach spaces endowed with a partial order.

- 3 木村泰紀(東邦大理) 収縮射影法の類似手法による写像族の共通不動点近似 15
Yasunori Kimura (Toho Univ.) Approximation of common fixed points by modified shrinking projection
methods

概要 This study deals with the approximation of fixed points for nonexpansive and other types of mappings. Using a modified shrinking projection method, we obtain the delta-convergence of generated approximation sequence to a fixed point of the mapping. We also consider a family of mappings and show a convergence theorem to their common fixed point.

- 4 青山耕治(千葉大社会) 吸引点に関する平均収束定理 15
Koji Aoyama (Chiba Univ.) Mean convergence theorems with respect to attractive points in a Hilbert
space

概要 We show a mean convergence theorem for a mapping with an attractive point in a Hilbert space by using a quasinonexpansive extension of the mapping and a mean convergence theorem for a quasinonexpansive mapping. Then using the theorem, we establish a mean convergence theorem for a λ -hybrid mapping in the sense of Aoyama, Iemoto, Kohosaka, and Takahashi (2010).

- 5 波多野修也(中大理工) モレー・ローレンツ空間に対する真の埋め込み 15
Naoya Hatano (Chuo Univ.) Proper embedding for Morrey-Lorentz spaces

概要 The embeddings for Morrey-Lorentz spaces have been previously introduced by Ragusa. This paper describes the major differences between these embedding types.

- 6 D. I. Hakim (Bandung Inst. of Tech.) Bourgain–Morrey spaces and its applications 15
 波多野修也 (中大理工)
 野ヶ山徹 (中大理工)
 澤野嘉宏 (中大理工)
 Denny Iwanal Hakim (Bandung Inst. of Tech.) Bourgain–Morrey spaces and its applications
 Naoya Hatano (Chuo Univ.)
 Toru Nogayama (Chuo Univ.)
 Yoshihiro Sawano (Chuo Univ.)

概要 In this talk, we introduce the Bourgain–Morrey space which is a one generalization of Morrey spaces. By using this space, it is showed that we can refine the Strichartz estimate for Schrödinger equations. We consider some properties of this space from the viewpoints of harmonic analysis and functional analysis.

- 7 野ヶ山徹 (中大理工) Complex interpolation for mixed Morrey spaces 15
 Toru Nogayama (Chuo Univ.) Complex interpolation for mixed Morrey spaces

概要 In this talk, we consider the complex interpolation for mixed Morrey spaces via the Calderón product. The mixed Morrey space $\mathcal{M}_q^p(\mathbb{R}^n)$ is a function space which is a generalization of Morrey spaces. The results in this talk are extensions for classical Morrey spaces.

- 8 谷口晃一 (東北大 AIMR) Besov 空間上の合成作用素の有界性 15
 池田正弘 (理化学研 AIP・慶大理工)
 石川勲 (愛媛大データサイエンスセンター)
 Koichi Taniguchi (Tohoku Univ.) Boundedness of composition operators on Besov spaces
 Masahiro Ikeda (RIKEN/Keio Univ.)
 Isao Ishikawa (Ehime Univ.)

概要 In this talk we consider boundedness of composition operators on Besov spaces $B_{p,q}^s$ on one dimensional Euclidean space. The boundedness has been well studied in the lower derivative case $0 < s < 1$. In contrast, there is almost no research on the higher order case $s \geq 1$, and many problems remain unsolved. The purpose of this talk is to give necessary and sufficient conditions for the boundedness when $s > 1 + 1/p$. Similar results for composition operators on Triebel–Lizorkin spaces are provided.

- 9 原 宇信 (北大情報) Strong barriers for weighted quasilinear equations 15
 Takanobu Hara (Hokkaido Univ.) Strong barriers for weighted quasilinear equations

概要 We construct strong barriers for weighted quasilinear elliptic equations. There are two applications: (i) solvability of Poisson-type equations with boundary singular data, and (ii) a geometric version of Hardy inequality.

14:15~15:25

- 10 川崎敏治 拡張積分可能関数族と拡張原始関数族の差について 15
 (玉川大工・千葉大法経・千葉大理)
 Toshiharu Kawasaki On the difference between the families of extended integrable functions
 (Tamagawa Univ./Chiba Univ./Chiba Univ.) and extended primitive functions

概要 In this talk, we discuss the difference between the families of extended integrable functions and extended primitive functions.

- 11 国定亮一 On topologically invariant means and almost convergence 15
Ryoichi Kunisada On topologically invariant means and almost convergence

概要 In this talk, we discuss applications of topologically invariant means and almost convergence for the theory of analytic functions on the half plane.

- 12 寺江拓海 (東工大情報理工) 非加法的測度論における歪測度の上限増分の一一致条件 15
室伏俊明 (東工大情報理工)
Takumi Terae (Tokyo Tech) An equality condition for supremum increments of distorted measures
Toshiaki Murofushi (Tokyo Tech) in non-additive measure theory

概要 In non-additive measure theory, there exist two definitions of null sets, which are weak null sets and strong null sets. Several theorems in measure theory such as the Egoroff theorem and the Lebesgue theorem are described by using the concept of null sets, therefore there exist weak forms and strong forms of these theorems as definitions in non-additive measure theory. The concept of supremum increment is important for studying about the strong forms of them. This paper has an argument that gives a necessary and sufficient condition for two different distorted measures having an equal supremum increment for a simple case. In addition, we define the concept of strong supremum increments in preparation for generalizing this argument.

- 13 岡崎悦明 (ファジィシステム研) 非加法的測度の定める集合環上の Fréchet–Nikodym 一様位相 15
福田亮治 (大分大理工)
本田あおい (九工大情報工)
Yoshiaki Okazaki Fréchet–Nikodym uniformity on a ring determined by a non-additive
(Fuzzy Logic Systems Inst.) measure
Ryoji Fukuda (Oita Univ.)
Aoi Honda (Kyushu Inst. of Tech.)

概要 Fréchet–Nikodym difference on a measure ring \mathcal{R} determined by a non-additive measure μ is defined by $\mu(A \ominus B)$. We prove that if μ is quasi-monotone and satisfies (p.g.p)-condition, then the entourages $\{(A, B) \mid \mu(A \ominus B) \leq \epsilon\}$, $\epsilon > 0$ is a basis of entourages of a uniform structure on the measure ring \mathcal{R} .

15:40～16:40 2022年度(第21回)日本数学会解析学賞受賞特別講演

- 中井英一 (茨城大理) 平均振動量が一樣でないコンパクト空間とそれに関連する話題
Eiichi Nakai (Ibaraki Univ.) Generalized Campanato spaces with variable growth condition and related topics

概要 Generalized Campanato spaces with variable growth condition were introduced by Nakai and Yabuta in 1985 to characterize the space of pointwise multipliers on $BMO(\mathbb{R}^n)$. This characterization was extended from \mathbb{R}^n to spaces of homogeneous type in 1997, to probability spaces with filtration in 2014, and, to \mathbb{R}^n with non-doubling measures in 2018. On the other hand, this characterization was used by Lerner in 2005 to study the class $\mathcal{P}(\mathbb{R}^n)$ of functions $p(\cdot)$ for which the Hardy-Littlewood maximal operator is bounded on the Lebesgue spaces $L^{p(\cdot)}(\mathbb{R}^n)$ with variable exponent, and positively solve a conjecture of Deining in 2005 saying that there are discontinuous functions belonging to $\mathcal{P}(\mathbb{R}^n)$. Moreover, in 2012 it was discovered by Nakai and Sawano that the dual of generalized Hardy space $H^{p(\cdot)}(\mathbb{R}^n)$ with variable exponent is a generalized Campanato space with variable growth condition. In this talk we discuss generalized Campanato spaces with variable growth condition and related topics.

17:00~18:00 特別講演

田中 仁 (筑波技術大) 2進立方体解析・2進直方体解析
 Hitoshi Tanaka Dyadic cubes analysis, dyadic rectangles analysis
 (Tsukuba Univ. of Tech.)

概要 After reviewing a history of the dyadic cubes analysis, with rectangular doubling weight, a generalized Hardy–Littlewood–Sobolev inequality for rectangular fractional integral operators is verified. The result is a nice application of M -linear embedding theorem for dyadic rectangles.

3月18日(土) 第V会場

9:00~12:00

- 14 赤木 剛朗 (東北大理) 非整数階時間微分を含む発展方程式の可解性とその応用 15
 中島 慶人 (東北大理)
 Goro Akagi (Tohoku Univ.) Solvability of evolution equations involving time-fractional derivatives
 Yoshihito Nakajima (Tohoku Univ.) and applications

概要 This talk is concerned with local and global in time solvability of abstract evolution equations involving time-fractional derivatives as well as subdifferential operators. An application to the initial-boundary value problem for degenerate parabolic equations involving time-fractional derivatives as well as blow-up nonlinearity will also be discussed.

- 15 喜多 航佑 (早大理工) Mosco convergence of functionals associated with Laplacian under non-
 大谷 光春 (早大理工) linear boundary conditions 15
 Kosuke Kita (Waseda Univ.) Mosco convergence of functionals associated with Laplacian under non-
 Ôtani Mitsuharu (Waseda Univ.) linear boundary conditions

概要 This talk is devoted to studying the convergence of functionals associated with Laplacian under nonlinear boundary conditions and the asymptotic behavior of solutions to nonlinear evolution equations governed by subdifferential operators of the functionals. To be more precise, we show the continuous dependence of the solution on the power of the nonlinear term of boundary conditions. The proof is based on some modifications of the argument of Mosco convergence by Attouch.

- 16 黒田隆徳 (早大理工) Asymptotic behaviors of complex solutions for the semilinear heat equation with arbitrarily large initial energy 15
 大谷光春 (早大理工)
 Takanori Kuroda (Waseda Univ.) Asymptotic behaviors of complex solutions for the semilinear heat equation with arbitrarily large initial energy
 Mitsuharu Ôtani (Waseda Univ.)

概要 In this talk, we consider the semilinear heat equation (P):

$$u_t - \Delta u - |u|^{q-2}u = 0, \quad u(0) = u_0$$

and the semilinear elliptic equation (E), which corresponds to the stationary problem of (P):

$$-\Delta u - |u|^{q-2}u = 0,$$

with the homogeneous Dirichlet boundary condition in bounded domains $\Omega \subset \mathbb{R}^N$, ($N \geq 3$) which are radially symmetric with respect to the first two variables. We seek solutions which takes complex values of the following form:

$$u_k(x_1, x_2, \tilde{x}) = e^{ik\theta} v_k(r, \tilde{x}), \quad (x_1, x_2, \tilde{x}) = (r \cos \theta, r \sin \theta, \tilde{x}) \in \Omega, \quad k \in \mathbb{N}.$$

We show the energy of solutions u_k of (E) tends to infinity as $k \rightarrow \infty$. Making use of this fact, for initial data of arbitrarily large energy, we construct two kinds of sets, i.e., stable set W and blow-up set V such that if $u_0 \in W$ (resp. $u_0 \in V$), then u can be continued globally up to $(0, \infty)$ (resp. u blows up in finite time).

- 17 小波津晶平 (東京理大理) Behavior of weak solutions to a Keller–Segel system with gradient dependent chemotactic coefficient 15
 横田智巳 (東京理大理)
 Shohei Kohatsu (Tokyo Univ. of Sci.) Behavior of weak solutions to a Keller–Segel system with gradient dependent chemotactic coefficient
 Tomomi Yokota (Tokyo Univ. of Sci.)

概要 We consider behavior of weak solutions to a Keller–Segel system with gradient dependent chemotactic coefficient:

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u |\nabla v|^{p-2} \nabla v), \\ v_t = \Delta v - v + u, \end{cases}$$

where $\chi > 0$, $p \in (1, \infty)$.

- 18 千代祐太朗 (東京理大理) Boundedness and stabilization in a quasilinear attraction-repulsion chemotaxis system 15
 Yutaro Chiyo (Tokyo Univ. of Sci.) Boundedness and stabilization in a quasilinear attraction-repulsion chemotaxis system

概要 This talk deals with a quasilinear parabolic–parabolic–elliptic attraction-repulsion chemotaxis system. In the case that the second equation is simplified by the elliptic one, boundedness and stabilization have already been obtained. The purpose of this talk is to derive boundedness and stabilization in the case that the second equation is parabolic.

- 19 T. Black (Paderborn Univ.) Possible points of blow-up in a chemotaxis system with environmental
M. Fuest dependent logistic source 15
(Leibniz Univ. Hannover)
J. Lankeit
(Leibniz Univ. Hannover)
水上雅昭 (京都教育大)
Tobias Black (Paderborn Univ.) Possible points of blow-up in a chemotaxis system with environmental
Mario Fuest (Leibniz Univ. Hannover) dependent logistic source
Johannes Lankeit
(Leibniz Univ. Hannover)
Masaaki Mizukami
(Kyoto Univ. of Edu.)

概要 We discuss the influence of possible spatial inhomogeneities in the coefficients of logistic source terms in parabolic-elliptic chemotaxis-growth systems in two-dimensional smoothly bounded domains. Assuming that the coefficient functions is nonnegative, we prove that finite-time blow-up of the classical solution can only occur in points where the coefficient function is zero.

- 20 赤川佳穂 (岐阜工高専) 仮似変分不等式で記述される塑弾性モデルについて 15
加納理成 (高知大教育)
深尾武史 (京都教育大)
Yoshiho Akagawa A quasi-variational inequality in plasticity model
(Gifu Nat. Coll. of Tech.)
Risei Kano (Kochi Univ.)
Takeshi Fukao (Kyoto Univ. of Edu.)

概要 This paper discusses the a quasi-variational inequality in plasticity model. In physical terms, it is natural that the threshold function depends on some unknown variable. This model is expected to describe plastic deformation due to the time-nonlocal dependence of the constraint on the unknown function. This is a difference from the perfect plasticity model. To prove the existence of a solution of the plasticity model, Banach fixed point theorem is applied, taking care to choose the test function.

- 21 森村晃子 (日本女大理) 多孔質媒体における水分移動モデルの解の存在と一意性について 15
愛木豊彦 (日本女大理)
Akiko Morimura On existence and uniqueness of solutions to the moisture transport
(Japan Women's Univ.) model in porous materials
Toyohiko Aiki (Japan Women's Univ.)

概要 We consider the initial-boundary value problems for nonlinear parabolic equations describing moisture transport in a porous material occupying a one-dimensional interval. Our problem is obtained by simplification of the model proposed by Green, Dabiri and Weinaug. Precisely, their system consists of two equations corresponding to conservation law for water in air and liquid regions, and in our problem the water distribution in air is given. The unknown function of the problem indicates the chemical potential of water. Also, we approximate the equation which is type of elliptic-parabolic, originally, as a first step in this research. The aim of this talk is to establish existence and uniqueness of solutions to the approximate problem by applying the standard fixed-point argument.

- 22 小杉千春 (日本女大理) 圧縮性弾性体の伸縮運動を表す初期値境界値問題の強解に対する ω 極限
愛木豊彦 (日本女大理) 集合について 15
Chiharu Kosugi (Japan Women's Univ.) Omega-limit sets for the strong solutions to initial and boundary value
Toyohiko Aiki (Japan Women's Univ.) problems for compressible elastic curves

概要 We talk about omega-limit sets for strong solutions to the initial and boundary value problem for the beam equation with the viscosity term and the nonlinear stress function having a singularity. Thanks to the singularity, we obtained the uniform estimate of lower bounds for the strain, and already proved existence and uniqueness of weak and strong solutions. By effect of the viscosity, it seems to be natural that the solution converges to the stationary solution. On this conjecture we show existence of a subsequence of the orbit $\{u(t)|t \geq 0\}$ such that it converges to a strong solution to the steady problem. We note that similar result on the weak solutions was reported at the previous meeting.

- 23 熊崎耕太 (長崎大教育) 多孔質媒体の湿潤・乾燥現象を表すマルチスケールモデルについて 15
A. Muntean (Karlstads Univ.)
Kota Kumazaki (Nagasaki Univ.) A multiscale model describing the wetting-drying phenomenon in porous
Adrian Muntean (Karlstads Univ.) materials

概要 In this talk, we propose a multiscale model describing the wetting-drying phenomenon in porous materials. This model consists of a diffusion equation for the relative humidity distributed in materials and a free boundary problem describing the wetting-drying process in microscopic pores. We consider each microscopic pore as a one-dimensional interval and correspond the interval to each point of materials. In our previous results, for a given relative humidity we showed the well-posedness of the free boundary problem. In this talk, we impose a governing equation for the relative humidity and discuss the existence and uniqueness of a locally-in-time solution of this model.

14:15~16:30

- 24 水野大樹 (千葉大教育) 擬放物型全変動流の適切性と解の正則性 15
白川 健 (千葉大教育)
Daiki Mizuno (Chiba Univ.) Well-posedness and regularity results for a class of total variation flows
Ken Shirakawa (Chiba Univ.) of pseudo-parabolic types

概要 In this talk, a class of initial-boundary value problems of total variation flows of pseudo-parabolic types is considered. Total variation flow has been adopted as governing equation of various nonlinear phenomena, such as grain boundary motion, image denoising, and so on. The principal characteristic of our problem is in the velocity term of diffusion flux, and the velocity would bring stronger regularity than that as in standard parabolic PDE. Meanwhile, our total variation flow contains singular diffusion, and the singularity would degenerate the regularity of solution. The objective of this paper will be to clarify the power balance between these conflicted effects. Consequently, the mathematical results concerned with well-posedness and regularity of solution will be discussed in the Main Theorems of this talk.

- 25 久保田翔大 (神奈川大工) 結晶粒界運動を記述する Warren-Kobayashi-Lobkovsky-Carter 型システムにおける時間周期解 15
Shodai Kubota (Kanagawa Univ.) Periodic solutions for Warren-Kobayashi-Lobkovsky-Carter type systems of grain boundary motions

概要 In this talk, we consider the periodic solutions for the parabolic PDE systems denoted by (S). Each system (S) is associated with the phase field model of grain boundary motion, proposed by [Warren, et al.; *Acta Materialia*, 51(20): 6035-6058, 2003]. Under suitable assumptions, the existence of the periodic solution to the system (S) will be reported as the main theorem of this talk.

- 26 渡邊 紘 (大分大理工) 3次元結晶粒界運動のフェーズフィールドモデルに対する解の存在 15
白川 健 (千葉大教育)
 S. Moll (Univ. València)
Hiroshi Watanabe (Oita Univ.) Existence of solutions to a phase-field model of 3D-grain boundary mo-
Ken Shirakawa (Chiba Univ.) tion
 Salvador Moll (Univ. València)

概要 We consider a phase-field model of 3D-grain boundary motion. The model is based on the three dimensional Kobayashi–Warren model for the dynamics of polycrystals. To formulate our 3D-model, we use a quaternion formulation for the orientation variable. In this talk, we obtain existence of solutions to the L^2 -gradient descent flow of the constrained energy functional via several approximating problems. Moreover, we also obtain an invariance principle for the orientation variable.

- 27 深尾 武史 (京都教育大) 前方後方動的境界条件を持つ Cahn–Hilliard 方程式に対する LW モデル
 P. Colli (Univ. of Pavia) からの接近 15
 L. Scarpa
 (Politecnico di Milano)
Takeshi Fukao (Kyoto Univ. of Edu.) The Cahn–Hilliard system of LW model with forward-backward dy-
 Pierluigi Colli (Univ. of Pavia) namic boundary condition
 Luca Scarpa (Politecnico di Milano)

概要 For Cahn–Hilliard equations with forward-backward dynamical boundary conditions, we consider the vanishing viscosity approach from LW model. In the previous study, the approach from GMS model was discussed. The same idea is used for the LW model. Since the regularity of the solution is lowered in the limit equations, we define a weak solution using the subdifferential with suitable function spaces. In the LW model unlike the GMS model, it is subject to the mass conservations at the bulk and the boundary, respectively, which makes some difficulty. In this talk, we will especially introduce the idea of the uniform estimate.

- 28 山崎 教昭 (神奈川大工) Approximation of singular optimal control problems for doubly quasi-
剣持 信幸 (千葉大*) variational evolution inclusions 15
白川 健 (千葉大教育)
Noriaki Yamazaki (Kanagawa Univ.) Approximation of singular optimal control problems for doubly quasi-
 Nobuyuki Kenmochi (Chiba Univ.*) variational evolution inclusions
 Ken Shirakawa (Chiba Univ.)

概要 In this talk, we establish an approximate procedure for singular optimal control problems for abstract doubly quasi-variational evolution inclusions governed by time-dependent subdifferentials with the unknown-dependent constraints. Indeed, we consider the parameter-dependent state inclusion and its control problems. Then, we show the relationship between the original control problem and its approximate one.

- 29 伊藤 昭夫 Tumor invasion model with quasi-variational structural porous medium diffusion 15
Akio Ito Tumor invasion model with quasi-variational structural porous medium diffusion

概要 We consider an initial-boundary value problem of a tumor invasion of Chaplain–Anderson type, in which the coefficient of random motility of tumor cells depends on the extracellular matrix and the diffusion flux of tumor cells is porous medium type whose power depends upon the extracellular matrix. From these points of view, the Cauchy problem considered in this talk has a quasi-variational structure and this fact makes it more difficult and complicate to analyze this model mathematically. The aim of this talk, we give the global existence of strong solutions by using the general theory, which was established in the paper, Evolution inclusion on a real Hilbert space with quasi-variational structure for inner products, Journal of Convex Analysis, 26 (2019), No. 4, 1185–1252.

- 30 松本 敏隆 (静岡大理) 一般化安定性条件の下での線形発展作用素の生成 15
田中 直樹 (静岡大理)
Toshitaka Matsumoto (Shizuoka Univ.) Generation of evolution operators under a generalized stability condition
Naoki Tanaka (Shizuoka Univ.)

概要 A generation theorem of a linear evolution operator on a Banach spaces is given under a generalized stability condition for a difference scheme. The denseness and the time-independence of the domains of generators are not assumed. A differentiability theorem of an evolution operator is also given.

17:00~18:00 特別講演

- 来間 俊介 (東京理大理) フェーズフィールドシステムに対する時間離散化法
Shunsuke Kurima (Tokyo Univ. of Sci.) Time discretization methods for phase field systems

概要 The speaker has studied approaches by time discretization to a phase field system of Caginalp type and a singular phase field system of conserved type with Pierluigi Colli. The time discretization method is a method of capturing the derivative with respect to time with the limit of the difference quotient, and it has the advantage that evolution equations can be reduced to stationary problems of elliptic equations. In this talk, for phase field systems with inertial term which has already been proved to have solutions, we will confirm if we can apply time discretization methods to show existence of solutions and derive error estimates for the time step. Also, in this talk, for singular nonlocal phase field systems with inertial term which has not yet been proved to have solutions, we will show existence of solutions by time discretization methods.

函数解析学

3月15日(水) 第VI会場

9:30~12:00

- 1 岩田 順 敬 (関西大化学生命工) 非有界作用素の対数表現によるフォン・ノイマン方程式の一般化 15
 Yoritaka Iwata (Kansai Univ.) Generalization of von Neumann equation based on the logarithmic representation of unbounded operators

概要 In this paper, by replacing the commutator product by the logarithmic representation of unbounded operators [1-9], the generalized/unbounded version of von Neumann equation is presented. The proposed equation is general enough to be valid to bounded situations (i.e., classical mechanics). By taking some concrete examples in the quantum mechanics, the derivation of generalized von Neumann equation is explained.

- 2 渡辺 秀 司 (群馬大理工) 超伝導のBCSギャップ方程式の新たな作用素論的扱いと2次相転移への応用 15
 Shuji Watanabe (Gunma Univ.) A new operator-theoretical treatment of the BCS gap equation of superconductivity and its application to the second-order phase transition

概要 We give a new operator-theoretical proof of the existence, uniqueness and smoothness of the solution to the BCS-Bogoliubov gap equation of superconductivity and apply the results to the second-order phase transition.

- 3 青木 貴 史 (近畿大*) 整関数の空間における連続準同型の微分作用素表示 15
 石村 隆 一 (千葉大*)
 岡田 靖 則 (千葉大理工)
 Takashi Aoki (Kindai Univ.*) Differential operator representations of continuous homomorphisms on
 Ryuichi Ishimura (Chiba Univ.*) some spaces of entire functions
 Yasunori Okada (Chiba Univ.)

概要 The space of entire functions at most of a given order and of normal type (resp. minimal type) is endowed with a locally convex topology as an inductive limit (resp. projective limit) of Banach spaces. Continuous endomorphisms of such spaces include differential operators, translations, dilations, etc., and they can be represented as formal differential operators of infinite order satisfying suitable conditions. We consider continuous homomorphisms between spaces of entire functions with growth given by proximate orders instead of orders, and give their differential operator representations. We also reveal the merits of treating proximate orders and homomorphisms from the view point of inductive and projective descriptions.

- 4 廣島 文 生 (九大数理) 並行移動不変なネルソン模型の正值改良性 15
 Fumio Hiroshima (Kyushu Univ.) Positivity improving for the translation invariant Nelson model

概要 Let $H(P)$ be the renormalized Nelson Hamiltonian with total momentum P . We show that $\exp(-TH(P))$ is positivity improving for any P .

- 5 佐々木格 (信州大理) Analyticity of the ground state for the pair interaction model 15
 松澤泰道 (信州大教育)
 泉真之介 (信州大理)
 井村航太 (長野県富士見高)
 Itaru Sasaki (Shinshu Univ.) Analyticity of the ground state for the pair interaction model
 Yasumichi Matsuzawa (Shinshu Univ.)
 Shinnosuke Izumi (Shinshu Univ.)
 Kota Imura
 (Nagano Pref. Fujimi High School)

概要 We consider the pair interaction model which is a models in the quantum field theory. In our previous work, we diagonalize its Hamiltonian by a Bogoliubov transformation. As a consequence, the existence and uniqueness of the ground states are proven. In addition, the explicit formula for the ground state energy has been given. In this talk, we show that the ground state and the ground state energy are analytic in the coupling constant. Their radii of convergence are always finite.

- 6 吉田尚矢 (立命館大理工) Bohr–Sommerfeld quantization condition for self-adjoint Dirac operators 15
 Naoya Yoshida (Ritsumeikan Univ.) Bohr–Sommerfeld quantization condition for self-adjoint Dirac operators

概要 We consider the eigenvalue problem for self-adjoint Dirac operators. It is known that, near an energy level where the square of the potential makes a simple well, the eigenvalues are approximated by a Bohr–Sommerfeld type quantization rule. A remarkable difference from the Schroedinger case appears in the Maslov correction term. This fact was recently found by Hirota (2017) under the analyticity condition of the potential using a complex WKB method. In this talk, we approach this problem with a microlocal technique focusing on the asymptotic behavior of the eigenfunction along the characteristic set to generalize the result to smooth potentials.

- 7 板倉恭平 (東大数理) On resonances for inverted harmonic oscillators 15
 Kyohei Itakura (Univ. of Tokyo) On resonances for inverted harmonic oscillators

概要 We characterize resonance free domain for semiclassical inverted harmonic oscillators by employing method of Briet–Combes–Duclos. For fixed non-zero energy E , we assume a certain non-trapping condition; the whole domain can be split into the classically forbidden region and region where virial is bounded by negative constant from above. Then we obtain the absence of resonances in a complex neighborhood of E .

- 8 宇治野広大 (九大数理) Exact Hausdorff dimension of the spectral measure for the graph Laplacian on a sparse tree 15
 Kota Ujino (Kyushu Univ.) Exact Hausdorff dimension of the spectral measure for the graph Laplacian on a sparse tree

概要 It is known that Schroedinger operators with sparse potentials have singular continuous spectrum. The graph Laplacian on a sparse tree is identified with one-dimensional discrete Schroedinger operators, and has singular continuous spectrum. The intermittency function describes the behavior of the time averaged momentum and gives the upper bound of the Hausdorff dimension of the spectral measure. We show the intermittency function explicitly and give the Hausdorff dimension of the spectral measure for the graph Laplacian on a sparse tree.

- 9 宇治野 広大 (九大数理) No eigenvectors embedded in the singular continuous spectrum of Schrödinger operators 15
 Kota Ujino (Kyushu Univ.) No eigenvectors embedded in the singular continuous spectrum of Schrödinger operators

概要 It is known that the spectrum of Schrödinger operators with sparse potentials consists of singular continuous spectrum. We give a sufficient condition so that the edge of the singular continuous spectrum is not an eigenvalue and construct examples with singular continuous spectrum which have no eigenvalues and which have a single negative eigenvalue.

14:15~16:15

- 10 柳下 浩紀 (京都産大理) 無限次元空間 \mathbb{R}^∞ 上の Laplace 作用素の変分法的定式化と自己共役性 .. 10
 Hiroki Yagisita (Kyoto Sangyo Univ.) Variational formulation and self-adjointness of Laplace operator in infinite dimensional space \mathbb{R}^∞

概要 In a variational method like Lax–Milgram theorem, we define Laplacian on the infinite direct product space $\mathbb{R}^\infty := \prod_{n \in \mathbb{N}} \mathbb{R}$ and show that it is a self-adjoint operator. Here, the most important thing is to define Hilbert space that corresponds to Sobolev space $H^1(\mathbb{R}^\infty)$. To do it, we use square root of density introduced by Hormander in context of Fourier integral operator theory. However, here, we use a measure-theoretic method rather than a geometric method to give a very elementary introduction at least for analyst (inspired by statistical interpretation of quantum mechanical wave function by Born and Heisenberg) of square root of density. All in all, only basic knowledge of analysis (as it is quite possible to study undergraduate) is required.

- 11 柳下 浩紀 (京都産大理) \mathbb{R}^∞ 上の無限次元干渉 Brown 運動に関する解析的半群 5
 Hiroki Yagisita (Kyoto Sangyo Univ.) Analytic semigroup of infinite dimensional interacting Brownian motion on \mathbb{R}^∞

概要 We define the generator of a holomorphic semigroup for a countably infinite interacting Brownian particle system on a countable direct product space $\mathbb{R}^\infty := \prod_{n \in \mathbb{N}} \mathbb{R}$ as a maximal monotone self-adjoint operator $\sum_{k=1}^{\infty} ((-\frac{\partial}{\partial x_k} + H_k)(\frac{\partial}{\partial x_k} + H_k))$. At first glance, this may seem like a pretty difficult problem, but it is not too difficult because $\frac{\partial}{\partial x_k}$ on \mathbb{R}^∞ can be defined as just a (generalized) partial derivative. The most important thing is to define a H -partial derivative $\partial_{H,k} := \frac{\partial}{\partial x_k} + H_k$ as a closed operator for a sequence $\{H_k\}_{k=1}^{\infty}$ of measurable functions whose domain are measurable “subsets” of \mathbb{R}^∞ . As with the previous talk, only basic knowledge of analysis (and part of the previous talk) is required.

- 12 望月 清 (中大理工・都立大*) 摩擦項をもつ磁場中の波動方程式に対する極限振幅の原理 15
 中澤 秀夫 (日本医大)
 Kiyoshi Mochizuki (Chuo Univ./Tokyo Metro. Univ.*) The principle of limiting amplitude for dissipative wave equations in magnetic fields
 Hideo Nakazawa (Nippon Medical School)

概要 The dissipative wave propagation problem with magnetic Laplacian in an exterior domain is considered. The uniform estimates and Hölder conditions of the resolvent are studied for the reduced wave operator without dissipation. Based on these results, the validity of the principle of limiting amplitude is proved for wave propagation problem with dissipation. As byproduct, the result by Mizohata–Mochizuki in 1966 are improved.

- 13 大屋博一 (サレジオ工高専) 指数関数を重みにもつ ソボレフ空間の埋め込みに関する一考察 15
 Hirokazu Ohya (Salesian Polytech.) Embedding properties for some Weighted Sobolev spaces with growing weight

概要 In this talk we are concerned with the embedding properties for several kind of functional spaces in R^N . We will consider the functional spaces called as Weighted Sobolev space with some weight function. If we choose some suitable growing functions in the weight functions, we can derive the continuity and compact embedding for these functional spaces. We also obtain the sufficient conditions for the weight functions to show the above properties. Our method is mainly owed to usual Sobolev's inequality.

- 14 樋口健太 (愛媛大理工) 複数の接触交差が生成する小さな擬交差間の 2 準位断熱遷移確率 15
 渡部拓也 (立命館大理工)
 Kenta Higuchi (Ehime Univ.) Two-level adiabatic transition probability for small avoided crossings
 Takuya Watanabe (Ritsumeikan Univ.) generated by tangential intersections

概要 We consider a first order ordinary differential system with two small parameters, which are an adiabatic parameter and an interaction one. This system is a model of a time-dependent Schrödinger equation which describes that two energy-levels with tangential intersections avoid crossings. In this talk, we give asymptotic expansion of the transition probability between avoided crossings as two parameters tend to zero as well as their ratio determined by the tangential order does.

- 15 宮西吉久 (信州大理工) 薄型領域におけるノイマン・ポアンカレ作用素のスペクトル 10
 Yoshihisa Miyanishi (Shinshu Univ.) Spectrum of the Neumann–Poincaré operator on thin domains

概要 The Neumann–Poincaré operator (abbreviated by NP) is a boundary integral linear operator known as one of the important tools associated with boundary value problems in the field of partial differential equations. Our purpose here is to introduce the spectral structure of the NP operator. In an asymptotic sense, the NP spectrum distributed in the interval $[-1/2, 1/2)$ as the domains become longer in 2D. In an asymptotic sense, we also find that the NP spectrum densely distributed in the interval $[-1/2, 1/2)$ (resp. $[0, 1/2)$), as the domains which are defined by spheroids become flatter (resp. longer) in 3D.

- 16 穴原義弘 (横浜国大先進実践) 流出入を伴う 1 次元格子量子ウォークの極限定理 15
 今野紀雄 (横浜国大工)
 森岡悠 (愛媛大理工)
 瀬川悦生 (横浜国大環境情報)
 Yoshihiro Anahara (Yokohama Nat. Univ.) Limit theorem for quantum walk on one dimensional lattice with inflows and outflows
 Norio Konno (Yokohama Nat. Univ.)
 Morioka Hisashi (Ehime Univ.)
 Etsuo Segawa (Yokohama Nat. Univ.)

概要 To obtain a stationary state of quantum walks on the finite path, we set the sink and source to the boundaries of the path so that the internal path receives the inflow and radiates the outflow at every time step. In this talk we consider one dimensional lattice as finite graph and we show the weak convergence theorem for the scaled limit distribution of the comfortability in the limit of the length of the path.

- 17 樋口 健太 (愛媛大理工) Resonance expansion for finitely perturbed free quantum walks on the
 森岡 悠 (愛媛大理工) line 15
 瀬川 悦生 (横浜国大環境情報)
 Kenta Higuchi (Ehime Univ.) Resonance expansion for finitely perturbed free quantum walks on the
 Hisashi Morioka (Ehime Univ.) line
 Etsuo Segawa (Yokohama Nat. Univ.)

概要 We introduce resonances to a class of quantum walks on the line as poles inside the unit circle of the outgoing resolvent. With each resonance, a resonant state which behaves like an eigenvector is associated. We show that the time evolution with an initial state with compact incoming support exactly coincides with the evolution of a sum of resonant states in each compact set after a finite length of time has passed. Consequently, the exponential decay rate of the time evolution is given by the modulus of resonances. We note that in the special case where a resonance is algebraically multiple, the decay becomes polynomially slower.

16:30~17:30 特別講演

- 高江洲 俊光 (群馬大理工) 相互作用する場の量子論の系のスペクトル解析について
 Toshimitsu Takaesu (Gunma Univ.) On the spectral analysis of interacting quantum field systems

概要 In this talk, interacting systems of quantum field models are investigated. We mainly consider the system of quantum electrodynamics, which is the interaction system of a Dirac field and the radiation field in the Coulomb gauge, but the Yukawa model and ϕ^4 model are also considered. By introducing momentum and spatial cutoffs, the Hamiltonians are self-adjoint operators on a Hilbert space. Focusing on ground states, the spectral analysis of these systems will be explained.

3月16日(木) 第VI会場

9:30~12:00

- 18 里見 貴志 (東大数理) ユニモジュラー局所コンパクト群上の畳み込みと再配分に関する不等式
 と関連する不等式的最適定数 15
 Takashi Satomi (Univ. of Tokyo) Inequality for the convolutions on unimodular locally compact groups
 and the optimal constant of related inequalities

概要 Let μ be the Haar measure of a unimodular locally compact group G and $m(G)$ be the infimum of the volumes of all open subgroups of G . The main result is that $\int_G f \circ (\phi_1 * \phi_2)(g) dg \leq \int_{\mathbb{R}} f \circ (\phi_1^* * \phi_2^*)(x) dx$ holds for any measurable functions $\phi_1, \phi_2: G \rightarrow \mathbb{R}_{\geq 0}$ with $\mu(\text{supp } \phi_1) + \mu(\text{supp } \phi_2) \leq m(G)$ and any convex function $f: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$ with $f(0) = 0$. Here ϕ^* is the rearrangement of ϕ .

Let $Y_O(P, G)$ and $Y_R(P, G)$ denote the optimal constants of Young's and the reverse Young's inequality, respectively, on G for the pair of the exponents $P = (p_1, p_2)$ under the assumption $\mu(\text{supp } \phi_1) + \mu(\text{supp } \phi_2) \leq m(G)$. Then we have $Y_O(P, G) \leq Y_O(P, \mathbb{R})$ and $Y_R(P, G) \geq Y_R(P, \mathbb{R})$ as a corollary of the main result. Thus, we obtain that $m(G) = \infty$ holds if and only if $H(p, G) \leq H(p, \mathbb{R})$ in the case of $p' := p/(p-1) \in 2\mathbb{Z}$, where $H(p, G)$ is the optimal constant of the Hausdorff-Young inequality on G for the exponent p .

- 19 久保利久 (龍谷大経済) On the standardness of homomorphisms between generalized Verma modules for $\mathfrak{sl}(3, \mathbb{C})$ 15
 Toshihisa Kubo (Ryukoku Univ.) On the standardness of homomorphisms between generalized Verma modules for $\mathfrak{sl}(3, \mathbb{C})$

概要 Let \mathfrak{g} be a finite dimensional complex simple Lie algebra and \mathfrak{b} a Borel subalgebra. Let $\mathfrak{p} \supset \mathfrak{b}$ be a standard parabolic subalgebra. In 1977, Lepowsky showed that a homomorphism $\varphi: M_{\mathfrak{b}}(\nu) \rightarrow M_{\mathfrak{b}}(\zeta)$ between Verma modules induces a map $\varphi_{\text{std}}: M_{\mathfrak{p}}(\nu) \rightarrow M_{\mathfrak{p}}(\zeta)$ between the corresponding generalized Verma modules, provided that ν and ζ satisfy a certain integral condition. He called such a map φ_{std} *standard*. It was also observed that the standard map φ_{std} could be zero and even in the case there could be a non-zero homomorphism $\varphi_{\text{nonstd}}: M_{\mathfrak{p}}(\nu) \rightarrow M_{\mathfrak{p}}(\zeta)$. Any map that is not standard is called *non-standard*. The classification of standard/non-standard maps is still an open problem.

In this talk, for $\mathfrak{g} = \mathfrak{sl}(3, \mathbb{C})$ with a maximal parabolic subalgebra \mathfrak{p} , we shall show that all non-zero homomorphisms $\psi: M_{\mathfrak{p}}(\nu) \rightarrow M_{\mathfrak{p}}(\zeta)$, where $M_{\mathfrak{p}}(\zeta)$ is of scalar type, are standard.

- 20 織田 寛 (拓殖大工) BC 型 Opdam–Cherednik 変換の逆変換公式 15
 Hiroshi Oda (Takushoku Univ.) Inversion formula for Opdam–Cherednik transform associated with a root system of type BC

概要 We give the inversion formula and the Plancherel formula for Opdam–Cherednik transform associated with a root system of type BC , when the multiplicity parameters are not necessarily nonnegative.

- 21 貝塚 公一 (日本医大) Some remarks on the Dirac operator on symmetric spaces 15
 Koichi Kaizuka Some remarks on the Dirac operator on symmetric spaces
 (Nippon Medical School)

概要 We develop the spectral analysis for the Dirac operator on symmetric spaces of noncompact type based on harmonic analysis on Lie groups. In this talk, we compute the continuous spectrum of the Dirac operator on irreducible symmetric spaces of noncompact type. We show that the continuous spectrum has a spectral gap if and only if the symmetric space is isomorphic to a coset space of the special pseudo-unitary group of odd matrix size. Furthermore, we give a uniform weighted resolvent estimate for the Dirac operator under a certain assumption on the symmetric space.

- 22 中濱 良祐 Computation of weighted Bergman inner products on block off-diagonal matrices in bounded symmetric domains for $Sp(2r, \mathbb{R})$ 15
 (NTT 基礎数学研究センター)
 Ryosuke Nakahama Computation of weighted Bergman inner products on block off-diagonal matrices in bounded symmetric domains for $Sp(2r, \mathbb{R})$
 (NTT Inst. for Funda. Math.)

概要 Let $(G, G_1) = (G, (G^\sigma)_0)$ be a symmetric pair of holomorphic type, and we consider a pair of Hermitian symmetric spaces $D_1 = G_1/K_1 \subset D = G/K$, realized as bounded symmetric domains in complex vector spaces $\mathfrak{p}_1^+ := (\mathfrak{p}^+)^\sigma \subset \mathfrak{p}^+$ respectively. Then the universal covering group \tilde{G} of G acts unitarily on the weighted Bergman space $\mathcal{H}_\lambda(D) \subset \mathcal{O}(D) = \mathcal{O}_\lambda(D)$ on D for sufficiently large λ . Its restriction to the subgroup \tilde{G}_1 decomposes discretely and multiplicity-freely, and its branching law is given in terms of the \tilde{K}_1 -decomposition of the space $\mathcal{P}(\mathfrak{p}_2^+)$ of polynomials on $\mathfrak{p}_2^+ := (\mathfrak{p}^+)^{-\sigma} \subset \mathfrak{p}^+$. Our goal is to understand the decomposition of the restriction $\mathcal{H}_\lambda(D)|_{\tilde{G}_1}$ by studying the weighted Bergman inner product on each \tilde{K}_1 -type in $\mathcal{P}(\mathfrak{p}_2^+) \subset \mathcal{H}_\lambda(D)$. Today we deal with the symmetric pair $(G, G_1) = (Sp(2r, \mathbb{R}), Sp(r, \mathbb{R}) \times Sp(r, \mathbb{R}))$.

- 23 田中雄一郎 (東大数理) 無重複性のユニタリトリックについて 15
 Yuichiro Tanaka (Univ. of Tokyo) A unitary trick for the multiplicity-freeness property

概要 With the aim of uniform treatment of multiplicity-free representations of Lie groups including infinite-dimensional representations, T. Kobayashi introduced the notion of visible actions on complex manifolds. In this talk I would like to show a trick for obtaining the multiplicity-freeness property of infinite-dimensional representations of real semisimple Lie groups from that of finite-dimensional ones of compact Lie groups through visible actions on flag varieties.

- 24 V. Pérez-Valdés (東大数理) 3次元球面上のベクトル束から2次元球面上の直線束への共形同変な微分対称性破れ作用素の構成と分類 15
 Víctor Pérez-Valdés (Univ. of Tokyo) Construction and classification of conformally equivariant differential symmetry breaking operators from a vector bundle over S^3 to a line bundle over S^2

概要 In this talk, we explain about the construction and classification problems of all differential symmetry breaking operators $D : C^\infty(S^3, \mathcal{V}_\lambda^{2N+1}) \rightarrow C^\infty(S^2, \mathcal{L}_{m,\nu})$ between the section spaces of $\mathcal{V}_\lambda^{2N+1} \rightarrow S^3$, a rank $2N+1$ vector bundle over S^3 , and $\mathcal{L}_{m,\nu} \rightarrow S^2$, a line bundle over S^2 .

- 25 中島秀斗 (統計数理研) 等質開凸錐に付随するゼータ関数のガンマ行列の分解定理 15
 Hideto Nakashima (Inst. of Stat. Math.) Decomposition formulas for gamma matrices of zeta functions associated with homogeneous cones

概要 Associated with homogeneous cones, we are able to construct solvable prehomogeneous vector spaces. In the previous paper, we calculate gamma matrices of the associated zeta functions explicitly. In this talk, we show that the gamma matrices can be decomposed into variable-wise matrices. Moreover, under suitable condition, we show that the associated zeta functions admit a kind of completion forms.

13:10~14:10 特別講演

橋本隆司 シンプレクティック・ベクトル空間上の運動量写像の量子化と極小表現
 (鳥取大教育支援・国際交流推進機構)

Takashi Hashimoto (Tottori Univ.) Quantization of the moment map on symplectic vector spaces and minimal representation

概要 Let G be $\mathrm{Sp}(n, \mathbb{R})$, $\mathrm{U}(p, q)$, or $\mathrm{O}^*(2n)$, which acts on a symplectic vector space W in a Hamiltonian way with moment map μ . The purpose of this talk is twofold: the first one is to show that the canonical quantization of μ with a choice of a Lagrangian subspace of W provides the oscillator representations of G , which have been extensively studied in relation to the Howe duality and to minimal representations. The second one is to construct the minimal representation of $\mathrm{O}(p, q)$ by applying this method to $\mathrm{O}(p, q)$ and using the fact that $(\mathrm{O}(p, q), \mathrm{SL}_2(\mathbb{R}))$ is a Howe's dual pair.

3月17日(金) 第VI会場

9:30~12:00

- 26 大井志穂 (新潟大理) 行列値リプシッツ環上の全射線形等距離写像の代数的反射性 15
 Shiho Oi (Niigata Univ.) Algebraic reflexivity of isometry groups of Lipschitz algebras

概要 We consider Banach algebras of Lipschitz maps taking the values in a Banach algebra of complex matrices of degree n . The unital subjective linear isometries on the Banach algebras have been obtained. In this talk, we study the algebraic reflexivity of the set of subjective linear isometries. The subset S of the set of all bounded linear operators from A_1 into A_2 is called algebraically reflexive if a bounded linear operator $T : A_1 \rightarrow A_2$ which satisfies $Ta \in Sa$ for every $a \in A_1$ implies $T \in S$. We show the set of unital subjective linear isometries is algebraic reflexivity.

- 27 榎並優太 (新潟大自然) Surjective isometries on the Banach algebra of analytic functions with
三浦毅 (新潟大理) C^n -boundary values 12
Yuta Enami (Niigata Univ.) Surjective isometries on the Banach algebra of analytic functions with
Takeshi Miura (Niigata Univ.) C^n -boundary values

概要 Let \mathbb{D} , $\bar{\mathbb{D}}$ and \mathbb{T} be the open unit disk, closed unit disk and unit circle in the complex plane \mathbb{C} . The disk algebra $A(\bar{\mathbb{D}})$ is the algebra of all continuous functions on $\bar{\mathbb{D}}$ that is holomorphic in \mathbb{D} . We consider the subalgebra $A^n(\bar{\mathbb{D}})$ consisting of all $f \in A(\bar{\mathbb{D}})$ whose restrictions $f|_{\mathbb{T}}$ are of class C^n . For each $f \in A^n(\bar{\mathbb{D}})$, we denote by $D^k(f)$ the k -th derivative of $f|_{\mathbb{T}}$. The algebra $A^n(\bar{\mathbb{D}})$ is a Banach algebra, with the norm $\|f\|_{\Sigma} = \sum_{k=0}^n \|D^k(f)\|_{\infty}/k!$, where $\|g\|_{\infty} = \sup_{z \in \mathbb{T}} |g(z)|$. We characterized surjective, not necessarily linear, isometries on $A^n(\bar{\mathbb{D}})$ with respect to the norm $\|\cdot\|_{\Sigma}$.

- 28 三浦毅 (新潟大理) Tingley's problem for uniformly closed function algebras 15
廣田大輔 (新潟大自然)
Takeshi Miura (Niigata Univ.) Tingley's problem for uniformly closed function algebras
Daisuke Hirota (Niigata Univ.)

概要 Let A and B be uniformly closed function algebras on locally compact Hausdorff spaces X and Y , respectively. We shall present a positive answer to Tingley's problem for A and B . More explicitly, we prove that every surjective isometry between the unit spheres, $S(A)$ and $S(B)$, of A and B admits an extension to a surjective real linear isometry between whole spaces A and B .

- 29 佐野隆志 (山形大理)^b クラウス行列の固有値分布について 10
武内和希 (山形大理工)
Takashi Sano (Yamagata Univ.) Inertia of Kraus matrices
Kazuki Takeuchi (Yamagata Univ.)

概要 Let $r, p_0, p_1 < \dots < p_n$ be positive numbers. We consider the $n \times n$ matrix whose (i, j) entry is given as $\frac{1}{p_i - p_j} \left(\frac{p_i^r - p_0^r}{p_i - p_0} - \frac{p_j^r - p_0^r}{p_j - p_0} \right)$. We call this a Kraus matrix associated with t^r . We show the inertia of this Kraus matrix.

- 30 渚勝 行列の p -ノルム 10
(立命館大理工・千葉大理)
Masaru Nagisa The p -norm of some matrices
(Ritsumeikan Univ./Chiba Univ.)

概要 We compute the operator p -norm of some $n \times n$ complex matrices, which can be seen as bounded linear operators on the n -dimensional Banach space $\ell^p(n)$. In particular, a p -norm of a matrix $A = \begin{pmatrix} 8 & 1 & 6 \\ 3 & 5 & 7 \\ 4 & 9 & 2 \end{pmatrix}$ which corresponds to a magic square is 15 for any $p \in [1, \infty]$.

- 31 遠山宏明 (前橋工科大) Operator inequalities related to Young's inequality 15
亀井栄三郎
渡邊雅之 (前橋工科大)
Hiroaki Tohyama Operator inequalities related to Young's inequality
(Maebashi Inst. of Tech.)
Eizaburo Kamei
Masayuki Watanabe
(Maebashi Inst. of Tech.)

概要 In this talk, we show several operator inequalities concerning Young's inequality by using relative operator entropies and operator valued α -divergence. One of them is partially a refinement of Young's inequality.

- 32 山崎 丈明 (東洋大理工) Limit of iterated induced Aluthge transformations of centered operators
 大坂 博幸 (立命館大理工) 15
 Takeaki Yamazaki (Toyo Univ.) Limit of iterated induced Aluthge transformations of centered operators
 Hiroyuki Osaka (Ritsumeikan Univ.)

概要 In this talk, we shall discuss on bounded linear operators on a complex Hilbert space. We will introduce convergence property of an operator sequence which is generated by iteration of the induced Aluthge transformation. In this talk, we consider the case of centered operators.

- 33 大坂 博幸 (立命館大理工) On a class of K -entanglement witnesses 15
 T. Młynik (Univ. of Gdańsk)
 M. Marciniak (Univ. of Gdańsk)
 Hiroyuki Osaka (Ritsumeikan Univ.) On a class of K -entanglement witnesses
 Tomasz Młynik (Univ. of Gdańsk)
 Marcin Marciniak (Univ. of Gdańsk)

概要 Recently, Yang et al. showed that each 2-positive map acting from $\mathcal{M}_3(\mathbb{C})$ into itself is decomposable. It is equivalent to the statement that each PPT state on $\mathbb{C}^3 \otimes \mathbb{C}^3$ has Schmidt number at most 2. It is a generalization of Perez–Horodecki criterion which states that each PPT state on $\mathbb{C}^2 \otimes \mathbb{C}^2$ or $\mathbb{C}^2 \otimes \mathbb{C}^3$ has Schmidt rank 1 i.e. is separable. Natural question arises whether the result of Yang et al. stays true for PPT states on $\mathbb{C}^3 \otimes \mathbb{C}^4$. This question can be considered also in higher dimensions. We construct a positive maps which is suspected for being a counterexample. More generally, we provide a class of positive maps between matrix algebras whose k -positivity properties can be easily controlled.

- 34 瀬尾 祐貴 (大阪教育大教育) 量子 \mathfrak{h}_α -Rényi ダイヴァージェンスのデータ処理不等式について 15
 Yuki Seo (Osaka Kyoiku Univ.) On data processing inequality for quantum \mathfrak{h}_α -Rényi divergences of positive order

概要 In this talk, we show some fundamental properties of quantum \mathfrak{h}_α -Rényi divergences of real order. In particular, we show that the data processing inequality holds for all $\alpha > 0$.

14:15~15:30

- 35 井上 寛 (九州産大経済) An unbounded generalization of Tomita's observable algebras II 15
 Hiroshi Inoue (Kyushu Sangyo Univ.) An unbounded generalization of Tomita's observable algebras II

概要 In a previous talk we tried to build the basic theory of unbounded Tomita's observable algebras called T^\dagger -algebras which are related to unbounded operator algebras, especially unbounded Tomita–Takesaki theory, operator algebras on Krein spaces, studies of positive linear functionals on $*$ -algebras and so on. And we defined the notions of regularity, semisimplicity and singularity of T^\dagger -algebras and characterized them. In this talk we shall proceed further studies of T^\dagger -algebras and investigate whether a T^\dagger -algebra is decomposable into a regular part and a singular part.

- 36 伊藤 慧 (東大数理) 複素力学系に付随する C^* -環のカルトン部分環 10
 Kei Ito (Univ. of Tokyo) Cartan subalgebras of C^* -algebras associated with complex dynamical systems

概要 Deaconu and Muhly was defined C^* -algebras associated with branched coverings. But their algebras lack information of branch points. Kajiwara and Watatani tried to modify Deaconu–Muhly algebras and defined C^* -algebras associated with complex dynamical systems. Let R be a rational function and J be its Julia set. It is known that the C^* -algebra $C(J)$ of complex-valued continuous functions on J is a Cartan subalgebra of the Deaconu–Muhly algebra for the dynamical system (J, R) , but it was not known whether $C(J)$ is a Cartan subalgebra of the Kajiwara–Watatani algebra for (J, R) or not. In this talk, I explain that $C(J)$ may not be a Cartan subalgebra of the Kajiwara–Watatani algebra for (J, R) .

- 37 有本 諒也 (京大数理研) On the type of the von Neumann algebra of an open subgroup of the Neretin group 15
 Ryoya Arimoto (Kyoto Univ.) On the type of the von Neumann algebra of an open subgroup of the Neretin group

概要 The Neretin group is the totally disconnected locally compact group consisting of almost automorphisms of the tree. In 2021, P.-E. Caprace, A. Le Boudec, and N. Matte Bon proved that the Neretin group is not of type I. We prove that a distinguished open subgroup of the Neretin group is not of type I either.

- 38 後藤 聡史 (上智大理工) On flat part commuting squares 10
 Satoshi Goto (Sophia Univ.) On flat part commuting squares

概要 When a non-degenerate commuting square is given, we can construct a hyperfinite II_1 subfactor $N \subset M$ generated by the tower of the given non-degenerate commuting squares. From the original commuting square, we can also construct the flat part commuting squares. If the subfactor $N \subset M$ is of finite depth, we will show that the composite connection of the original and the flat part ones (in the sense of N. Sato) produces a subfactor which is isomorphic to $N \subset M$. We also show an application of this result.

15:45~16:45 特別講演

- 渚 勝 コンパクト環上の非線型トレースとマジョリゼーション
 (立命館大理工・千葉大理)
 Masaru Nagisa Non-linear traces on the algebra of compact operators and majorization
 (Ritsumeikan Univ./Chiba Univ.)

概要 We study non-linear traces of Choquet type and Sugeno type on the algebra of compact operators. They have certain partial additivities. We show that these partial additivities characterize non-linear traces of both Choquet type and Sugeno type respectively. There exists a close relation between non-linear traces of Choquet type and majorization theory. We study trace class operators for non-linear traces of Choquet type. More generally we discuss Schatten–von Neumann p -class operators for non-linear traces of Choquet type. We determine when they form Banach spaces. This is an attempt of non-commutative integration theory for non-linear traces of Choquet type on the algebra of compact operators. We also consider the triangle inequality for non-linear traces of Sugeno type.

統計数学

3月15日(水) 第VIII会場

10:00~11:30

- 1 久保田直樹 (日大理工) フロッグモデルの時間定数におけるリップシツ型評価 15
Naoki Kubota (Nihon Univ.) Lipschitz-type estimates for the time constant of the frog model

概要 We consider the frog model on the multidimensional lattice, whose initial configuration is according to a Bernoulli distribution. The main object of interest is the so-called time constant, which describes the speed of the spread of frogs. The time constant depends on the parameter of the Bernoulli distribution. Hence, in this talk, we discuss how the change in the parameter of the Bernoulli distribution affects the time constant. More precisely, we provide some Lipschitz-type estimates for the time constant.

- 2 大井拓夢 (京大数理研) Convergence of time-changed α -stable processes by GMC 15
Takumu Ooi (Kyoto Univ.) Convergence of time-changed α -stable processes by GMC

概要 We consider Gaussian fields whose covariance kernels are 1-order green functions of d -dimensional symmetric α -stable processes, Gaussian multiplicative chaos (GMC) and time-changed α -stable processes by GMC. In the case of both d and α are 2, these are the Gaussian free field, the Liouville measure and the Liouville Brownian motion, respectively. In this talk, we explain weak convergence of the time-changed α -stable processes by GMC with Skorokhod's topology.

- 3 広兼巳紀雄 (阪大基礎工) A limit theorem for generalized tempered stable processes and their
深澤正彰 (阪大基礎工) quadratic variations with stable index tending to two 15
Mikio Hirokane (Osaka Univ.) A limit theorem for generalized tempered stable processes and their
Masaaki Fukasawa (Osaka Univ.) quadratic variations with stable index tending to two

概要 Motivated by Yoshida's martingale expansion theory, we study the limit of the joint distribution of a multidimensional Generalized Tempered Stable (GTS) process and its quadratic covariation process when the stable index tends to two. Under a proper scaling, the GTS processes converges to a Brownian motion that is a stable process with stable index two. We renormalize their quadratic covariation processes so that they have a nondegenerate limit distribution. We show that the limit is a stable process with stable index one and is independent of the limit Brownian motion of the GTS processes.

- 4 鈴木由紀 (慶大医) A diffusion process with a symmetric stable potential on the negative
高橋弘 (慶大商) side in \mathbb{R} 15
田村要造
Yuki Suzuki (Keio Univ.) A diffusion process with a symmetric stable potential on the negative
Hiroshi Takahashi (Keio Univ.) side in \mathbb{R}
Yozo Tamura

概要 We investigate a diffusion process with a symmetric stable potential on the negative side in \mathbb{R} . In the case of a Brownian potential on the negative side, this process was examined by Kawazu, Suzuki and Tanaka (2001, 2006). We study a long-time behavior of our process by a different method from theirs. We use a method based on a theory of bi-generalized diffusion processes studied by Ogura (1989).

- 5 鈴木 由紀 (慶大 医) Limit theorems for a diffusion process with a non-selfsimilar random potential 15
 Yuki Suzuki (Keio Univ.) Limit theorems for a diffusion process with a non-selfsimilar random potential

概要 We study a one-dimensional diffusion process with a non-selfsimilar random potential. The type of the potential is different on the left and the right hand sides of the origin. We show that the long-time behavior of the process is different from the behaviors of any other previous models.

- 6 小川 重義 (立命館大理工) Another representation of the mean value theorem for noncausal stochastic integrals 10
 Shigeyoshi Ogawa (Ritsumeikan Univ.) Another representation of the mean value theorem for noncausal stochastic integrals

概要 This is the follow up to the preceding lecture “Mean value theorems for the noncausal stochastic integral” that was presented at the 2022 annual meeting of Math Society of Japan held in Sapporo, in October 2022. There we showed a basic fact about the validity of a mean value theorem for the noncausal stochastic integral, also called ogawa integral, with respect to Brownian motion W_t of the form $\int_s^t f(X_r) d_* W_r$ where X_t is an Itô process. We derived formulae of mean value theorem for Itô integral $\int_s^t f(X_r) d_0 W_r$ and for the noncausal integral $\int_s^t f(\tilde{X}_r) d_* W_r$ where $\tilde{X}_t(\phi)$ is a noncausal variation of X_t .

The aim of the present talk is to give an elementary but important comment on the basic result and show another formula of mean value theorem for each of three cases mentioned above.

14:15~15:15 特別講演

- 中島 秀太 (明大 理工) ファーストパッセージパーコレーションと関連する話題
 Shuta Nakajima (Meiji Univ.) First-passage percolation and its related topics

概要 In this talk, we discuss recent progress on First-passage percolation (FPP). FPP was originally introduced by Hammersley and Welsh as a model of the spread of fluid in a random medium. The model also can be regarded as a dynamic version of percolation. In FPP, a graph with random weights is given, and we consider the optimization problem of the passage time between two fixed vertices. The minimum value is called the first passage time and it represents the time when the fluid reaches from one point to the other. From the viewpoint of an optimization problem, properties of the optimal path that attains minimum value are also of interest. Over 50 years, many mathematical techniques have been developed to study these objects. This talk mainly focuses on the fluctuations and the large deviations of the first passage times, and the geometric properties of geodesics. We also discuss related models such as the chemical distance.

15:30~16:30 特別講演

梁 松 (早大教育) ブラウン運動と古典力学系
 Song Liang (Waseda Univ.) A mechanical model of Brownian motion

概要 We provide a connection between Brownian motion and a classical Newton mechanical system. Brownian motion is a well-known physical phenomenon concerning the erratic motion of a small particle immersed into a fluid in equilibrium. The physical explanation of Brownian motion is often presented in the following way: since the massive particle is interacted by a large number of very light environmental particles, if we could assume that the interactions from each light particle at each time are independent, then by the central limit theorem for the sums of *i.i.d.* random variables, this will give in a suitable limit the Brownian motion. We notice that this assumption of independence can hardly be justified, even in a collision model, because of the possibility of re-collisions. This becomes a more evident and significant drawback when considering the model of interactions caused by potentials.

In this talk, we consider a mechanical model consisting of a massive particle in an ideal gas, evolved according to non-random Newton mechanical principles, via interaction potentials. We prove that under certain conditions, the (position, velocity)-process of the massive particle converges to a diffusion process, when the mass of the environmental particles converges to 0, while the density and the velocities of them go to infinity promptly. The precise expression of the limiting process is also given.

3月16日(木) 第VIII会場

10:20~11:10

- 7 田中晴喜 (和歌山県立医大) On dimension estimates in nonconformal graph iterated function systems via asymptotic perturbation 15
 Haruyoshi Tanaka (Wakayama Med. Univ.) On dimension estimates in nonconformal graph iterated function systems via asymptotic perturbation

概要 We consider infinite graph-directed iterated function systems (GIFSs) whose contraction mappings are nonconformal. As our main result, we formulate asymptotic perturbations from conformal GIFSs to nonconformal GIFSs, and give the asymptotic behaviour of the Hausdorff dimension of the limit set of the perturbed system. We also investigate perturbed self-affine sets as special cases.

- 8 高野凌史 (阪大基礎工) Large deviation principle for the Lyons–Victoir extension 15
 Ryoji Takano (Osaka Univ.) Large deviation principle for the Lyons–Victoir extension

概要 We study the large deviation principle for the Lyons–Victoir extension. The Lyons–Victoir extension is known as a method to lift a general continuous path of finite p -variation to a geometric q -variation rough path, where q is greater than p . We use the Lyons–Victoir extension as a method for lifting continuous stochastic processes to geometric rough paths. The main idea of this study is that by using the Lyons–Victoir extension, lifting the large deviation principle for continuous stochastic processes to that for geometric rough paths. To do so, we will prove the boundedness of the Lyons–Victoir extension map. As consequence of this, the large deviation principle for rough differential equation driven by the Lyons–Victoir extension can be obtained.

- 9 柳下浩紀 (京都産大理) 無限次元空間 \mathbb{R}^∞ 上の Gibbs 測度は, コンパクト空間 $(\mathbb{R} \cup \{\infty\})^\infty$ 上の局所性をもつ正則 Dirichlet 形式を定める 10

Hiroki Yagisita (Kyoto Sangyo Univ.) Any Gibbs measure on infinite-dimensional space \mathbb{R}^∞ defines a regular Dirichlet form with local property on compact space $(\mathbb{R} \cup \{\infty\})^\infty$

概要 For Gibbs measure μ on \mathbb{R}^∞ , we consider a classical symmetric form formally denoted $E_{CD}^\mu(f_1, f_2) := \sum_{k \in \mathbb{N}} \int_{\mathbb{R}^\infty} \frac{\partial f_1}{\partial x_k} \frac{\partial f_2}{\partial x_k} d\mu$. Following, for example, Albeverio, Kondratiev, Rockner, etc., we adopt definition of Gibbs measure by establishment of integration by parts formula. In our main result, “ μ determines a regular Dirichlet form with local property”, there are no constraints other than that μ is (what we say) Gibbs measure. It seems that Gibbs measure (as we call it) also includes fairly marginal probability measure in that it is “spatially spread out”. Note that we see \mathbb{R}^∞ as a direct product of Riemannian manifolds.

- 10 中川雄太 (京大人間環境) ギブス点過程と非正のポテンシャルに関するシュレディンガー作用素の状態密度 10

Yuta Nakagawa (Kyoto Univ.) Density of states for Schrödinger operators associated with Gibbs point processes and nonpositive potentials

概要 In this talk, we consider the random Schrödinger operator for the potential generated by the nonpositive single site potential and the points randomly located according to the Gibbs point process for the pairwise interaction. We determine the leading term of the asymptotic behavior of the integrated density of states (IDS) for the Schrödinger operator. This IDS vanishes much faster near $-\infty$ than the IDS for the Poisson point process, of which asymptotic behavior is known.

11:20~11:50 統計数学分科会総会

3月17日(金) 第VIII会場

9:50~11:35

- 11 田中輝雄 (広島市大情報) 最適停止問題の預言者の不等式 —差の評価— 10

Teruo Tanaka (Hiroshima City Univ.) Prophet inequalities for finite stage multiparameter optimal stopping problems

概要 We consider the optimal stopping problem for discrete time multiparameter stochastic processes. In the classical optimal stopping problems, the comparisons between the expected reward of a player with complete foresight and the expected reward of a player using nonanticipating stop rules, known as prophet inequalities, have been studied by many authors. We shall study the prophet inequalities for optimal stopping problems for finite stage multiparameter stochastic processes.

- 12 藤田敏治 (九工大工) 合流型推移をもつマルコフ決定過程 —乗法型評価— 10

Toshiharu Fujita (Kyushu Inst. of Tech.) Markov decision process with converging branch system —Multiplicative reward system—

概要 We consider multiplicative reward system on a certain finite-stage stochastic converging decision process model. Converging decision process has converging transition, which is one of the nonserial branch systems proposed by Nemhauser. We apply bidecision approach to our model and give a recursive method by dynamic programming technique.

- 13 高橋 勇人 (Random Data Lab.) Posterior distributions weakly converge to Martin-Löf random parameters 15
 Hayato Takahashi (Random Data Lab.) Posterior distributions weakly converge to Martin-Löf random parameters

概要 In parametric models estimators that are consistent at all parameters are concerned, while in Bayes models those that are consistent at almost all parameters are concerned. The identification of the points at which the posterior distributions weakly converge constitutes the problem (Diaconis and Freedman 1986). We show that for joint probabilities on complete separable metric spaces, the posterior distributions are consistent at almost all parameters if and only if the posterior distributions are consistent at ML-random parameters. Reference: Hayato Takahashi, Bayesian definition of random sequences with respect to conditional probabilities, preprint

- 14 高橋 勇人 (Random Data Lab.) Some explicit formulae for the distributions of words 15
 Hayato Takahashi (Random Data Lab.) Some explicit formulae for the distributions of words

概要 Bassino et al. 2010 and Regnier et al. 1998 showed generating functions of the distributions of words for all sample sizes. Robin et al. 1999 presented generating functions of the distributions for the return time of words and demonstrated a recurrence formula for these distributions. These generating functions are rational functions; except for simple cases, it is difficult to expand them into power series. In this paper, we study finite-dimensional generating functions of the distributions of nonoverlapping words and some overlapping words for each fixed sample size, respectively and demonstrate the explicit formulae for the distributions of words for the Bernoulli models.

- 15 地 壽 頌 子 (大阪工大情報) Optimality of spanning bipartite block designs II 15
 栗 木 進 二 (阪 府 大*)
 藤 原 良 (筑 波 大*)
 宮 本 暢 子 (東京理大理工)
 Shoko Chisaki (Osaka Inst. of Tech.) Optimality of spanning bipartite block designs II
 Shinji Kuriki (Osaka Pref. Univ.*)
 Ryoh Fuji-Hara (Univ. of Tsukuba*)
 Nobuko Miyamoto (Tokyo Univ. of Sci.)

概要 It is usually to design an experiment using treatments and its blocks in the design of experiments. Consider a set of edges of a complete bipartite graph as a treatments set and suppose the treatments have a structure. Then, we proposed a spanning bipartite block design (SBBD) to achieve better estimation accuracy. In this talk, using an SBBD as a statistical model, we prove that the design matrix of an SBBD is A-optimal under some conditions.

- 16 種 市 信 裕 (北教大札幌) 多項分布の適合度検定統計量選択の指針とその応用 15
 関 谷 祐 里 (北教大釧路)
 Nobuhiro Taneichi An index for selecting the multinomial goodness-of-fit test statistics and
 (Hokkaido Univ. of Edu.) its applications
 Yuri Sekiya (Hokkaido Univ. of Edu.)

概要 For the goodness-of-fit test for a multinomial distribution, we consider phi-divergence family of goodness-of-fit test statistics. Members of phi-divergence family of statistics all have an equivalent limiting chi-square distribution under the null hypothesis. We consider a second-order correction term as an index whether the distribution of statistic is close to the limiting chi-square distribution. We derive important properties of the index for selecting a phi-divergence goodness-of-fit statistic. We also discuss an application of the index to the other kinds of tests.

- 17 栗田 絵梨 (東京理大理) 多変量歪度を用いた多変量正規性検定統計量の帰無分布について 15
 瀬尾 隆 (東京理大理)
 Eri Kurita (Tokyo Univ. of Sci.) On the null distribution of multivariate normality test statistics using
 Takashi Seo (Tokyo Univ. of Sci.) multivariate skewness

概要 In this talk, we consider on the null distribution of multivariate normality test statistic using multivariate skewness. Although there are several definitions of multivariate skewness and kurtosis, this talk focuses on Mardia's multivariate skewness and proposes a new chi-square approximation with adjusted degrees of freedom for the null distribution of the normality test statistic using it. Finally, some simulation results are presented to investigate the accuracy of the chi-square approximation of the test statistic proposed in this talk.

14:25~15:25 特別講演

- 松田 孟留 行列の縮小推定
 (東大情報理工・理化学研)
 Takeru Matsuda Shrinkage estimation: from vector to matrix
 (Univ. of Tokyo/RIKEN)

概要 In the estimation of a normal mean vector under the quadratic loss, the maximum likelihood estimator (MLE) is inadmissible and dominated by shrinkage estimators (e.g. James–Stein estimator) when the dimension is greater than or equal to three (Stein's paradox). In particular, generalized Bayes estimators with respect to superharmonic priors (e.g. Stein's prior) are minimax and dominate MLE. Note that a function is said to be superharmonic if its average value on a hypersphere is always not greater than its value at the center.

In this talk, I will introduce recent studies on generalizations of the above results to matrices. First, we develop a superharmonic prior for matrices that shrinks singular values, which can be viewed as a natural generalization of Stein's prior. This prior is motivated from the Efron–Morris estimator, which is an extension of the James–Stein estimator to matrices. The generalized Bayes estimator with respect to this prior is minimax and dominates MLE under the Frobenius loss. In particular, since it shrinks to the space of low-rank matrices, it attains large risk reduction when the unknown matrix is close to low-rank (e.g. reduced-rank regression). Next, we construct a theory of shrinkage estimation under the “matrix quadratic loss”, which is a matrix-valued loss function suitable for matrix estimation. A notion of “matrix superharmonicity” for matrix-variate functions is introduced and the generalized Bayes estimator with respect to a matrix superharmonic prior is shown to be minimax under the matrix quadratic loss. The matrix-variate improper t-priors are matrix superharmonic and this class includes the above generalization of Stein's prior. Applications include matrix completion and nonparametric estimation.

15:40~16:40 特別講演

- 上原 悠 槓 (関西大システム理工) Information criteria for jump diffusion models
 Yuma Uehara (Kansai Univ.) Information criteria for jump diffusion models

概要 In this talk, we consider ergodic jump diffusion models with high-frequency observations. We mainly focus on the model selection problem for the models. More specifically, our interest is to select all drift and diffusion coefficients, and jump distribution among candidates. For such a problem, the threshold based transition density approximation which corresponds to the case where at most one jump occur within one observation interval is useful. We evaluate such an approximation, and propose explicit AIC-type information criterion constructed by the threshold based quasi-likelihood function.

3月18日(土) 第VIII会場

10:00~11:10

- 18 八木文香 (東京理大理) On the test for adequacy in growth curve model with two-step monotone missing data 15
尾崎冬弥 (東京理大理)
瀬尾隆 (東京理大理)
Ayaka Yagi (Tokyo Univ. of Sci.) On the test for adequacy in growth curve model with two-step monotone missing data
Toya Ozaki (Tokyo Univ. of Sci.)
Takashi Seo (Tokyo Univ. of Sci.)

概要 We consider the test for adequacy in growth curve model when the data set has a two-step monotone missing pattern. For the complete data case, to test the adequacy hypothesis, Hotelling's T^2 statistic is presented in Fujikoshi, Ulyanov and Shimizu (2010). In this talk, we propose a test statistic for the adequacy hypothesis with two-step monotone missing data based on the above result. Further, we give the approximation to the upper percentile of this statistic. Finally, we investigate the accuracy of the approximation by Monte Carlo simulation.

- 19 小池祐太 (東大数理) 対数凹な独立同分布確率ベクトルの和に対する高次元中心極限定理 15
Xiao Fang
 (Chinese Univ. of Hong Kong)
Yuta Koike (Univ. of Tokyo) A high-dimensional central limit theorem for sums of i. i. d log-concave random vectors
Xiao Fang
 (Chinese Univ. of Hong Kong)

概要 We consider the problem of bounding the normal approximation error over rectangles for a sum of n independent d -dimensional random vectors. Here, our aim is to establish such a bound with poly-logarithmic dependence on the dimension d . It is known that such a bound is available with a nearly $n^{-1/2}$ convergence rate when the covariance matrix of the sum is non-degenerate. In this talk, we show that, if the random vectors are i.i.d. and log-concave, we can derive an error bound with a nearly $n^{-1/2}$ convergence rate and poly-log dependence on d without any restriction on the covariance matrix.

- 20 山形浩一 (国立情報学研) 局所漸近正規性を有する量子統計モデルに対する推定量の有効性 15
藤原彰夫 (阪大理)
Koichi Yamagata Efficiency of estimators for locally asymptotically normal quantum statistical models
 (Nat. Inst. of Informatics)
Akio Fujiwara (Osaka Univ.)

概要 We herein establish an asymptotic representation theorem for locally asymptotically normal quantum statistical models. This theorem enables us to study the asymptotic efficiency of quantum estimators such as quantum regular estimators and quantum minimax estimators, leading to a universal tight lower bound beyond the i.i.d. assumption. This formulation complements the theory of quantum contiguity developed in the previous paper [Fujiwara and Yamagata, *Bernoulli* **26** (2020) 2105–2141], providing a solid foundation of the theory of weak quantum local asymptotic normality.

- 21 中村慶太 (東京理大理工) Aitchison 幾何を用いた正方分割表の対称性 15
中川智之 (東京理大理工)
田畑耕治 (東京理大理工)
Keita Nakamura (Tokyo Univ. of Sci.) Symmetry in square contingency tables using aitchison geometry
Tomoyuki Nakagawa
(Tokyo Univ. of Sci.)
Kouji Tahata (Tokyo Univ. of Sci.)

概要 The probability table obtained from a contingency table can be viewed as an element in the simplex. Symmetric probability tables are identified as a linear subspace using the Aitchison geometry of the simplex. Given a probability table, we can obtain the closest symmetric table by orthogonal projection onto the symmetric subspace. The original probability table is decomposed into the symmetric and the skew-symmetric tables that are orthogonal to each other. A procedure to test symmetry of a contingency table, based on a multinomial simulation, is developed and examples of tests carried out are given.

応 用 数 学

3月15日(水) 第VII会場

9:50~11:50

- 1 田中優帆 (早大理工) Jacobi 多項式とデザイン理論について 15
三枝崎剛 (早大理工)
大浦学 (金沢大理工)
H. S. Chakraborty
(Shahjalal Univ. of Sci. Tech.)
- Yuhō Tanaka (Waseda Univ.) Jacobi polynomials and design theory
Tsuayoshi Miezaki (Waseda Univ.)
Manabu Oura (Kanazawa Univ.)
Himadri Shekhar Chakraborty
(Shahjalal Univ. of Sci. Tech.)

概要 There are Jacobi polynomials as an analogue to Jacobi forms as a powerful generalization of theta series of Lattices. It is a generalization of weight enumerators that M. Ozeki introduced for codes. A. Bonnetcaze et al. constructed various types of designs and did some studies to establish a relationship between Jacobi polynomials and designs by focusing on weight enumerators, Jacobi polynomials, and the Molien series. In this talk, we introduce the concept of multiple Jacobi polynomials and give the MacWilliams type duality for them. We also give generalizations and analogies of some results in their paper. In addition, we introduce a criterion for the existence of (generalized) designs using Jacobi polynomials.

- 2 須田庄 (防衛大) 4-(11,5,1) デザインに関連するアソシエーション・スキームの一意性 .. 10
A. L. Gavriljuk (島根大総合理工)
- Sho Suda (Nat. Defense Acad. of Japan) Uniqueness of an association scheme related to the 4-(11,5,1)-design
Alexander L. Gavriljuk
(Shimane Univ.)

概要 It follows from the Delsarte theory that the Witt 4-(11,5,1) design gives rise to a (Q -polynomial) association scheme \mathcal{W} defined on the set of its blocks. In this talk, we observe that \mathcal{W} is unique, i.e., defined up to isomorphism by its parameters.

- 3 尾白典文 (豊田工大工) 準巡回符号における LCD 性と反転不変性 15
松井一 (豊田工大工)
- Norifumi Ojiro (Toyota Tech. Inst.) LCD property and reversibility in quasi-cyclic codes
Hajime Matsui (Toyota Tech. Inst.)

概要 We show a necessary and sufficient condition for which quasi-cyclic codes represented by polynomial matrices are to be LCD (Linear Complimentary Dual), and using this result, a connection between LCD quasi-cyclic codes and reversible quasi-cyclic codes is investigated.

- 4 佐竹翔平 (明大総合数理) 高い内周をもつ弧推移的エクスパンダーグラフ 15
Hyungrok Jo (横浜国大 IAS)
- Shohei Satake (Meiji Univ.) On high-girth arc-transitive expander graphs
Hyungrok Jo (Yokohama Nat. Univ.)

概要 Expander graphs are highly-connected sparse graphs which have fruitful connections to mathematics, computer science and cryptography. One of the fundamental problems about expander graphs is to construct explicit high-girth expander graphs, which can be applied for designing cryptographic hash functions, for example. In this talk we introduce new explicit constructions of high-girth expander graphs that are arc-transitive.

- 5 藤田 慎也 (横浜市大データサイエンス) Connectivity keeping paths in highly connected graphs 15
Shinya Fujita (Yokohama City Univ.) Connectivity keeping paths in highly connected graphs

概要 Some recent results concerning connectivity keeping paths in highly connected graphs will be reviewed.

- 6 藤村 丞 (福岡大情報基盤センター) On the generation of all Euler trails for an Eulerian graph with 2^n edges 10
白石 修二 (福岡大理) Sho Fujimura (Fukuoka Univ.) On the generation of all Euler trails for an Eulerian graph with 2^n edges
Shuji SHIRAISHI (Fukuoka Univ.)

概要 For an Eulerian graph G with 2^n edges, we give a new method of generating all Euler trails in G in parallel efficiently by using 2-chain cover of G .

- 7 町出 智也 (国立情報学研) Boolean 多項式の連立方程式の彩色問題への応用 15
Tomoya Machide (Nat. Inst. of Informatics) An application of systems of Boolean polynomial equations to list color problems

概要 I will talk an application of systems of Boolean polynomial equations to list color problems, in which not only vertex-coloring conditions but also edge-coloring conditions are imposed under Grünbaum method.

- 8 高橋 正 (甲南大知能情報) グレブナー基底を用いた学習モデルにおける補題証明 10
鷲野 朋広 (甲南大知能情報) Tadashi Takahashi (Konan Univ.) On the proof of lemma in the learning model using Groebner basis
Tomohiro Washino (Konan Univ.)

概要 We consider the statistical model of three-layer neural network with H hidden units. By using Taylor expansion of activation function, the set of parameters that true density function is realizable by a statistical model is a common zero points defined by finite polynomials. First, in this case $H = 2$, we calculate equations define the algebraic set. The algebraic set is the set of parameters that true density function is realizable by a statistical model with one or zero hidden unit. Next, we determine equations define the algebraic set. The algebraic set is the set of parameters that true density function is realizable by a statistical model with $H_0 < H$ hidden unit. We prove the lemma in the learning model applying Hilbert's basis theorem using Groebner basis.

14:15~15:25

- 9 辻 栄周平 (北教大旭川) グラフ埋め込みと辺に付随する互換の積 15
内海 凌 (北大理) Shuhei Tsujie (Hokkaido Univ. of Edu.) Graph embedding and the product of transpositions associated with
Ryo Uchiumi (Hokkaido Univ.) edges

概要 Given a graph, we associate each edge with the transposition which exchanges the endvertices. Fixing a linear order on the edge set, we obtain a permutation of the vertices. Dénes proved that the permutation is a full cyclic permutation for any linear order if and only if the graph is a tree.

In this article, we characterize graphs having a linear order such that the associated permutation is a full cyclic permutation in terms of graph embeddings.

- 10 山田 祐見 (筑波大システム情報) 格子凸多面体の free sum の Ehrhart 多項式の根の分布 15
八森 正泰 (筑波大システム情報)
東谷 章弘 (阪大情報)
Yumi Yamada (Univ. of Tsukuba) The root distributions of Ehrhart polynomials of free sums of lattice
Masahiro Hachimori polytopes
(Univ. of Tsukuba)
Akihiro Higashitani (Osaka Univ.)

概要 In this talk, we study the root distributions of Ehrhart polynomials of free sums of certain lattice polytopes. We investigate cases where the roots of the Ehrhart polynomials of the free sums of A_d^* 's or A_d 's lie on the canonical line $Re(z) = -1/2$ on the complex plane \mathbb{C} , where A_d denotes the root polytope of type A of dimension d and A_d^* denotes its polar dual. For example, it is proved that $A_m^* \oplus A_n^*$ with $\min\{m, n\} \leq 1$ or $m + n \leq 7$, $A_2^* \oplus (A_1^*)^{\oplus n}$ and $A_3^* \oplus (A_1^*)^{\oplus n}$ for any n satisfy this property.

- 11 東谷 章弘 (阪大情報) 行列のスウィッチングと付随する単体的複体による歪多項式環の分類 15
上山 健太 (弘前大教育)
Akihiro Higashitani (Osaka Univ.) Classification of skew polynomial algebras by switching of matrices and
Kenta Ueyama (Hirosaki Univ.) its associated simplicial complexes

概要 Classifying skew polynomial rings is important in noncommutative algebraic geometry because a skew polynomial ring can be regarded as a noncommutative version of the homogeneous coordinate ring of the projective space. In this talk, we focus on a particular class of skew polynomial rings and discuss whether the classification of such skew polynomial rings up to equivalence of their graded module categories can be done by using the switching of matrices and the simplicial complexes associated to switching equivalence classes. We prove that we can do it if the sizes of the matrices are relatively small.

- 12 栗原 大武 (山口大創成) 根付き木を用いた等質ラグランジュ部分代数の構成 15
橋永 貴弘 (佐賀大教育)
Hirotake Kurihara (Yamaguchi Univ.) Construction of homogeneous Lagrangian subalgebras using rooted trees
Takahiro Hashinaga (Saga Univ.)

概要 In this talk, we consider the construction and classification problems of homogeneous Lagrangian submanifolds of the noncompact Hermitian symmetric space $M = \mathrm{Sp}(n, \mathbb{R})/\mathrm{U}(n)$ of rank n . We give that rooted trees contribute to the classification of homogeneous Lagrangian submanifolds in $M = \mathrm{Sp}(n, \mathbb{R})/\mathrm{U}(n)$.

15:45~16:45 特別講演

- 野口 健太 (東京理大理工) 閉曲面上のグラフ, とくに三角形分割と四角形分割
Kenta Noguchi (Tokyo Univ. of Sci.) Graphs on closed surfaces, expressly triangulations and quadrangulations

概要 Graphs on closed surfaces are widely studied in the literature. We have been interested in and studied the triangulations and quadrangulations, which are important classes of such graphs. Specifically, we have been investigated the relationship between them to reveal their several properties, such as the chromatic number and the related topics. In this talk, we will present the basic background and some problems, and give recent results on the extension from a quadrangulation to a triangulation and spanning quadrangulation subgraphs of triangulations.

3月16日(木) 第VII会場

10:00~11:45

- 13 佐藤 巖 (小山工高専) Metzler/Zeta 対応 15
 井手 勇介 (日大文理)
 小松 堯
 (産業数理研究所Calc・広島大先進理工)
 今野 紀雄 (横浜国大工)
 Iwao Sato (Oyama Nat. Coll. of Tech.) Metzler/Zeta correspondence
 Yusuke Ide (Nihon Univ.)
 Takashi Komatsu
 (Math. Res. Inst. Calc for Industry/Hiroshima Univ.)
 Norio Konno (Yokohama Nat. Univ.)

概要 We present an explicit formula for the determinant on the Metzler matrix of a digraph D . Furthermore, we introduce a walk-type zeta function with respect to this Metzler matrix of the symmetric digraph of a finite torus, and express its limit formula by using the integral expression.

- 14 佐藤 巖 (小山工高専) Alternating walk/Zeta 対応 15
 小松 堯
 (産業数理研究所Calc・広島大先進理工)
 今野 紀雄 (横浜国大工)
 Iwao Sato (Oyama Nat. Coll. of Tech.) Alternating walk/Zeta correspondence
 Takashi Kmatsu
 (Math. Res. Inst. Calc for Industry/Hiroshima Univ.)
 Norio Konno (Yokohama Nat. Univ.)

概要 We express the alternating zeta function of a graph G by using the Ihara zeta function of G . Next, we define a generalized alternating zeta function of a graph, and express it of a vertex-transitive regular graph by spectra of the transition probability matrix of the symmetric simple random walk on it and its Laplacian. Furthermore, we present an integral expression for the limit of the generalized alternating zeta functions of a series of vertex-transitive regular graphs. As an example, we treat the generalized alternating zeta functions of a finite torus. Finally, we treat the relation between the Mahler measure and the alternating zeta function of a graph.

- 15 穴原 義弘 (横浜国大先進実践) 流出入を伴う 1次元格子上量子ウォークの収束速度 15
 瀬川 悦生 (横浜国大環境情報)
 森岡 悠 (愛媛大理工)
 Yoshihiro Anahara (Yokohama Nat. Univ.) Convergence speed for quantum walks on one dimensional lattice with
 inflows and outflows
 Etsuo Segawa (Yokohama Nat. Univ.)
 Morioka Hisashi (Ehime Univ.)

概要 For quantum walks on finite graphs whose time evolution is unitary operator, it is difficult to find a stationary state which is not equal to the initial state. We can consider a quantum walk model with a stationary state on a connected finite graph by introducing infinite length tails and inserting a quantum walker into the internal graph at every time step from the tail. In this talk focusing on the finite path as the internal graph, and also on the spectrum of the truncated submatrix of the unitary time evolution operator with respect to the internal graph, we estimate a convergence speed to the stationary state.

- 16 成松 明廣 (福知山公大数理・データサイエンスセンター) 完全状態遷移, グラフの衝平分割と連続時間量子ウォークによる探索アル
井手 勇介 (日大文理) ゴリズム 15
Akihiro Narimatsu Perfect state transfer, equitable partition and continuous-time quantum
(Univ. of Fukuchiyama) walk based search
Yusuke Ide (Nihon Univ.)

概要 We consider a continuous-time quantum walk based search algorithm. We introduce equitable partition of the graph and perfect state transfer on it. By these two methods, we can calculate the success probability and the finding time of the search algorithm. In addition, we gave some examples of graphs that we can calculate the success probability and the finding time.

- 17 藪 奥 哲史 (北九州工高専) Enumeration of connected bipartite graphs with given Betti number · · 15
蓮井 太朗 (九大数理)
白井 朋之 (九大IMI)
Satoshi Yabuoku Enumeration of connected bipartite graphs with given Betti number
(Kitakyushu Nat. Coll. of Tech.)
Taro Hasui (Kyushu Univ.)
Tomoyuki Shirai (Kyushu Univ.)

概要 Let $G = (V, E)$ be a simple connected bipartite graph with bipartition $V = V_1 \sqcup V_2$, and $f(r, s, q)$ denote the number of G s such that $|V_1| = r, |V_2| = s$ and $|E| = q$. In this talk, we focus on the values $f(r, s, r + s + k - 1)$ for given Betti number k . For $k = 0$, this is the number of spanning trees on $K_{r,s}$, and Scoins proved $f(r, s, r + s - 1) = r^{s-1}s^{r-1}$. By recurrence equation, we obtain the explicit form of exponential generating functions of $f(r, s, r + s + k - 1)$ for general k , and we derive the asymptotic behavior of their coefficients. We also give another expression of the generating functions as the summation of rational functions of those for the number of bipartite rooted spanning trees.

- 18 山岸 弘幸 (都立産業技術高専) 完全グラフに対応する ℓ^p 離散ソボレフ不等式の最良定数 15
渡辺 宏太郎 (防衛大)
Hiroyuki Yamagishi The best constant of the discrete ℓ^p Sobolev inequality on the complete
(Tokyo Metro. Coll. of Ind. Tech.) graph
Kohtaro Watanabe
(Nat. Defense Acad. of Japan)

概要 We assume a classical mechanical model of the complete graph K_N with N vertices. We put the N atoms on vertices of K_N . Its neighboring two atoms are connected to a linear spring. The spring constant is 1. Let p and q be real numbers satisfying $1/p + 1/q = 1$ ($p, q > 1$). If $u(i)$ takes a real value, then $u(i)$ represents a deviation from the steady state. The discrete ℓ^p Sobolev inequality shows that the maximum of deviation $|u(i)|^p$ is estimated in a constant multiple of the potential energy on K_N . Hence, the best constants represent the rigidity of the mechanical model on K_N .

13:00~14:10

- 19 盧 曉南 (岐阜大工) Covering arrays guaranteeing a passing test 15
Xiao-Nan Lu (Gifu Univ.) Covering arrays guaranteeing a passing test

概要 Covering arrays (CAs) are a kind of combinatorial designs proposed for applications in combinatorial testing (CT). For adaptive decoding algorithms in CT using CAs, it is required that a passing test should be contained in the CA. In this talk, a few results on such binary CAs that guarantee at least one passing test are introduced. In particular, it is shown that for no more than two faulty interactions, the known optimal construction for binary CAs is good enough. Then, a general sufficient condition for such CAs in terms of domination numbers of graphs associated to the CAs is proposed.

- 20 齋藤 明 (日大文理) 安全支配数に関する2つの上界…………… 15
 出川 慎悟 (日大文理)
 増田 合氣 (日大文理)
 西田 大輝 (日大文理)

Akira Saito (Nihon Univ.) Two upper bounds on secure domination number
 Shingo Degawa (Nihon Univ.)
 Aiki Masuda (Nihon Univ.)
 Hiroki Nishida (Nihon Univ.)

概要 A set of vertices S in a graph G is a dominating set if every $v \in V(G) - S$ satisfies $N_G(v) \cap S \neq \emptyset$. A dominating set S is secure if for every $v \in V(G) - S$, there exists $x \in S$ such that $(S - \{x\}) \cup \{v\}$ is a dominating set. The order of a smallest secure dominating set of G is called the secure domination number of G and denoted by $\gamma_s(G)$. In this talk, we report an upper bound $\gamma_s(G) \leq \frac{3}{5}|G|$ for a graph G of minimum degree at least 2. Then we generalize the notion of a secure dominating set. A dominating set $S = \{x_1, \dots, x_k\}$ in G is weakly secure if for every $v \in V(G) - S$, there exists $y_i \in N_G(x_i) \cup \{x_i\}$ for each i with $1 \leq i \leq k$ such that $\{y_1, \dots, y_k\}$ is a dominating set and $v \in \{y_1, \dots, y_k\}$. The order of a smallest weakly secure dominating set of G is called the weakly secure domination number and denoted by $\gamma_{ws}(G)$. We report $\gamma_{ws}(G) \leq \lceil |G|/2 \rceil$ for every connected graph G of minimum degree at least 1.

- 21 吉田 裕一 (北見工大工) 連結単純グラフのマッチングに付随する3種の不変量…………… 15
 松田 一徳 (北見工大工)

Yuichi Yoshida (Kitami Inst. of Tech.) On the three graph invariants related to matching of finite simple graphs
 Kazunori Matsuda (Kitami Inst. of Tech.)

概要 We determine the relationship between the matching number, the minimum matching number, the induced matching number and the number of vertices.

- 22 R. E. L. Aldred (Univ. of Otago) 閉曲面上の三角形分割でない5-連結グラフにおけるマッチング拡張問題
 藤沢 潤 (慶大商) …………… 15

Robert E. L. Aldred (Univ. of Otago) Distance restricted matching extension in 5-connected graphs on surfaces which are not triangulation
 Jun Fujisawa (Keio Univ.)

概要 In this talk, we show the following result. Let G be a 5-connected graph of even order which is 2-cell embedded in a surface F^2 with sufficiently large face-width, and let M be a matching of G such that any two edges in M are at distance at least $2 \left(20 - \frac{12\chi(F^2)+25}{|M|} \right)$ in the face subdivision G . Then G has a perfect matching containing M .

3月17日(金) 第VII会場

9:50~11:50

- 23 三好 啓也 (埼玉大理工) Derivation of Brownian motion from the Goldstein–Taylor model…………… 15
 Hironari Miyoshi (Saitama Univ.) Derivation of Brownian motion from the Goldstein–Taylor model

概要 In this lecture, we consider a random walk associated with the initial value problem of the Goldstein–Taylor model. First, we show the existence of the solution for a scaled Goldstein–Taylor model by using the corresponding telegraph equation. Next, we find the probability density function $p_\varepsilon(x, t)$ associated with a random process of particles for the Goldstein–Taylor model. Finally, we prove that the scaling limit of the cumulative distribution function for $p_\varepsilon(x, t)$ converges to the Brownian motion in the sense of distributions.

- 27 米田 剛 (一橋大経済) 深層ニューラルネットワークのミニバッチ勾配降下に対する各点収束定理
 15

Tsuyoshi Yoneda (Hitotsubashi Univ.) Pointwise convergence theorem of mini-batch gradient descent in terms of deep neural network

概要 The theoretical understanding of deep neural network (DNN) has not yet been fully understood. Nevertheless, even in this recent circumstance, Imaizumi-Fukumizu (2019) and Suzuki (2019) clarified that the learning ability of DNN is superior to the previous theories when the target function is non-smooth functions. In this talk we give a convergence problem in more practical setting. More precisely, we prove that a certain prescribed function pointwisely converges to a step function provided by mini-batch gradient descent process with ReLU DNN.

- 28 奥村真善美 (北大電子研) 毛包形成の数理モデルの数値計算とその考察 15
 小林康明 (北大電子研)
 長山雅晴 (北大電子研)
 藤原裕展 (理化学研)
 安ヶ平祐介 (日立製作所)
 大野航太 (中大理工)

Makoto Okumura (Hokkaido Univ.) Numerical simulations of a mathematical model of hair follicle formation
 Yasuaki Kobayashi (Hokkaido Univ.)
 Masaharu Nagayama (Hokkaido Univ.)
 Hironobu Fujiwara (RIKEN)
 Yasugahira Yusuke (Hitachi, Ltd.)
 Kota Ohno (Chuo Univ.)

概要 As an appendage of the skin, a hair follicle is the source of the hair. Moreover, the hair follicle has a cylinder-like structure. In this study, we attempt to examine the mechanism of hair follicle formation from a mathematical viewpoint by applying a mathematical model of the basement membrane, which enables us to describe plastic deformation. In this presentation, we will introduce numerical examples of our mathematical model and discuss what assumptions play an essential role in shaping the cylinder shape of the hair follicle. If possible, we will also present the numerical results that consider the epithelium, while previous mathematical models have considered only the basal layer, basement membrane, and dermal structure.

- 29 シュワドレンカカレル 構造材料の弾塑性有限変形の変分解析 15
 (京大理)

Karel Svadlenka (Kyoto Univ.) Variational analysis of finite strain elastoplasticity for structured materials

概要 Structured materials, such as metallic alloys with atomic-scale layers, show peculiar deformation patterns, which may have significant implications on material properties. In this talk, we discuss one possible approach to modeling and understanding of this kind of pattern formation through the so-called rate-independent evolution in the variational setting of finite-strain elastoplasticity. Besides mentioning connections to homogenization via Gamma-convergence, we present the underlying mathematical theory and show numerical simulations in comparison to experimental measurements.

14:15~14:40 2022年度日本数学会応用数学賞授賞式

14:45~16:30

- 30 土屋拓也 (八戸工大) de Sitter 時空中における半線形 Klein-Gordon 方程式の精度と数値安定
中村誠 (阪大情報) 性について 15
Takuya Tsuchiya Numerical accuracy and stability of semilinear Klein-Gordon equation
(Hachinohe Inst. of Tech.) in de Sitter spacetime
Makoto Nakamura (Osaka Univ.)

概要 The Klein-Gordon equation is known as the fundamental wave equation in relativistic phenomena such as near light speed phenomena. We perform numerical simulations of this equation using the structure-preserving scheme in the de Sitter spacetime. The discretization equation proposed in our previous work (J. Comput. Appl. Math. 361 pp.364-412) has good accuracy but poor numerical stability. In this talk, we propose a new discretization equation with the structure-preserving scheme and show numerical results with high-accuracy and high-stability.

- 31 土屋拓也 (八戸工大) 収縮する de Sitter 時空中における半線形 Klein-Gordon 方程式の解の挙動
中村誠 (阪大情報) について 15
Takuya Tsuchiya On behaviors of solutions of semilinear Klein-Gordon equation in con-
(Hachinohe Inst. of Tech.) tracting de Sitter spacetime
Makoto Nakamura (Osaka Univ.)

概要 de Sitter spacetime is a well-known model of expanding or contracting universes. In this spacetime, we investigate numerical solutions of the semilinear Klein-Gordon equation. In particular, in the case of contracting spacetime, it is difficult to investigate the behaviors of the solutions even by numerical calculations because of the blow-up solutions. Therefore, we apply the structure-preserving scheme to the Klein-Gordon equation to obtain accurate and stable numerical solutions and show the behaviors of the solutions.

- 32 竹田航太 (京大理) HMC による球面上 N 点渦系不変測度の計算 15
坂上貴之 (京大理)
Kota Takeda (Kyoto Univ.) Computing an invariant measure of the N -vortex problem on the sphere
Takashi Sakajo (Kyoto Univ.) by Hamiltonian Monte Carlo

概要 Hamiltonian Monte Carlo (HMC) is a stochastic method to approximate high-dimensional integrals by the samples from the target distribution, efficiently by utilizing the Hamiltonian dynamics. In general, the efficiency of HMC is characterized by a geometric convergence rate of the associated Markov chain, known as a property ‘geometric ergodicity’. For a class of target distributions on the Euclidean space, geometric ergodicity for HMC has been established, however, that has not yet been established on manifolds. Hence, we prove geometric ergodicity for HMC on compact manifolds. As an application, we consider the sampling problem from the invariant measure of the N -vortex problem on the sphere, which is relevant to two-dimensional turbulence on geophysical fluid.

- 33 藤原宏志 (京大情報) On a Cauchy-type singular integral equation for x-ray computerized tomography with partial measurement 15
 K. Sadiq
 (Johann Radon Inst. for Comput. Appl. Math.)
 A. Tamasan
 (Univ. Central Florida)
- Hiroshi Fujiwara (Kyoto Univ.) On a Cauchy-type singular integral equation for x-ray computerized tomography with partial measurement
 Kamran Sadiq
 (Johann Radon Inst. for Comput. Appl. Math.)
 Alexandru Tamasan
 (Univ. Central Florida)

概要 In this talk we propose a direct numerical reconstruction procedure for x-ray tomography with partial measurement. The proposed method is based on Bukhgeim–Cauchy boundary integral formula developed for an inverse source problem of the transport equation, and is characterized by a singular integral equation with Cauchy kernel for the considering partial measurement case. We also show its quantitative stability in terms of the condition number with a particular discretization for reliable numerical reconstruction.

- 34 石毛利昌 (千葉大融合理工) ある K3 曲面族についての Picard–Fuchs 微分方程式のモノドロミーに対する精度保証付き数値計算 15
 高安亮紀 (筑波大システム情報)
 Toshimasa Ishige (Chiba Univ.) Rigorous numerics for finding the monodromy of Picard–Fuchs differential equations for a family of K3 toric hypersurfaces
 Akitoshi Takayasu (Univ. of Tsukuba)

概要 In this talk, we introduce a method for finding monodromy matrices of linear differential equations with finite-dimensional solution spaces using rigorous numerics. We also provide a result of computer-assisted proof that yields monodromy matrices of Picard–Fuchs differential equations for a certain family of K3 toric hypersurfaces.

- 35 木下武彦 (佐賀大理工) 2 階楕円型線形作用素に対する近似逆作用素ノルムの収束オーダー評価 15
 渡部善隆
 (九大情報基盤研究開発センター)
 中尾充宏 (早大理工)
 Takehiko Kinoshita (Saga Univ.) Convergence order estimations of approximate inverse operator norm for second-order elliptic linear operators
 Yoshitaka Watanabe (Kyushu Univ.)
 Mitsuhiro T. Nakao (Waseda Univ.)

概要 In this talk, we consider the convergence order of the approximate inverse operator norm for second-order elliptic linear operators. We show that the general order of convergence is $O(\sqrt{h})$, and give concrete computable error bounds for achieving the same order.

16:45～17:45 特別講演

- 榎原航也 (岡山理大理・理化学研) 界面現象の構造保存型数値解析
 Koya Sakakibara Structure-preserving numerical analysis of interfacial phenomena
 (Okayama Univ. of Sci./RIKEN)

概要 Soap bubbles, falling water droplets, and other phenomena that deform from moment to moment are all around us. The problem of mathematically investigating these phenomena is known as the moving boundary problem, which has been studied in mathematics, physics, and various other fields. Numerical analysis of interfacial phenomena is more complicated than the usual numerical analysis of differential equations on a fixed domain because the problem domain changes from moment to moment. In this talk, we will review what has been known about attempts to design structure-preserving numerical methods for mathematical models describing interfacial phenomena. Further directions of research development will also be discussed.

3月18日(土) 第VII会場

9:50~11:50

- 36 國谷紀良 (神戸大システム情報) 隔離と時間遅れを考慮した感染症モデルにおけるホップ分岐 15
Toshikazu Kuniya (Kobe Univ.) Hopf bifurcation in a delayed epidemic model with quarantine

概要 In this study, we construct a time-delayed epidemic model with quarantine and obtain sufficient conditions for a Hopf bifurcation, which represents the recurrent epidemic waves. We show that if the basic reproduction number \mathcal{R}_0 is less than 1, then the disease-free equilibrium is globally asymptotically stable, whereas if $\mathcal{R}_0 > 1$, then the unique endemic equilibrium exists and it can be destabilized and a nontrivial periodic solution can arise in a Hopf bifurcation. Our numerical simulation suggests that the quarantine and time delay play important roles in the occurrence of recurrent epidemic waves.

- 37 竹口佳輝 (京大情報) 力学系理論の手法による感染症数理モデル SEIR 系の最適制御 15
矢ヶ崎一幸 (京大情報)
Yoshiki Takeguchi (Kyoto Univ.) Optimal control of the SEIR epidemic model using a dynamical systems
Kazuyuki Yagasaki (Kyoto Univ.) approach

概要 We consider the susceptible-exposed-infected-removed (SEIR) epidemic model and apply optimal control to it. Our approach is to reduce the computation of the optimal control input to that of the stable manifold of an invariant manifold in a Hamiltonian system. Some numerical examples in which the computer software AUTO is used to numerically compute the stable manifold are given to demonstrate the usefulness of our approach for the optimal control in the SEIR model. Our study suggests how we can decrease the number of infected individuals quickly before a critical situation occurs while keeping social and economic burdens small.

- 38 田崎創平 (北大理) 腸管上皮細胞社会における脱分化現象の数理解析 15
長井広樹
(東大薬・東北大学際フロンティア研)
ルイスアウグストエイジナガイ
(東大薬)
中戸隆一郎 (東大定量生命研)
三浦正幸 (東大薬)
中嶋悠一郎
(東大薬・東北大学際フロンティア研)
Sohei Tasaki (Hokkaido Univ.) Mathematical analysis of dedifferentiation in the intestinal epithelial cell
Hiroki Nagai community
(Univ. of Tokyo/Tohoku Univ.)
Luis Augusto Eijy Nagai
(Univ. of Tokyo)
Ryuichiro Nakato (Univ. of Tokyo)
Masayuki Miura (Univ. of Tokyo)
Yu-ichiro Nakajima
(Univ. of Tokyo/Tohoku Univ.)

概要 The intestinal epithelium is a highly flexible and adaptive cell community that is formed and regulated by the proliferation and differentiation of intestinal stem cells (ISCs). We have quantitatively analyzed the nutrient-dependent tissue expansion of the adult *Drosophila* intestinal epithelium just after eclosion. We have shown that stem cell division alone cannot explain the increase in stem cell number, and have proposed a novel dedifferentiation phenomenon as a possible compensatory mechanism. We have developed a mathematical model of intestinal epithelial cell society and established a simulation system that enables quantitative prediction of cell population dynamics. The analysis is revealing that enteroendocrine cell dedifferentiation is a good way to efficiently achieve intestinal enlargement while maintaining cell diversity.

- 39 松江 要 常微分方程式の爆発解の漸近展開と無限遠ダイナミクス 15

(九大IMI・九大I2CNER)

落合啓之(九大IMI)

小谷久寿(九大IMI)

高安亮紀(筑波大システム情報)

Kaname Matsue

(Kyushu Univ./Kyushu Univ.)

Hirofumi Ochiai (Kyushu Univ.)

Hisatoshi Kodani (Kyushu Univ.)

Akitoshi Takayasu (Univ. of Tsukuba)

Correspondence between asymptotic expansions of blow-up solutions for ODEs and dynamics at infinity

概要 We provide a natural correspondence of eigenstructures of Jacobian matrices associated with equilibria for appropriately transformed two systems (describing asymptotic expansions and dynamics at infinity) describing finite-time blow-ups for ODEs with quasi-homogeneity in an asymptotic sense. As a corollary, we see that asymptotic expansions of blow-ups themselves provide a criterion of the existence of blow-ups with an intrinsic gap structure of stability information among two systems.

- 40 劉 逸 侃(北大電子研) 非整数階拡散方程式系の初期値境界値問題について 15

Yikan Liu (Hokkaido Univ.)

Initial-boundary value problems for coupled systems of time-fractional diffusion equations

概要 This talk is concerned with the initial-boundary value problem for a moderately coupled system of time-fractional diffusion equations with Caputo derivatives of orders in $(0, 1)$. Regarding the coupled part of components as a source term, we define the mild solution by an integral equation. Then we establish fundamental unique existence, limited smoothing property and long-time asymptotic behavior of the solution, which mostly inherit those of a single equation.

- 41 宮路智行(京大理) A conjecture on the asymptotic reflection rule of a self-propelled particle
R. Sinclair 15

Tomoyuki Miyaji (Kyoto Univ.)

A conjecture on the asymptotic reflection rule of a self-propelled particle

Robert Sinclair

概要 We consider a four-dimensional ODE model of a self-propelled particle that interacts repulsively with the domain boundary. It is derived from certain planar reaction-diffusion systems. A previous study proves that the particle governed by this model moves almost in a straight line and reflects from the boundary without collision. In addition, numerical simulations suggest that the angle of reflection is greater than that of incidence, which is proved under certain conditions. Recently, we have proposed a conjecture on the functional relationship between the angles. We will discuss how we arrived at the conjecture and became convinced.

- 42 石井宙志(京大ASHBi) 十分弱い非局所効果を持つ反応拡散方程式における積分核形状に依存したパターンダイナミクス 15

Hiroshi Ishii (Kyoto Univ.)

Pattern dynamics depending on integral kernel shape in reaction-diffusion equations with sufficiently weak nonlocal effect

概要 In this talk, we consider reaction-diffusion equations with nonlocal effects described by the convolution with an appropriate integral kernel. Such a nonlocal effect is used to describe the diffusion process of substances and cell-cell interactions, for example. This work is particularly concerned with the behavior of the widths of phase-separated patterns. The equation of motion of the width is derived and applied to several integral kernels as examples to explain how the pattern dynamics of the solution depend on the shape of the integral kernel.

14:15~15:05

- 43 大野航太 (中大理工) Stability of traveling wave in nonlocally coupled oscillator system 15
小川知之 (明大先端数理)
Kota Ohno (Chuo Univ.) Stability of traveling wave in nonlocally coupled oscillator system
Toshiyuki Ogawa (Meiji Univ.)

概要 Coupled oscillator systems have been studied in various fields of physical and biological phenomena. Among them, systems of nonlocally coupled oscillators can exhibit chimera states, which consist of spatially coherent and incoherent states. However, the stability and bifurcation origin of chimera states have not been understood. Stuart–Landau type nonlinearity enables us to formulate the stability problem of traveling waves precisely through Floquet theory. We analyzed Floquet multiplier and confirmed the transition of stability of traveling wave by changing the parameters of nonlocal coupling. In this study, we will state the relationship between the stability of traveling waves and chimera states.

- 44 関坂歩幹 (明大研究・知財) 振動場反応拡散系に現れるセグメントパターンの安定性について 15
小川知之 (明大先端数理)
Ayuki Sekisaka (Meiji Univ.) Stability problem of segmented pattern for oscillatory reaction-diffusion
Toshiyuki Ogawa (Meiji Univ.) systems

概要 In reaction-diffusion systems of oscillatory fields, especially those in which wave bifurcations occur, it is known that traveling waves with periodic profiles appear. Even if the traveling waves are asymptotically stable, starting from a random initial value, several segments are formed, each consisting of different periodic traveling waves. Pattern dynamics is then observed in which the segments change from one to another. The transition between different segments is considered to be the so-called defect solution. In this lecture, we will discuss whether the defect solution can be regarded as a steady-state observation of the segment pattern.

- 45 小川知之 (明大先端数理) Compact metric graph 上の Turing 不安定性 15
小林俊介 (宮崎大工)
Toshiyuki Ogawa (Meiji Univ.) Turing instability on compact metric graph
Shunsuke Kobayashi
(Univ. of Miyazaki)

概要 Studies on the behavior of reaction-diffusion systems on a one-dimensional domain with junctions (metric graph) have been increasing recently. It is expected that joint research with cell physiology and ecology will progress more and more in the future. So, we would like to clarify from the basics what kind of pattern formation occurs in compact metric graphs. First, by preparing 2 or 3 bounded intervals, let us classify the metric graph that is created by connecting them, and consider the eigenvalue problem there. Then, we describe the conditions under which Turing instability occurs for each modes, and discuss the differences in pattern formation by metric graphs.

15:20~16:20 特別講演

西 慧 (京都産大 理) 3種反応拡散方程式のパルスダイナミクスについて

Kei Nishi (Kyoto Sangyo Univ.) Pulse dynamics in a three-component reaction-diffusion system

概要 The pulse dynamics in a three-component FitzHugh–Nagumo system is considered both numerically and analytically. The system admits a pulse solution of bistable type, which exhibits a variety of interface dynamics, not observed for the two-component FitzHugh–Nagumo systems. In this talk, we focus on the parameter regime $\alpha > 0$, $\beta < 0$, where there appear not only two types of pulse solutions, whose profiles are mutually inverted, but also multiple-pulse solutions which show collective motions such as clustered traveling pulses and an array of oscillating pulses. In order to analytically investigate these pulse behaviors, we apply the multiple scales method to the original reaction-diffusion system, and derive finite-dimensional ordinary differential equations which describe the motions of the pulse interfaces. The reduced ODEs enable us to reveal the global bifurcation structures of the pulse solutions, as well as the clustered multiple-pulse solutions, and to elucidate the mechanism for the rich variety of pulse dynamics in a clear, unified manner.

トポロジー

3月15日(水) 第Ⅲ会場

9:30~12:00

- 1 黒木慎太郎(岡山理大理) Orlik–Raymond type classification of simply connected 6-dimensional torus manifolds with vanishing odd degree cohomology 15
 Shintaro Kuroki Orlik–Raymond type classification of simply connected 6-dimensional
 (Okayama Univ. of Sci.) torus manifolds with vanishing odd degree cohomology

概要 A torus manifold is defined as an even-dimensional compact manifold with a half-dimensional torus action which has fixed points. This notion was introduced by Hattori–Masuda in 2003 as the generalization of the toric and the quasitoric manifolds. The aim of this talk is to classify simply connected 6-dimensional torus manifolds with vanishing odd-degree cohomology. It is shown that there is a one-to-one correspondence between equivariant diffeomorphism types of such manifolds and 3-valent torus graphs introduced by Maeda–Masuda–Panov. Using this correspondence and combinatorial arguments, we prove that such a manifold is equivariantly diffeomorphic to the 6-dimensional sphere or an equivariant connected sum of copies of 6-dimensional quasitoric manifolds or 4-dimensional sphere bundles over the 2-dimensional sphere.

- 2 那須正和 The limits of resolving directions of automorphisms of sofic systems · · 15
 Masakazu Nasu The limits of resolving directions of automorphisms of sofic systems

概要 We present the results that an automorphism of a sofic system has the same limits of resolving directions as those of its canonical support by way of the Krieger cover and that a similar result to this holds for an automorphism of a transitive sofic system and its canonical support by way of the Fischer cover. One of the corollaries to these proves that no automorphism of a transitive sofic system has an irrational unique non-expansive direction.

- 3 白根竹人(徳島大理工) Splitting invariants and a π_1 -equivalent Zariski pair of conic-line ar-
 M. Amram rangements 15
 (Shamoon Coll. of Eng.)
 坂内真三(岡山理大理)
 U. Sinichkin (Tel Aviv Univ.)
 徳永浩雄(都立大理)
 Taketo Shirane (Tokushima Univ.) Splitting invariants and a π_1 -equivalent Zariski pair of conic-line ar-
 Meirav Amram (Shamoon Coll. of Eng.) rangements
 Shinzo Bannai (Okayama Univ. of Sci.)
 Uriel Sinichkin (Tel Aviv Univ.)
 Hiro-o Tokunaga (Tokyo Metro. Univ.)

概要 In this talk, we consider the embedded topology of plane curves on the complex projective plane \mathbb{P}^2 . It is known that there are π_1 -equivalent Zariski pairs $(\mathcal{C}_1, \mathcal{C}_2)$ of plane curves $\mathcal{C}_1, \mathcal{C}_2 \subset \mathbb{P}^2$, i.e., \mathcal{C}_1 and \mathcal{C}_2 have the same embedded topology in their tubular neighborhoods but different embedded topology in \mathbb{P}^2 , and $\pi_1(\mathbb{P}^2 \setminus \mathcal{C}_1) \cong \pi_1(\mathbb{P}^2 \setminus \mathcal{C}_2)$. Known examples of π_1 -equivalent Zariski pairs consist of curves containing a component with either singularities or genus $g \geq 1$. The aim of this talk is to give a π_1 -equivalent Zariski pair of conic-line arrangements, i.e., a pair of curves consisting of smooth rational curves.

- 4 山崎 薫里 (高崎経大経済) Extensions of continuous increasing functions 10
 Kaori Yamazaki Extensions of continuous increasing functions
 (Takasaki City Univ. of Econ.)

概要 In this talk, we study extensions of continuous increasing functions, and provide an increasing version of Gillman–Jerison theorem.

- 5 深沢 尚希 (都立大理) Persistent cohomology と cup length function の計算 15
 Naoki Fukasawa (Tokyo Metro. Univ.) Persistent cohomology and computation of cup length function

概要 We introduce the structure theorem of the persistent cohomology to describe the representative cocycles in detail. Then, taking cup products of representative cocycles, we can compute the cup length function of a finite metric space.

- 6 田嶋 優 (北大理) グラフのマグニチュードホモトピー型と Mayer–Vietoris 型定理 15
 吉永 正彦 (阪大理)
 Yu Tajima (Hokkaido Univ.) Magnitude homotopy type of graphs and Mayer–Vietoris type theorem
 Masahiko Yoshinaga (Osaka Univ.)

概要 Magnitude homology is defined by Hepworth and Willerton as a categorification of the magnitude (an invariant for metric spaces). They proved the Kunneth formula and Mayer–Vietoris type theorem for magnitude homology of graphs. Recently, Asao–Izumihara introduced the pair of simplicial complexes (K, K') whose homology groups are isomorphic to direct summands of graph magnitude homology groups. We call the CW complex K/K' magnitude homotopy type. In this talk, we give a new proof of Mayer–Vietoris type theorem using Discrete Morse theory on magnitude homotopy types. If time permits, we also talk about generalizations to metric spaces.

- 7 原子 秀一 (東大数理) The modular class of an orientable ρ -Q-manifold 15
 Shuichi Harako (Univ. of Tokyo) The modular class of an orientable ρ -Q-manifold

概要 As one generalization of Lie algebras and commutative algebras, we have ρ -Lie algebras and ρ -commutative algebras, respectively. We will introduce a concept of a ρ -manifold, whose functional algebra is locally isomorphic to a ρ -commutative algebra, and discuss its modular class where the ρ -manifold is orientable and equipped with a Q-structure.

- 8 若月 駿 (名大多元数理)^b Independence complexes of grid graphs 10
 松下 尚弘 (琉球大理)
 Shun Wakatsuki (Nagoya Univ.) Independence complexes of grid graphs
 Takahiro Matsushita
 (Univ. of Ryukyus)

概要 We determine the homotopy types of the independence complexes of $(n \times 4)$, $(n \times 5)$ and $(n \times 6)$ -square grid graphs. In fact, they are homotopy equivalent to wedges of spheres.

14:20~15:20 特別講演

小川 竜 (東海大理) 凸シンプレクティック多様体の Liouville 構造と Weinstein 構造
Noboru Ogawa (Tokai Univ.) Liouville and Weinstein structures on convex symplectic manifolds

概要 In symplectic geometry, there are two important classes on convexity: Liouville structures and Weinstein structures. The former is a symplectic manifold equipped with a conformally expanding symplectic vector field which is pointing outward. The latter is imposed so that the vector field is gradient-like. Such structures have played a significant role in the study of symplectic and contact topology. For example, the convexity provides a constraint on the behavior of pseudo-holomorphic curves by the maximum principle. By definition, a Weinstein structure is a Liouville structure, but the converse is not true in general; McDuff gave the first example which is a Liouville manifold without any Weinstein structures. The purpose of this talk is to present recent advances on the difference between the two structures, up to homotopy. In particular, I will show that the stabilization of McDuff's example admits a flexible Weinstein structure. The main part is based on a joint work with Yakov Eliashberg (Stanford University) and Toru Yoshiyasu (Kyoto University of Education). If time permits, I would like to discuss some open questions, progress and perspective.

15:40~17:30

- 9 知念直紹 (防衛大) Isometry groups of symmetric products of $(\mathbb{R}^k)_p^q$ 15
Naotsugu Chinen Isometry groups of symmetric products of $(\mathbb{R}^k)_p^q$
(Nat. Defense Acad. of Japan)

概要 By $F_n(X)$, $n \geq 1$, we denote the n -th symmetric product of a metric space (X, d) as the space of the nonempty finite subsets of X with at most n elements endowed with the Hausdorff metric d_H . By $\text{Iso}(X)$ we denote the topological group of all isometries from X onto itself with the topology of pointwise convergence. We are interested in what topological group $\text{Iso}(F_n(X))$ is. The aim is to decide a topological group structure of $\text{Iso}(F_n((\mathbb{R}^k)_p^q))$ for any $(n, k, p, q) \in \mathbb{N} \times \mathbb{N} \times (\mathbb{N} \cup \{\infty\}) \times (\mathbb{N} \cup \{\infty\})$.

- 10 森 淳秀 (大阪歯大) 多様体の相互 Bayes 学習 15
Atsuhide Mori (Osaka Dent. Univ.) Mutual Bayesian learning for manifolds

概要 In previous work, the author described the geometry of Bayesian learning on a manifold. In this talk, we take the square of any manifold and a positive function which takes the maximum 1 on the diagonal set. Then we define the mutual Bayesian learning between the factors of the square. Particularly, we consider mutual learning between two copies of the parameter space of a statistical model, and associate the above function with the relative entropy. Although the mutual learning forgets all about the model except the relative entropy, it still substitutes for the usual Bayesian estimation of the parameter in a certain case.

- 11 狩野隼輔 (東北大RACMaS) Unbounded \mathfrak{sl}_3 -laminations and their shear coordinates 15
石橋 典 (東北大理)
Shunsuke Kano (Tohoku Univ.) Unbounded \mathfrak{sl}_3 -laminations and their shear coordinates
Tsukasa Ishibashi (Tohoku Univ.)

概要 Generalizing the work of Fock–Goncharov on rational unbounded laminations, we give a geometric model of the tropical points of the moduli space $\mathcal{X}_{PGL_3, \Sigma}$ of framed PGL_3 -local systems on a marked surface Σ based on the Kuperberg's \mathfrak{sl}_3 -webs. We introduce their tropical cluster coordinates as an \mathfrak{sl}_3 -analogue of the Thurston's shear coordinates associated with any ideal triangulation. We also describe tropical points of Goncharov–Shen's moduli space $\mathcal{P}_{PGL_3, \Sigma}$. Then we give a tropical analogue of gluing morphisms among the moduli spaces as a geometric gluing procedure of \mathfrak{sl}_3 -laminations with “shearings”. We also investigate a relation to the graphical basis of the \mathfrak{sl}_3 -skein algebra by Ishibashi–Yuasa, which conjecturally leads to a quantum duality map.

- 12 越野 克久 (神奈川大工) A characterization of compact sets in L^p -spaces and its application ... 15
 Katsuhisa Koshino (Kanagawa Univ.) A characterization of compact sets in L^p -spaces and its application

概要 In this talk, we give a characterization of compact sets in L^p -spaces on metric measure spaces. As its application, we investigate the topological types of the subspace consisting of lipschitz functions with bounded supports.

- 13 足立 真訓 (静岡大理) Harmonic measures and rigidity for surface group actions on the circle
 松田 能文 (青学大理工) 15
 野沢 啓 (立命館大理工)
 Masanori Adachi (Shizuoka Univ.) Harmonic measures and rigidity for surface group actions on the circle
 Yoshifumi Matsuda
 (Aoyama Gakuin Univ.)
 Hiraku Nozawa (Ritsumeikan Univ.)

概要 We study rigidity properties of surface group actions on the circle via foliated harmonic measures on the suspension bundles. We show a curvature estimate and a Gauss–Bonnet formula for an S^1 -connection obtained by taking the average of the flat connection on the suspension bundle with respect to a harmonic measure. As consequences, we give a precise description of the harmonic measure on suspension foliations with maximal Euler number and an alternative proof of rigidity theorems of Matsumoto and Burger–Iozzi–Wienhard.

3月16日(木) 第Ⅲ会場

9:30~12:00

- 14 山本 卓宏 (東京学大教育) 連続写像のファイバー特異点 15
 Takahiro Yamamoto Fiber singularities of continuous maps
 (Tokyo Gakugei Univ.)

概要 The notion of fiber-singularities of continuous maps between topological manifolds are introduced. Then, we investigate some examples of fiber-singularities and, study fiber-singularities of simplicial maps of PL maps of surfaces.

- 15 濱田 直希 (KLab(株)) 多目的最適化問題における制約関数芽および実行可能領域芽の分類と認識
 早野 健太 (慶大理工) 15
 寺本 央 (関西大システム理工)
 Naoki Hamada (KLab Inc.) Classification and recognition of constraint function-germs and feasible
 Kenta Hayano (Keio Univ.) set-germs in multiobjective optimization
 Hiroshi Teramoto (Kansai Univ.)

概要 For functions g_1, \dots, g_q and h_1, \dots, h_r on the Euclidian space, we consider the minimizing problems of some given functions on the set $M(g, h) = \{x \mid \forall i, g_i(x) \leq 0, \forall j, h_j(x) = 0\}$. The functions g_i 's, h_j 's and the set $M(g, h)$ are respectively called the constraint functions and the feasible set for the problem. In this talk, we will explain the results on classification and recognition of constraint function-germs under suitable equivalence for them (called $\mathcal{K}[G]$ -equivalence). If time permits, we will also discuss how the results are related with classification of feasible set-germs.

- 16 久野恵理香 (阪大 理) 向き付け不可能曲面の非分離曲線グラフの一樣双曲性 10
 Erika Kuno (Osaka Univ.) Uniform hyperbolicity of nonseparating curve graphs of nonorientable surface

概要 Let N be a connected nonorientable surface with or without boundary components and punctures. We prove that the graph consists of the isotopy classes of nonseparating curves of N is connected and Gromov hyperbolic with a constant which does not depend on the topological type of the surface by using the bicorn curves introduced by Przytycki and Sisto. The proof is based on the argument by Rasmussen on the uniform hyperbolicity for the graphs consists of the isotopy classes of nonseparating curves for orientable surfaces.

- 17 久野恵理香 (阪大 理) 向き付け不可能曲面のファイン曲線グラフの一樣双曲性 10
 木村満晃 (京大 理)
 Erika Kuno (Osaka Univ.) Uniform hyperbolicity of fine curve graphs of nonorientable surfaces
 Mitsuaki Kimura (Kyoto Univ.)

概要 A fine curve graph defined by Bowden, Hensel, and Webb (J. Amer. Math. Soc. (2021)) is a new curve graph consists of all essential simple closed curves on a surface. They proved that the fine curve graph of any closed orientable surface of genus $g \geq 1$ is uniformly hyperbolic in the sense of Gromov. We proved that the fine curve graph of any closed nonorientable surface of genus $g \geq 2$ is also uniformly hyperbolic.

- 18 小林竜馬 (石川工高専) 向き付け不可能閉曲面のレベル 2 写像類群による非分離単純閉曲線への
 井本奈緒 作用 10
 Ryoma Kobayashi Action on non-separating simple closed curves by the level 2 mapping
 (Ishikawa Nat. Coll. of Tech.) class group of a non-orientable closed surface
 Nao Imoto

概要 The level 2 mapping class group $\mathcal{M}_2(N)$ of a non-orientable closed surface N is the subgroup of the mapping class group of N consisting of elements acting trivially on $H_1(N; \mathbb{Z}/2\mathbb{Z})$. In our work, we proved that for any non-separating simple closed curves α and β of N , there is $\varphi \in \mathcal{M}_2(N)$ such that $\varphi(\alpha) = \beta$ if and only if α and β are equal as elements in $H_1(N; \mathbb{Z}/2\mathbb{Z})$.

- 19 小林竜馬 (石川工高専) 向き付け不可能閉曲面の非分離単純閉曲線に沿った Dehn ツイストの 2 乗
 井本奈緒 が生成する群 15
 Ryoma Kobayashi The group generated by squares of Dehn twists about non-separating
 (Ishikawa Nat. Coll. of Tech.) simple closed curves of a non-orientable closed surface
 Nao Imoto

概要 The level 2 mapping class group of an orientable closed surface is equal to the group generated by squares of Dehn twists about non-separating simple closed curves. In our work, we proved that the twist subgroup of the level 2 mapping class group of a non-orientable closed surface is not equal to the group generated by squares of Dehn twists about non-separating simple closed curves and Dehn twists about separating simple closed curves. As an application, we gave a finite generating set for this group.

- 20 大森源城 (東京理大理工) A finite presentation for the balanced superelliptic handlebody group
 15
 Genki Omori (Tokyo Univ. of Sci.) A finite presentation for the balanced superelliptic handlebody group

概要 The balanced superelliptic handlebody group is the normalizer of the transformation group of the balanced superelliptic covering space in the handlebody group of the total space. We give a finite presentation for the balanced superelliptic handlebody group. To give this presentation, we construct a finite presentation for the liftable Hilden group.

- 21 大森源城 (東京理大理工) A small generating set for the balanced superelliptic handlebody group
 10
 Genki Omori (Tokyo Univ. of Sci.) A small generating set for the balanced superelliptic handlebody group

概要 The balanced superelliptic handlebody group is the normalizer of the transformation group of the balanced superelliptic covering space in the handlebody group of the total space. We prove that the balanced superelliptic mapping class group is generated by four elements. To prove this, we also proved that the liftable Hilden group is generated by three elements. This generating set for the liftable Hilden group is minimal except for some hyperelliptic cases and the generating set for the balanced superelliptic mapping class group above is also minimal for several cases.

13:00~14:00

- 22 北野晃朗 (創価大理工) Morse–Smale flow を用いた Reidemeister torsion のトーラス和公式につ
 清水達郎 10
 (東京電機大システムデザイン工)
 Teruaki Kitano (Soka Univ.) On the torus sum formula of Reidemeister torsion defined by Morse–
 Tatsuro Shimizu (Tokyo Denki Univ.) Smale flows

概要 We consider Reidemeister torsion for a 3-manifold. It can be defined by several ways. For a 3-manifold with a torus decomposition, the torus sum formula of Reidemeister torsion is well known. In this talk we give a proof of the torus sum formula by using Morse–Smale flows.

- 23 寺垣内政一 (広島大教育) 双曲的 L 空間結び目の Upsilon 不変量について 10
 Masakazu Teragaito (Hiroshima Univ.) Hyperbolic L-space knots and their Upsilon invariants

概要 For a knot in the 3-sphere, the Upsilon invariant is a piecewise linear function defined on the interval $[0, 2]$. It is known that for an L-space knot, the Upsilon invariant is determined only by the Alexander polynomial. We exhibit infinitely many pairs of hyperbolic L-space knots such that two knots of each pair have distinct Alexander polynomials, so they are not concordant, but share the same non-zero Upsilon invariant. Conversely, we examine the restorability of the Alexander polynomial of an L-space knot from the Upsilon invariant through the Legendre–Fenchel transformation.

- 24 門上晃久 (金沢大理工) The Ma–Qiu index and the Nakanishi index for a fibered knot are equal,
 and ω -solvability 15
 Teruhisa Kadokami (Kanazawa Univ.) The Ma–Qiu index and the Nakanishi index for a fibered knot are equal,
 and ω -solvability

概要 For a knot K , let $G(K)$ be the knot group of K , $a(K)$ the Ma–Qiu index, which is the minimal number of normal generators of the commutator subgroup of $G(K)$, and $m(K)$ the Nakanishi index of K , which is the minimal number of generators of the Alexander module of K . We generalize the notions for a pair of a group G and its normal subgroup N , and we denote them by $a(G, N)$ and $m(G, N)$ respectively. Then it is easy to see $m(G, N) \leq a(G, N)$ in general. We also introduce a notion “ ω -solvability” for a group. Our main theorem is that if N is ω -solvable, then we have $m(G, N) = a(G, N)$. As corollaries, for a fibered knot K , we have $m(K) = a(K)$, and we could determine the MQ indices of prime knots up to 9 crossings completely.

3月17日(金) 第Ⅲ会場

9:30~12:00

- 25 松土恵理 (日大文理自然研) Minimum numbers of Dehn colors and \mathcal{R} -palette graphs 10
 大城佳奈子 (上智大理工)
 山岸凱司 (上智大理工)
Eri Matsudo (Nihon Univ.) Minimum numbers of Dehn colors and \mathcal{R} -palette graphs
 Kanako Oshiro (Sophia Univ.)
 Gaishi Yamagishi (Sophia Univ.)

概要 We consider the number of colors for Dehn-colorings, and we define a \mathcal{R} -palette graphs. In this talk, for any Dehn p -colorable knot, we give a lower bound of the minimum number of colors on diagrams.

- 26 市原一裕 (日大文理) 絡み目群の全射二面体群表現と二相彩色について 10
 石川勝巳 (京大数理研)
 鈴木正明 (明大総合数理)
 松土恵理 (日大文理自然研)
Kazuhiro Ichihara (Nihon Univ.) Two-tone colorings and surjective dihedral representations for links
 Katsumi Ishikawa (Kyoto Univ.)
 Masaaki Suzuki (Meiji Univ.)
 Eri Matsudo (Nihon Univ.)

概要 It is well-known that a knot group admits a surjective homomorphism to the dihedral group of degree n if and only if the knot is Fox n -colorable. However, it is not true for links with two or more components. In this talk, I will introduce a two-tone coloring on a link diagram, and give a condition for links that the link groups admit surjective representations to dihedral groups.

- 27 野坂武史 (東工大理工) 閉3次元多様体の対称ラックコサイクル不変量 10
 Takefumi Nosaka (Tokyo Tech) Symmetric rack cocycle invariants of closed 3-manifolds

概要 We establish a new approach to obtain 3-manifold invariants via Dehn surgery. For this, we introduce racks with good involution and Property FR, and define cocycle invariants as 3-manifold invariants.

- 28 松崎尚作 空間曲面と多重群ラックコサイクル不変量 10
 (足利大共通教育センター)
村尾智 (早大GEC)
 Shosaku Matsuzaki Spatial surfaces and multiple group rack cocycle invariants
 (Int. Affairs Ashikaga Univ.)
Tomo Murao (Waseda Univ.)

概要 A spatial surface is a compact surface embedded in the 3-sphere. A multiple group rack is an algebraic structure motivated from Reidemeister moves of oriented spatial surfaces. In this talk, we introduce a multiple group rack cocycle invariant for oriented spatial surfaces and give some calculation examples.

- 29 谷口雄大 (阪大理工) 結び目の n -カンドルの中心拡大 15
 田中心 (東京学大教育)
Yuta Taniguchi (Osaka Univ.) A central extension of the n -quandle of a knot
 Kokoro Tanaka (Tokyo Gakugei Univ.)

概要 The fundamental quandle is a powerful invariant for knots. However, it is difficult to compare them. Then, we often use a certain quotient of the fundamental quandle of a knot, which is called an n -quandle of a knot. In this talk, we show that the fundamental quandle of the n -twist spun knot, which is a 2-knot introduced by Zeeman, is a central extension of the n -quandle. As a corollary, we determine the second quandle homology group of n -quandles.

- 30 田中 心 (東京学大教育) 二次元結び目の結び目カンドルと結び目群について 10
 谷口 雄大 (阪大 理)
 Kokoro Tanaka (Tokyo Gakugei Univ.) Knot quandles and knot groups for 2-knots
 Yuta Taniguchi (Osaka Univ.)

概要 We give examples of 2-knots with the same knot group but different knot quandles by analyzing the knot quandles of twist-spun knots. As a byproduct of our method, we give a classification of all twist-spun knots with finite knot quandles.

- 31 和田 康載 (神戸大 理) ブルンの性質をもつリボン 2 次元ストリング絡み目の有限型不変量 10
 Kodai Wada (Kobe Univ.) Finite type invariants of ribbon 2-dimensional string links with generalized Brunnian properties

概要 Gusarov and Habiro independently gave a topological interpretation of finite type invariants for 1-dimensional knots in terms of C_k -equivalence. As a 4-dimensional analogue, Watanabe introduced the notion of RC_k -equivalence on ribbon knotted surfaces and proved that two ribbon 2-dimensional knots are RC_k -equivalent if and only if they cannot be distinguished by any finite type invariant of degree less than k . In this talk, we provide an extension of the above result of Watanabe to ribbon 2-dimensional string links with generalized Brunnian properties.

14:20~15:20 特別講演

- 石井 敦 (筑波大数理物質) Alexander type invariants of groups, quandles and MCQs
 Atsushi Ishii (Univ. of Tsukuba) Alexander type invariants of groups, quandles and MCQs

概要 Groups, quandles and MCQs are algebraic objects appeared in low-dimensional topology. In particular, the notions of quandles and multiple conjugation quandles (MCQs) were introduced via the axioms corresponding to the Reidemeister moves for oriented knots and handlebody-knots, respectively. In this talk, we discuss Alexander type invariants of groups, quandles and MCQs, which include the Alexander ideal and the (twisted) Alexander polynomial.

15:40~17:00

- 32 若槇 洋平 (阪大 情報) いくつかの有理曲面のコルク 15
 Yohei Wakamaki (Osaka Univ.) Corks of some rational surfaces

概要 Corks of 4-manifolds are one of the most important objects in the study of exotic 4-manifolds. Nevertheless, we only have a few explicit examples of corks of standard simply-connected closed 4-manifolds with small second Betti numbers b_2 . In this talk, we give the first example of a cork of $\mathbb{C}P^2 \# 8\overline{\mathbb{C}P^2}$. This achieves the smallest b_2 where a cork of standard simply-connected closed 4-manifold has been found. We also talk about new examples of corks of $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P^2}$.

- 33 浅野喜敬 (津山工高専) Some lower bounds for the Kirby–Thompson invariant of 4-manifolds
 直江央寛 (中大理工) 15
 小川将輝 (埼玉大理工)

Nobutaka Asano (Tsuyama Nat. Coll. of Tech.) Some lower bounds for the Kirby–Thompson invariant of 4-manifolds
 Hironobu Naoe (Chuo Univ.)
 Masaki Ogawa (Saitama Univ.)

概要 A trisection is a decomposition of a 4-manifold into three 1-handlebodies. This decomposition is characterized by a diagram called a trisection diagram. A trisection diagram is a 4-tuple consisting of a closed surface and three cut systems. Kirby and Thompson introduced an invariant of a 4-manifold that is a kind of complexity defined by its trisection diagrams. This invariant is called the Kirby–Thompson invariant. In this talk, we give some lower bounds for the Kirby–Thompson invariant of certain 4-manifolds. As an application, we find the first example of a 4-manifold with non-trivial Kirby–Thompson invariant.

- 34 磯島司 (東工大) Gluck twist により得られる 4次元球面の trisection 15
 小川将輝 (埼玉大理工)

Tsukasa Isoshima (Tokyo Tech) Trisections of the 4-sphere obtained by Gluck twisting
 Masaki Ogawa (Saitama Univ.)

概要 A trisection of a 4-manifold, introduced by Gay and Kirby in 2012, is a decomposition of the 4-manifold into three 4-dimensional 1-handlebodies. A trisection can be described by a trisection diagram that is similar to a Heegaard diagram. In this talk, we construct infinitely many trisection diagrams of the 4-sphere obtained by Gluck twisting along some spun torus knots. We also study the standardness of the trisection diagrams.

- 35 大場貴裕 (阪大) Symplectic submanifolds in dimension 6 from Lefschetz fibrations 15
 Takahiro Oba (Osaka Univ.) Symplectic submanifolds in dimension 6 from Lefschetz fibrations

概要 Thanks to the affirmative answer to the symplectic Thom conjecture, all connected symplectic surfaces representing the same homology class in a closed symplectic 4-manifold are diffeomorphic. In higher dimension, for example, take $\mathbb{C}P^3$ with the Fubini–Study form. Then, it is known that its connected symplectic submanifolds representing the class $d[\mathbb{C}P^2]$ are mutually diffeomorphic as long as $1 \leq d \leq 3$. In this talk, I will present arbitrarily (finitely) many codimension 2 connected symplectic submanifolds in a closed symplectic 6-manifold which are mutually homotopy inequivalent but homologous. A key ingredient of the construction is hyperelliptic Lefschetz fibrations.

無限可積分系

3月17日(金) 第IX会場

9:45~10:45

- 1 上野 祐一 (皇學館大教育・神戸大理) Polynomial Hamiltonians for quantum Garnier system in two variables 10
 Yuichi Ueno (Kogakkan Univ./Kobe Univ.) Polynomial Hamiltonians for quantum Garnier system in two variables

概要 Garnier system is the extension of Painlevé equations and is given as Frobenius integrable multi-time Hamiltonian systems. In this talk, we construct and characterize the quantum Garnier system by certain holomorphic properties. The characterization of the classical Garnier system by holomorphic properties has been given by Sasano, and our result is a quantization of that work. In other words, we introduce quantum canonical transformations such that the Hamiltonian system of the Garnier system is again transformed into a polynomial Hamiltonian system, and show that the Hamiltonian can be uniquely characterized by this holomorphic properties.

- 2 中園 信孝 (東京農工大工) 離散 KdV 方程式の立方体周りの整合性 15
 Nobutaka Nakazono Consistency around a cube property of Hirota's discrete KdV equation
 (Tokyo Univ. of Agri. and Tech.)

概要 It has been unknown whether Hirota's discrete analogue of the Korteweg-de Vries (KdV) equation have the consistency around a cube (CAC) property, which is known as an integrability of 2-dimensional difference equation. In this talk, we show that Hirota's discrete KdV equation has the CAC property.

- 3 鈴木 貴雄 (近畿大理工) q -ガルニエ系に由来する q -超幾何級数へのアフィン・ワイル群作用 15
 Takao Suzuki (Kindai Univ.) An affine Weyl group action on the basic hypergeometric series arising from the q -Garnier system

概要 Recently, we formulated the q -Garnier system and its variations as translations of an extended affine Weyl group of type $A_{2n+1}^{(1)} \times A_1^{(1)} \times A_1^{(1)}$. On the other hand, those systems admit particular solutions in terms of the basic hypergeometric series ${}_{n+1}\phi_n$. In this talk, we investigate an action of the extended affine Weyl group on ${}_{n+1}\phi_n$. This talk is based on a joint work with Taiki Idomoto of Kindai University.

- 4 大山 陽介 (徳島大理工) Boundary behavior of q -Painlevé equation of type $A_4^{(1)}$ 15
 Yousuke Ohyama (Tokushima Univ.) Boundary behavior of q -Painlevé equation of type $A_4^{(1)}$

概要 We study asymptotic analysis on the q -Painleve V equation around the origin. We solve the q -Riemann-Hilbert problem for any asymptotic series of q -Painleve V functions. We show the Mano-type decomposition for q -Painleve V.

11:00~12:00 特別講演

井上 玲 (千葉大理) ワイル群のクラスター実現と表現論への応用
 Rei Inoue (Chiba Univ.) Cluster realization of Weyl groups and its applications to representation theory

概要 The cluster algebra is a commutative algebra introduced by Fomin and Zelevinsky around 2000. The characteristic operation in the algebra called ‘mutation’ is related to various notions in mathematics and mathematical physics.

In this talk we introduce a realization of Weyl groups in terms of cluster mutations, for finite dimensional semisimple Lie algebras. We discuss its applications to (1) the geometric R-matrix of symmetric power representations for the affine quantum algebra of A -type, and to (2) the q -characters of quantum non-twisted affine algebras introduced by Frenkel and Reshetikhin, when q is a root of unity. If time permits we also mention the application to higher Teichmüller theory.

This talk is based on joint works with Thomas Lam, Pavlo Pylyavskyy, Tsukasa Ishibashi, Hironori Oya, and Takao Yamazaki.

3月18日(土) 第IX会場

10:00~10:45

5 小見山 尚 (名大多元数理) Associators and the Grothendieck–Teichmüller group in mould theory 10

Nao Komiyama (Nagoya Univ.) Associators and the Grothendieck–Teichmüller group in mould theory

概要 In this talk, we explain the mould theory introduced by Jean Ecalle, and we introduce mould theoretic formulations of Drinfeld’s notion of the associator set and that of the Grothendieck–Teichmüller group. This talk is based on the joint work with Hidekazu Furusho and Minoru Hirose.

6 国場 敦夫 (東大総合文化) New solutions to the tetrahedron equation associated with quantized
 松生 周一郎 (東大総合文化) six-vertex models 15

米山 瑛仁 (東大総合文化)

Atsuo Kuniba (Univ. of Tokyo) New solutions to the tetrahedron equation associated with quantized
 Shuichiro Matsuike (Univ. of Tokyo) six-vertex models

Akihito Yoneyama (Univ. of Tokyo)

概要 In this talk, we present a family of new solutions to the tetrahedron equation of the form $RLLL = LLLR$, where L operator may be regarded as a quantized six-vertex model whose Boltzmann weights are specific representations of the q -oscillator or q -Weyl algebras. When the three L ’s are associated with the q -oscillator algebra, R coincides with the known intertwiner of the quantized coordinate ring $A_q(sl_3)$. On the other hand, L ’s based on the q -Weyl algebra lead to new R ’s whose elements are either factorized or expressed as a terminating q -hypergeometric type series.

7 中島 俊樹 (上智大理工) Geometric crystal on unipotent variety and crystal of modified quantum algebra 15

Toshiki Nakashima (Sophia Univ.) Geometric crystal on unipotent variety and crystal of modified quantum algebra

概要 We shall define a geometric crystal structure on the unipotent variety of a simple, simply connected complex algebraic group. In $SL(2)$ -case, certain relation to crystal base of a modified quantum algebra will be presented.

11:00~12:00 特別講演

T. Scrimshaw (北大教育推進機構) Krystal theory

Travis Scrimshaw (Hokkaido Univ.) Krystal theory

概要 Kashiwara's theory of crystal bases has been applied to many different problems in representation theory, physics, and geometry. In this talk, we will focus on their appearance in the K-theoretic Schubert calculus of the Grassmannian. Specifically, we look at the symmetric function representatives of K-theory classes that arise from Schubert varieties called (symmetric) Grothendieck polynomials $\{G_\lambda\}_\lambda$ and three other related bases: the dual basis and their "weak" analogs. We construct a \mathfrak{gl}_n -crystal using the combinatorial descriptions of Buch and Lam–Pylyavskyy to describe the decomposition of these bases into Schur functions and describe their multiplicities. We will then discuss extensions to the (refined) canonical Grothendieck polynomials introduced by Hwang *et al.* and Yeliussizov. This suggests an extension of Kashiwara's theory which we call K-crystals, or Krystals for short. We conclude with some open problems. This is based on joint works with Graham Hawkes, Cara Monical, Oliver Pechenik.