

🌸 日本数学会

2014年度秋季総合分科会

英文サマリ集

2014年9月

於 広島大学

2014 日本数学会

秋季総合分科会プログラム

期 日 2014年9月25日(木)～9月28日(日)
 会 場 広島大学東広島キャンパス
 〒739-8521 広島県東広島市鏡山1-7-1
 連絡先 広島大学大学院理学研究科数学教室
 〒739-8526 広島県東広島市鏡山1-3-1
 E-mail hiroshima14sept@mathsoc.jp
 (会期中) Tel 090-1791-3483
 一般社団法人 日 本 数 学 会
 Tel 03-3835-3483

会場 日時	第I会場 総合科学部 K110	第II会場 総合科学部 K108	第III会場 総合科学部 K107	第IV会場 総合科学部 L101	第V会場 総合科学部 L102	第VI会場 総合科学部 K211	第VII会場 総合科学部 K210	第VIII会場 総合科学部 L201	第IX会場 総合科学部 K314
25日 (木)	函 数 論 9:00～12:00 14:10～16:50	代 数 学 9:30～12:00 14:15～16:15	幾 何 学 9:40～12:00 14:15～16:10	函数解析学 14:15～16:35	函数方程式論 9:00～12:00 14:15～16:15	トポロジー 9:20～12:00 14:15～15:25	統計数学 9:30～11:50	応用数学 9:30～11:30 14:15～17:00	無限可積分系 10:00～12:00 14:15～15:30
	企画特別講演 13:00～14:00								
	特別講演 17:00～18:00	特別講演 16:30～17:30	特別講演 16:30～17:30	特別講演 17:00～18:00	特別講演 16:30～17:30	特別講演 15:45～16:45	特別講演 14:15～15:15 15:30～16:30		特別講演 15:45～16:45
26日 (金)	函 数 論 9:00～10:30	代 数 学 9:30～11:50	幾 何 学 10:00～11:50	函数解析学 9:00～12:00	函数方程式論 13:00～14:15	トポロジー 9:30～11:30	統計数学 9:30～11:30	応用数学 9:30～11:30	無限可積分系 10:00～12:00
	特別講演 13:20～14:20	特別講演 13:20～14:20	特別講演 10:50～11:50 13:15～14:15	特別講演 13:10～14:10	特別講演 13:15～14:15			特別講演 13:00～14:00	特別講演 13:00～14:00
	日本数学会賞授賞式(サタケメモリアルホール).....(14:50～15:20)								
	総 合 講 演 (") 日本数学会賞秋季賞受賞者.....(15:30～16:30)								
	大 鹿 健 一 (阪 大 理).....(16:45～17:45)								
懇 親 会 (西条HAKUWAホテル).....(18:15～20:15)									
27日 (土)	実函数論 9:00～11:45 14:15～16:20	代 数 学 9:30～11:25 15:30～17:30	幾 何 学 9:40～12:00 14:15～16:00	函数解析学 9:00～12:00 14:15～16:45	函数方程式論 9:00～12:00 14:15～16:15	トポロジー 9:20～11:50 14:15～15:40	統計数学 9:30～12:00	応用数学 14:15～16:30 特別セッション 9:30～11:55	数学基礎論 および歴史 9:30～11:40 14:15～15:40
	企画特別講演 13:00～14:00								
	特別講演 16:40～17:40	特別講演 14:15～15:15	特別講演 16:15～17:15	特別講演 17:00～18:00	特別講演 16:30～17:30	特別講演 16:00～17:00	特別講演 14:15～15:15 15:30～16:30	特別講演 16:45～17:45	特別講演 16:00～17:00
28日 (日)	実函数論 9:00～11:50 14:15～15:40	代 数 学 9:15～12:00			函数方程式論 9:00～12:00 14:15～16:15	トポロジー 9:20～12:00	統計数学 9:30～11:20 14:15～16:10	応用数学 9:30～11:50 14:15～16:15	数学基礎論 および歴史 9:30～11:30 15:00～16:10
	企画特別講演 13:00～14:00								
	特別講演 16:00～17:00	特別講演 14:15～15:15			特別講演 16:30～17:30			特別講演 16:30～17:30	

目 次

総合講演	1
企画特別講演	1
数学基礎論および歴史	5
9月27日(土)	5
9月28日(日)	9
代数学	12
9月25日(木)	12
9月26日(金)	17
9月27日(土)	20
9月28日(日)	23
幾何学	26
9月25日(木)	26
9月26日(金)	31
9月27日(土)	31
函数論	36
9月25日(木)	36
9月26日(金)	40
函数方程式論	43
9月25日(木)	43
9月26日(金)	49
9月27日(土)	54
9月28日(日)	60
実函数論	68
9月27日(土)	68
9月28日(日)	73

函数解析学	78
9月25日(木)	78
9月26日(金)	80
9月27日(土)	82
統計数学	87
9月25日(木)	87
9月26日(金)	90
9月27日(土)	92
9月28日(日)	96
応用数学	101
9月25日(木)	101
9月26日(金)	106
9月27日(土)	109
9月28日(日)	112
トポロジー	117
9月25日(木)	117
9月26日(金)	121
9月27日(土)	122
9月28日(日)	127
無限可積分系	130
9月25日(木)	130
9月26日(金)	132

総合講演

9月26日(金) サタケメモリアルホール

日本数学会賞秋季賞受賞者 (15:30~16:30)
MSJ Autumn Prize Winner

大鹿健一(阪大理) Klein群の幾何とその応用 (16:45~17:45)
Ken'ichi Ohshika (Osaka Univ.) Geometry of Kleinian groups and its applications

概要 The study of Kleinian groups using geometric method began with Thurston's paradigm-making work in 1980's. During a few decades after then, many specialists in this field worked to solve problems raised by Thurston. Now all of these problems are solved, and it is time to look further.

In this talk, I shall first explain what are Thurston's problems and what are their significances, and summarise how they have been solved. After that, I shall present how techniques developed in the process of solving these problems are useful in the study of three-manifolds and discrete groups, taking results I have recently obtained as examples.

企画特別講演

9月25日(木)

第II会場

Ming-Lun Hsieh (Nat. Taiwan Univ.)^b Modular forms and Iwasawa theory (13:00~14:00)
Ming-Lun Hsieh (Nat. Taiwan Univ.) Modular forms and Iwasawa theory

概要 One of the most intriguing formulas in algebraic number theory is Dirichlet class number formula, which relates the analytic objects: special L-values and algebraic objects: class numbers and Selmer groups. Many outstanding conjectures such as Bloch-Kato conjecture, Birch and Swinnerton-Dyer conjecture and Iwasawa main conjectures all share the same flavour. Number theorists have been using modular forms to study analytic or algebraic properties of special L-values by various period integral formulas and to study associated Galois representations and Selmer groups via the development of Langlands correspondence. One effective way to make a connection between special L-values and Selmer groups is via the method of congruence among modular forms. In this talk, we would like to talk about successful examples of this method as well as the recent progress on Iwasawa main conjectures and the application to Birch and Swinnerton-Dyer conjecture.

第V会場

- 小 俣 正 朗 (金 沢 大 理 工) 液 滴 ・ 泡 の 数 理 …………… (13:00~14:00)
 Seiro Omata (Kanazawa Univ.) Mathematical and numerical analysis of a droplet and bubble
 motion

概要 In this talk we discuss the motion of droplets or bubbles on a flat surface. This surface is represented by a plane, which serves as an obstacle. The motion is controlled mainly by the surface tension and the contact angle between the plane and the surface of the droplet. We introduce the governing equations for the evolution of the droplet surface. The equations are of wave type (hyperbolic type) with an obstacle and a free boundary. Since the droplet should keep its volume throughout the evolution, we need to impose a volume-preserving constraint. To establish an existence theorem, we approximate the governing equations by a discrete Morse flow.

Furthermore, we present an algorithm describing a motion of multiple bubbles, based on the so-called BMO algorithm. We modify the BMO algorithm in order to handle multiphase mean curvature or mean curvature acceleration problems under a volume constraint.

9月27日(土)

第II会場

- 柏 原 正 樹 (京 大 数 理 研) Riemann Hilbert correspondence for irregular holonomic D-
 modules …………… (13:00~14:00)
 Masaki Kashiwara (Kyoto Univ.) Riemann Hilbert correspondence for irregular holonomic D-
 modules

概要 The original Riemann-Hilbert problem is to construct a linear ordinary differential equation with regular singularities whose solutions have a given monodromy. Nowadays, it is formulated as a categorical equivalence between the category of regular holonomic D-modules and the category of perverse sheaves. However it is a long standing problem to describe holonomic D-modules with irregular singularities in a geometric or topological language.

Recently, I, with Andrea D'Agnolo, proved a Riemann-Hilbert correspondence for holonomic D-modules which are not necessarily regular (arXiv:1311.2374). In this correspondence, we have to replace the derived category of constructible sheaves with the enhanced sheaves category. This category is a quotient of the derived category of ind-sheaves (or subanalytic sheaves) on the product of the base space and the real projective line. Then we construct a fully faithful functor from the derived category of holonomic D-modules to the enhanced sheaves category.

第Ⅲ会場

河野俊丈(東大数理) 反復積分と de Rham ホモトピー理論 (13:00~14:00)
 Toshihiko Kohno (Univ. of Tokyo) Iterated integrals and de Rham homotopy theory

概要 The notation of iterated integrals of differential forms was initiated by K.-T. Chen in 1970's. The theory of iterated integrals provides a method to describe the de Rham cohomology of the loop spaces of simply connected manifolds. It also enables us to extract information about the nilpotent completion of the fundamental groups. The theory of iterated integrals is closely related to the rational homotopy theory due to D. Quillen and the de Rham homotopy theory due to D. Sullivan.

After reviewing these historical aspects, I will focus on more recent developments concerning the theory of iterated integrals. First, I describe the Kontsevich integral which gives a universal expression for finite type invariants of knots in terms of iterated integrals of logarithmic forms. The Kontsevich integral is relevant to the Drinfel'd associator and multiple zeta values as well.

There is also a new aspect for the algebraic structure of the homology of free loop spaces in the framework of string topology. Such structures are related to iterated integrals by means of Hochschild homology. I describe how loop products and the Batalin-Vilkovisky structure for the homology of free loop spaces are expressed in terms of iterated integrals. Finally, I will mention a description of volumes of spherical and hyperbolic simplices by iterated integrals based on the Schläfli formula.

第Ⅴ会場

古谷康雄(東海大理) コーシー積分—実解析と偏微分方程式の接点— (13:00~14:00)
 Yasuo Komori-Furuya (Tokai Univ.) Cauchy integral operator and related topics

概要 This talk is a survey for Cauchy integral operator and related topics. We recall the results by Calderón and Coifman, McIntosh and Meyer: L^p boundedness of Cauchy integral operator. We consider close relations between this problem and some problems in PDE: the Laplace equation on Lipschitz domains and Kato's square root problem.

9月28日(日)

第Ⅱ会場

島田伊知朗(広島大理) K3 surfaces and lattice theory (13:00~14:00)
 Ichiro Shimada (Hiroshima Univ.) K3 surfaces and lattice theory

概要 Thanks to the Torelli-type theorems for the period mapping, many geometric problems on K3 surfaces are reduced to computational problems in lattice theory, and the latter can often be solved by means of computer. In this talk, we explain how to use the lattice theory and computer in the study of K3 surfaces.

第Ⅲ会場

森本光生 建部賢弘生誕 350 周年 (13:00~14:00)
 (四日市大関孝和数学研・上智大*)
 Mitsuo Morimoto The 350th anniversary of Takebe Katahiro
 (Yokkaichi Univ./Sophia Univ.*)

概要 Takebe Katahiro (1664-1739) is a Japanese mathematician in the Edo era, a student and a coworker of Seki Takakazu (ca 1642-1708). Takebe's mathematical works include the *Kenki Sanpō* (1683), the *Endan Genkai* (1685) to the *Hatsubi Sanpō* (Seki Takakazu, 1674), the *Genkai Taisei* (1690) to the Chinese mathematics book *Suanxue Qimeng* (Zhu Shijie, 1299) and the *Tetsujutsu Sankei* (1722). He, together with his master and his elder brother Kata'akira, conceives the *Taisei Sankei*, an encyclopedic monograph of all mathematics known to them. The project starts in 1683 and ends in 1711 after 28 years of compilation.

To celebrate his 350th anniversary, an international conference on the "Traditional Mathematics of East Asia and Related Topics," the Takebe Conference 2014, is organized on August 25-30 at Ochanomizu University. The Takebe Conference 2014 is one of the satellite conferences of SEOUL ICM 2014 and supported by the Mathematical Society of Japan.

In this talk, we survey the mathematics of Takebe Katahiro and the events of his 350th anniversary.

数 学 基 礎 論 お よ び 歴 史

9月27日(土) 第IX会場

9:30~11:40

- 1 大 藪 卓 b Theory of H -theorems, 他5件 5
 Takashi Oyabu Theory of H -theorems, and other 5 talks

概要 1. Constructible world. All sets are constructible::Then::Conntinuum problem::General contunuum Problem is true::We stand on Kantian philosophy: $V = L$: $-problem$:

2. Theory of H -theorems. H -function in mathematics:Entropy concept in math... Then:: $dH/dt = < 0$ =====> 0 :

3. Representation theory. Representations of $Diff(M)::Aut(R)$:Poincaré conjectures hold:Real orthogonal representations:

4. Fluctuation. Fluctuation and prigogine theory: H -theorems: δ - H theorems:Gibbs free energy:::Helmholtz free energy:: $H = SdAU(\mu)$: $H = SdA\Sigma UilogUi$: $\delta - H = SdAdUlogdU$: $\delta - H = SdA\Sigma dUilogdUi$

5. Riemann surface. Taniyama-Simura conjecture:Auto-morphic functions:Modular functions:Modular curves: X =====> $E : V$::::Morphism:

6. Maximal ideal space. R =====> $SPEC(R)$:: R =====> $M(R)$: $M(R) = \{\text{Maximal ideals in } R\}$ Topology:::Topological space:Introduce topology in $M(R)$:This makes $M(R)$:::A manifold::

- 2 伊 東 由 文 (徳 島 大*) 順序数の概念の定義と存在定理 15
 Yoshifumi Ito (Univ. of Tokushima*) Definition and existence theorem of the concept of ordinal numbers

概要 In this paper we give the new definition of the concept of ordinal numbers and prove its existence theorem on the basis of the ZFC set theory. This is the generalization of Peano's system of axioms of finite ordinal numbers.

- 3 菊 池 誠 (神戸大システム情報) 不完全性定理の構成的性質について 15
 瀧 野 昌 (神戸大システム情報)
 Makoto Kikuchi (Kobe Univ.) On the constructive nature of the incompleteness theorem
 Sakaé Fuchino (Kobe Univ.)

概要 The first incompleteness theorem consists of two statements. One is that the Gödel sentence of a theory of arithmetic is unprovable if the theory is consistent, and the other is that the negation of the Gödel sentence is unprovable if the theory satisfies a certain soundness condition. We discuss the difference of the constructive nature of these two statements by interpreting their proofs in terms of the constructions of primitive primitive recursive functions.

- 4 菊池 誠 (神戸大システム情報) 算術の超準モデルにおける定理と証明について 15
 倉橋 太志 (木更津工高専)
Makoto Kikuchi (Kobe Univ.) On theorems and proofs in nonstandard models of arithmetic
Taishi Kurahashi
 (Kisarazu Nat. Coll. of Tech.)

概要 The set of theorems of arithmetic in a nonstandard model of arithmetic is equal to or is bigger than the standard set of theorems of arithmetic. We firstly show that if the set of theorems in such a model is bigger than the standard one, the set must contain true and untrue sentences. Then, we show the existence of nonstandard models which satisfy the condition that set of theorems on their consistent initial segments are always standard, and are always nonstandard. At last, we prove that there is a Rosser-type provability predicate and a nonstandard model such that the theory defined by the provability predicate in the model is true arithmetic.

- 5 田中 一之 (東北大理) Phase transitions and reverse mathematics 15
 F. Pelupessy (東北大理)
Kazuyuki Tanaka (Tohoku Univ.) Phase transitions and reverse mathematics
Florian Pelupessy (Tohoku Univ.)

概要 RM (reverse mathematics) and CMI (concrete mathematical independence) are two related areas of foundational interest which have emerged in mathematical logic. Recently, Andreas Weiermann innovated a new dichotomy principle called ‘phase transitions’ to illustrate an incompleteness phenomenon of CMI. In this talk, we are going to present some finer versions of the phase transition principle via abounding results of RM. Thus we can cross-link RM and CMI for a better understanding of foundational phenomena.

- 6 鈴木 登志雄 (首都大東京理工) Equilibrium points of an AND-OR tree: under constraints on probability 15
 仁井田 哲尚 (パテントリザルト)
Toshio Suzuki (Tokyo Metro. Univ.) Equilibrium points of an AND-OR tree: under constraints on probability
Yoshinao Niida (Patent Result Co.)

概要 We study a probability distribution d on the truth assignments to a uniform binary AND-OR tree. Liu and Tanaka [2007, Inform. Process. Lett.] showed the following: If d achieves the equilibrium among independent distributions (ID) then d is an independent identical distribution (IID). We show a stronger form of the above result. Given a real number r such that $0 < r < 1$, we consider a constraint that the probability of the root node having the value 0 is r . Our main result is the following: When we restrict ourselves to IDs satisfying this constraint, the above result of Liu and Tanaka still holds.

- 7 只木孝太郎 (中大研究開発機構) アルゴリズム的ランダムネスによる量子力学の再構成 15
 Kohtaro Tadaki (Chuo Univ.) Reformulating quantum mechanics by algorithmic randomness

概要 The notion of probability plays a crucial role in quantum mechanics. It appears through the so-called Born rule. In modern mathematics which describes quantum mechanics, however, probability theory means nothing other than measure theory, and therefore any operational characterization of the notion of probability is still missing in quantum mechanics. In this sense, the current form of quantum mechanics is considered to be imperfect as a physical theory which must stand on operational means. In this talk, we present an alternative rule to the Born rule based on algorithmic randomness without reference to the notion of probability. We use the notion of Martin-Löf randomness with respect to Bernoulli measure for specifying the property of the results of quantum measurements in an operational way. We consider the validity of the new rule, in particular, based on the many-worlds interpretation of quantum mechanics.

- 8 太田浩行 (東大情報理工) 帰納解析学における小さな計算量級 15
 河村彰星 (東大情報理工)
 Hiroyuki Ota (Univ. of Tokyo) Small complexity classes for computable analysis
 Akitoshi Kawamura (Univ. of Tokyo)

概要 Type-two Theory of Effectivity (TTE) provides a general framework for computable analysis. To refine it to polynomial-time computability while keeping as much generality as possible, Kawamura and Cook recently proposed a modification to TTE using machines that have random access to an oracle and run in time depending on the “size” of the oracle. They defined type-two analogues of P, NP, PSPACE and applied them to real functions and operators. We further refine their model and study computation below P: type-two analogues of the classes L, NC, and P-completeness under log-space reductions. As prototypical applications, we recast several facts (some in a stronger form than was known) about the complexity of numerical problems into our framework.

14:15~15:40

- 9 木原貴行 (北陸先端大情報) 無限次元ポーランド空間の σ -同相型と次数スペクトル 15
 A. Pauly (Univ. of Cambridge)
 Takayuki Kihara (JAIST) σ -homeomorphism types and point degree spectra of infinite dimensional spaces
 Arno Pauly (Univ. of Cambridge)

概要 The concept of a point degree spectrum links the study of degree structures in computability theory to the study of dimension in descriptive set theory and topology. We answer an open question whether there are more than two equivalence classes of perfect Polish spaces w.r.t. countable continuous isomorphisms in the positive.

- 10 渕野昌 (神戸大システム情報) ベール関数の概連続性 15
 Sakaé Fuchino (Kobe Univ.) Almost continuity of Baire functions

概要 Frankiewicz and Kunen (1987) proved that the assertion “for any Baire space X and any topological space Y with σ -disjoint open base, if f is a Baire function from X to Y then f is continuous on a co-meager subset of X ” is independent over ZFC and the negation of the assertion is equi-consistent with a measurable cardinal. We shall discuss some results related to this theorem.

- 11 南 裕明 (神戸大工) The dominating number of F_σ ideals on Katětov–Blass order 15
 酒井 拓史 (神戸大工) The dominating number of F_σ ideals on Katětov–Blass order
 Hiroaki Minami
 Hiroshi Sakai (Kobe Univ.)

概要 We will talk about the dominating number of F_σ ideals on Katětov–Blass order is equal to the dominating number \mathfrak{d} which is the least size of cofinal subsets of the family of functions from the natural numbers to the natural numbers ordered by the relation “almost dominating”.

- 12 阿部 吉弘 (神奈川大理) Rigidity and weakly normal ideals on $\mathcal{P}_\kappa\lambda$ 15
 Yoshihiro Abe (Kanagawa Univ.) Rigidity and weakly normal ideals on $\mathcal{P}_\kappa\lambda$

概要 On κ , selective ideals are rigid, and normal ideals selective. While on $\mathcal{P}_\kappa\lambda$, all normal ideals are not selective. We proved last year a weakly normal ideal is selective if the sup-function is one-to-one on a measure one set. We show the same assumption gives the rigidity to a weakly normal ideal.

- 13 薄葉 季路 (神戸大自然) Set-theoretic geology in HOD 15
 Toshimichi Usuba (Kobe Univ.) Set-theoretic geology in HOD

概要 Set-theoretic geology is a study of the structure of all ground models of the universe. We prove that under $V = HOD$ or a bit weaker assumption, the structure of all ground models behaves well as under $V = L[A]$.

16:00~17:00 特別講演

- 池上 大祐 (神戸大システム情報) Gödel’s program, large cardinals, and forcing axioms
 Daisuke Ikegami (Kobe Univ.) Gödel’s program, large cardinals, and forcing axioms

概要 After proving that one cannot refute the Continuum Hypothesis (CH) in Zermelo–Fraenkel set theory with the Axiom of Choice (ZFC), Gödel suspected that one cannot decide the truth-value of CH in ZFC, i.e., CH is *independent* from ZFC. To discuss the truth-values of CH and other mathematical statements, Gödel initiated the following program nowadays called **Gödel’s Program**: Decide the truth-values of mathematically interesting statements independent from ZFC in “well-justified” extensions of ZFC.

What Gödel had in mind at that time was the notion of *large cardinals*. He expected to decide the truth-value of CH by assuming the existence of a certain large cardinal in ZFC.

After Cohen introduced *forcing* which is now a basic tool to argue the unprovability of statements in set theory, it was shown by Lévy and Solovay that the existence of *any* large cardinal in ZFC cannot decide the truth-value of CH. However, using large cardinals in ZFC, one can decide the truth-values of many mathematical statements which are independent from ZFC alone, and furthermore one can obtain a clear understanding of the theory of the second-order structure $(\mathbb{N}, \mathcal{P}(\mathbb{N}), \in)$.

In this talk, we try to extend this understanding to the theory of the structure $(\omega_1, \mathcal{P}(\omega_1), \in)$ where ω_1 is the least uncountable cardinal, by using large cardinals, forcing axioms, and some hypothesis from inner model theory in ZFC. This is joint work with Matteo Viale.

9月28日(日) 第IX会場

9:30~11:30

- 14 池田宏一郎 (法政大経営) ジェネリック構造の単純性と強順序性 15
Koichiro Ikeda (Hosei Univ.) Simplicity and the strong order property of generic structures

概要 Evans and Wang proved that if an omega-categorical generic structure is not simple then it has SOP_3 . We consider a generalization of their result.

- 15 桔梗宏孝 (神戸大システム情報) On model complete generic structures 15
Hirotaka Kikyo (Kobe Univ.) On model complete generic structures

概要 We discuss the model completeness of countably categorical structures constructed by Hrushovski's amalgamation method. We obtained some positive results in the ab initio case with coefficient 1 for the predimension function and with a ternary relation. We will discuss the cases with coefficient other than 1.

- 16 小林宗広 (筑波大数理物質) NTP₂ 理論における dividing と forking の一致について –indiscernible
坪井明人 (筑波大数理物質) tree を用いた新証明– 15
Munehiro Kobayashi On the equivalence of dividing and forking in NTP₂ theories
(Univ. of Tsukuba)
Akito Tsuboi (Univ. of Tsukuba)

概要 In this talk, we see a new proof to Chernikov–Kaplan's theorem which insists the equivalence of dividing and forking over extension bases. We use indiscernible structures to prove the theorem, which are more elementary notions than the ones used in Chernikov and Kaplan's proof. The new proof was obtained by improving Tsuboi's idea.

- 17 倉田俊彦 (法政大経営) Concrete domain と層の圏論的同等性に関する考察 15
Toshihiko Kurata (Hosei Univ.) Categorical equivalence between concrete domains and sheaves

概要 We introduce a both-way translation between the category of concrete domains and a certain category of sheaves, based on which we furthermore study the equivalence of these two categories.

- 18 佐野勝彦 (北陸先端大情報) 四値論理の式計算体系のカット除去定理 15
Katsuhiko Sano (JAIST) Cut-elimination theorem for Belnap–Dunn's four-valued logic

概要 This talk concerns a proof theory for Belnap–Dunn's four-valued logic. First, we overview how we can extract Dunn's relational semantics for a given logical connective from a four-valued truth table for the connective. Second, we provide inference rules for the connective in a sequent calculus of Belnap–Dunn logic and establish an admissibility of the rule of cut in the expanded calculus with these rules.

- 19 鈴木 信行 (静岡大 理) 中間述語論理における existence property と disjunction property の独立性 15
 Nobu-Yuki Suzuki (Shizuoka Univ.) The independence of existence and disjunction properties in intermediate predicate logics

概要 An intermediate predicate logic \mathbf{L} is said to have the *existence property* (EP), if for every $\exists xA(x)$, $\mathbf{L} \vdash \exists xA(x)$ implies that there exists an individual variable v such that $\mathbf{L} \vdash A(v)$. A logic \mathbf{L} is said to have the *disjunction property* (DP), if for every A and every B , $\mathbf{L} \vdash A \vee B$ implies either $\mathbf{L} \vdash A$ or $\mathbf{L} \vdash B$. Since \exists is regarded as the generalized \vee , a natural question arises: *Does EP imply DP?* This is known as Ono's problem **P52**. We provide a negative solution to this problem. We also present an affirmative partial result under a natural condition for reasonable logics.

- 20 鹿島 亮 (東工大情報理工) Semilattice relevant logic について 15
 Ryo Kashima (Tokyo Tech) On semilattice relevant logics

概要 Urquhart's semilattice relevant logic is defined to be the set of valid formulas in semilattice models. We call it R_0 , which is slightly different from the orthodox relevant logic R . In this talk, semilattice models are extended and a new logic, called R_1 , is defined. R_1 is a proper subset of R_0 . The following formulas are elements of $R_0 - R_1$: $((p \rightarrow r) \wedge (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$, $((r \rightarrow (p \vee q)) \wedge (p \rightarrow q)) \rightarrow (r \rightarrow q)$; the former is an axiom of R , and the latter is not provable in R . We study certain properties of R_1 ; e.g., finding a sound and complete proof system.

11:30~12:00 数学基礎論および歴史分科会総会

- 14:15~14:45 歴史部門懇談会 第1部 「8代将軍吉宗に命じられ建部賢弘が指揮し作成した享保日本図について」
 久下実氏 (広島県立歴史博物館) による講演と討議

15:00~16:10

- 21 堀口 俊二 (新潟産大経済) 清水流測量術秘伝書の発見 —測量家たちの繋がり と宗教からの一考察—
 下斗米 哲明 15
 Shunzi Horiguchi A discovery of the Shimizu style surveying secret book —Considerations
 (Niigata Sangyo Univ.) from the connections of the surveyors and religions—
 Tetsuaki Shimotomai

概要 The Shimizu style surveying secret book of early stages in the Edo period (1603–1868) was found in 2007 in Sapporo. We consider the surveying from the connections of the surveyors and religions.

- 22 田中昭太郎 b 分数関数のべき級数表示法の厳格さ, 簡便さ —和田寧の定理, コーシーの積分公式, 循環級数の定理— 15
 Shotaro Tanaka Methods for power series expansions of rational functions. Strictness and handiness of their methods

概要 Example: rational function: $f(z) = (4z^2 - 7z + 11)/(2 + 5z)(2z - 3)^2 = 1/(2 + 5z) + 1/(2z - 3)^2$. By Wada's theorem, power series: $\sum_1 : k = 1 \rightarrow \infty, d_p(k) \equiv (k+p-1)!/p!(k-1)!$. $\sum_1 \{d_0(k)(-1)^{k-1}(5^{k-1}/2^k) + d_1(k)(2^{k-1}/3^{1+k})\}z^{k-1}$ ($|z| < 2/5$), $\sum_1 \{d_1(k)(2^{k-1}/3^{1+k})\}z^{k-1} + \sum_1 d_0(k)(-1)^{k-1}(2^{k-1}/5^k)(1/z^k)$ ($2/5 < |z| < 3/2$), $\sum_1 \{d_0(k)(-1)^{k-1}(2^{k-1}/5^k) + d_1(k)(3^{k-1}/2^{1+k})(1/z)\}(1/z^k)$ ($3/2 < |z|$). By Cauchy's integral formula, $\sum a_n(z-c)^n : n = (-\infty) \rightarrow \infty, a_n = (1/2\pi i) \oint \{f(\zeta)/(\zeta-c)^{n+1}\}d\zeta = (1/n!)f^{(n)}(c)$. If $c = 0$, then $a_n = (1/2\pi i) \oint \{f(\zeta)/\zeta^{n+1}\}d\zeta = (1/n!)f^{(n)}(0)$ is equal to the result from Wada's. It is not easy to find the general term from the recurring series.

- 23 増田 茂 (流体数理古典理論研) La valeur particulière and the eigenvalue 15
 Shigeru Masuda La valeur particulière and the eigenvalue
 (Res. Workshop of Classical Fluid Dynamics)

概要 We discuss the eigenvalue problem, especially, the coincidence between *la valeur particulière* and the eigenvalue. The eigenvalue problem is the model of the Schrödinger equations or the quantum equations, namely, the Sturm–Liouville type boundary value problem of heat diffusion is the model of the Schrödinger equations. Sturm and Liouville discuss *la valeur particulière*, without its corresponding eigenspace, and the definition of eigenvalue and eigenspace/eigenfunction are introduced by Hilbert. This handling of the value is traditionally relates to the studies of linear differential equations, such as by Laplace, Fourier, Poisson, Cauchy, et al. Especially, Poisson's preceding studies contribute to the study of this type differential equations by Sturm and Liouville and convergence of series in his heat theory.

- 24 増田 茂 (流体数理古典理論研) Prévo's study preceding Fourier of heat communication in the history of physico-mathematics 15
 Shigeru Masuda Prévo's study preceding Fourier of heat communication in the history of physico-mathematics
 (Res. Workshop of Classical Fluid Dynamics)

概要 We discuss Prévo's work on heat communication, which precedes Fourier. These situations owe to the arrival of continuum, on which we summarize the topics as the background from the viewpoint of mathematical history in 18–19th centuries as follows: 1) On the attraction and repulsion of molecule, Navier depends on Fourier's principle of heat molecule. The then physico-mathematicians had little evaluated Navier until the top of 20th century. 2) Poisson points strongly out Fourier's invalidity on the handling of De Gua's theory into the transcendental equations. 3) For formulation of heat motion in the fluid, Fourier had submitted this paper, however, until his death, he has not published it, in which he seems to aim the unity of hydro- and thermodynamics. 4) The hydrodynamicists, like Poisson and Cauchy propose the equations in unity of elasticity and fluid. Finally, we evaluate Prévo's work on these background.

代 数 学

9月25日(木) 第II会場

9:30~12:00

- 1 築場 広子 (広尾学園) オイラー陪関数のある方程式について 10
 穴倉 鷹宏 (広尾学園)
 Hiroko Yanaba (Hiroo Gakuen) On some results of the function associated with the Euler function $\varphi(a)$
 Takahiro Shishikura (Hiroo Gakuen)

概要 Given a positive integer a , we have the Euler function $\varphi(a)$. By s we denote the associated function of $\varphi(a)$, which is indicated by $\tilde{\varphi}(a)$.

Given x , we consider the equation $a - 2^{e+1}\tilde{\varphi}(a) = x$, e being a positive integer.

The main result is as follows.

If $x = 0$, then $e = 1$ and $a = 2^\varepsilon$ for some $\varepsilon > 0$. If $x = 1$ and $e \leq 4$, then the equation has no solution.

- 2 飯高 茂 (学習院大*) 約数関数の陪関数について 10
 Shigeru Itaka (Gakushuin Univ.*) On a function associated with $\sigma(a)$

概要 Associated with Euler totient function, we introduced a new function $\tilde{\varphi}(a)$, which is defined to be $\frac{\varphi(a)}{2^s}$, s being the number of distinct prime factors of a .

Here, associated with divisor (sum) function $\sigma(a)$, we introduce a yet another new function $\tilde{\sigma}(a)$, which is defined to be $\frac{\sigma(a)}{2^s}$.

If $2\tilde{\sigma}(a) - a = -1$ then $a = 2^e p$, where $p = 2^{e+1} + 1$ is a prime.

Here $e + 1$ turns out to be 2^m .

- 3 野村 泰敏 * 第2種スターリング数の素数性と可除性 10
 Yasutoshi Nomura Primality and divisibility of Stirling numbers of the 2nd kind

概要 We discuss in this talk when Stirling numbers $S(n, k)$ are prime and when they are divisible by prime powers.

- 4 木内 功 (山口大理)* On a sum involving the Möbius function 10
 南出 真 (山口大理)
 谷川 好男 (名大多元数理)
 Isao Kiuchi (Yamaguchi Univ.) On a sum involving the Möbius function
 Makoto Minamide (Yamaguchi Univ.)
 Yoshio Tanigawa (Nagoya Univ.)

概要 Let $c_q(n)$ be the Ramanujan sum, i.e., $c_q(n) = \sum_{d|(q, n)} d\mu(q/d)$, where μ is the Möbius function.

Chan and Kumchev gave formulas for $\sum_{n \leq y} \left(\sum_{q \leq x} c_q(n) \right)^k$ ($k = 1, 2$). In this our talk, we shall consider formulas for $\sum_{n \leq y} \left(\sum_{n \leq x} \hat{c}_q(n) \right)^k$ ($k = 1, 2$).

- 5 宮之原 永士 (早大理工) 自然数の k 進展開から生ずる超越数について 10
 Eiji Miyanohara (Waseda Univ.) Transcendence of digital expansions generated by a cyclic permutation and k -adic expansion

概要 We prove the transcendence of certain real numbers defined by using a cyclic permutation and k -adic expansion of natural numbers.

- 6 黒沢 健 (東京理大理) ある種の三角関数の q -成分の代数的独立性 10
 塩川 宇賢 (慶大*)
 Takeshi Kurosawa (Tokyo Univ. of Sci.) Algebraic independence of components of certain trigonometric functions
 Iekata Shiokawa (Keio Univ.*)

概要 Let $q \geq 2$ be an integer. We separate a given power series into q subseries, called q -components, according to the residue classes mod q of their powers. We study algebraic independence for values at an algebraic point of q -components for a certain trigonometric functions.

- 7 立谷 洋平 (弘前大理工) 一般約数関数の母関数値の数論的性質について 10
 Yohei Tachiya (Hirosaki Univ.) Arithmetical properties of the values of the generalized divisor function series

概要 Let q ($|q| \geq 2$) and $\ell \geq 2$ be rational integers. Then the ℓ numbers $1, \sum_{n=1}^{\infty} d_k(n)q^{-n}$ ($k = 2, 3, \dots, \ell$) are linearly independent over \mathbb{Q} , where $d_k(n)$ are the generalized divisor functions. This generalizes a result of Erdos who treated the case $\ell = 2$.

- 8 佐藤 一樹 (東北大理) 対角的三次曲面の有理点 10
 Kazuki Sato (Tohoku Univ.) Rational points on diagonal cubic surfaces

概要 We give a numerical sufficient condition for the existence of rational points on diagonal cubic surfaces over the rationals under the assumption that the Tate-Shafarevich group of any elliptic curve over the rationals is finite.

- 9 伊東 杏希子 (神奈川大工)* ある二次体の類数の 3-divisibility について (II) 10
 Akiko Ito (Kanagawa Univ.) On the 3-divisibility of the class numbers of certain quadratic fields (II)

概要 Let m_1 and m_2 be distinct square-free integers with $12 \nmid m_1 m_2$ and let S_+, S_-, S_0 be mutually disjoint finite sets of prime numbers not containing 2, 3, and the prime factors of $m_1 m_2$. We show that there exist infinitely many square-free integers d with $\gcd(m_1 m_2, d) = 1$ such that the class numbers of quadratic fields $\mathbb{Q}(\sqrt{m_1 d})$ and $\mathbb{Q}(\sqrt{m_2 d})$ are both divisible by 3, every prime number $p \in S_+$ splits in $\mathbb{Q}(\sqrt{d})$, every prime number $p \in S_-$ is inert in $\mathbb{Q}(\sqrt{d})$, and every prime number $p \in S_0$ is ramified in $\mathbb{Q}(\sqrt{d})$.

- 10 池田 創一 (名大多元数理)* 多重ゼータ関数の平均値 10
 松岡 謙晶 (名大多元数理)
 Soichi Ikeda (Nagoya Univ.) Mean values of multiple zeta-functions
 Kaneaki Matsuoka (Nagoya Univ.)

概要 We study mean values of the Euler–Zagier multiple zeta-functions.

- 11 小野塚 友一 (名大多元数理) 等号付き多重ゼータ関数の非零領域 10
 Tomokazu Onozuka (Nagoya Univ.) Zero-free regions of multiple zeta star functions

概要 First, as a generalization of the classical notion of Dirichlet series, we define multiple Dirichlet series and we give zero-free regions of multiple Dirichlet series. Next, we give zero-free regions of multiple zeta star functions.

- 12 広中 由美子 (早大教育)* 有限群の部分群に関するゼータ関数 15
 Yumiko Hironaka (Waseda Univ.) Zeta functions of finite groups by counting numbers of subgroups

概要 For a finite group G , we consider the zeta function $\zeta_G(s) = \sum_H |H|^{-s}$, where H runs over the subgroups of G .

We give examples of certain abelian p -group G and non-abelian p -group G' of order p^m , $m \geq 3$ for odd p (resp. 2^m , $m \geq 4$) for which $\zeta_G(s) = \zeta_{G'}(s)$. Hence we see there are many non-abelian groups whose zeta functions have symmetry and Euler product like the case of abelian groups.

On the other hand, we show that $\zeta_G(s)$ determines G within abelian groups.

We will explain some relation to the theory of Eisenstein series of GL_n and local densities of square matrices.

14:15~16:15

- 13 井上 公人 (九大数理) マイクスナー・ポーラチック多項式による完備リーマンゼータ関数の展開とその近似零点 10
 Hiroto Inoue (Kyushu Univ.) Expansion of the completed Riemann zeta function in Meixner–Pollaczek polynomials and its zeros

概要 We will introduce the expansion of the completed Riemann zeta function in the Meixner–Pollaczek polynomials, which satisfy the functional equation. It is showed that the expansion converges uniformly in every compact set in the critical strip. Then we are interested in investigating the zeros of the partial sums because these zeros approximate the nontrivial zeros of the Riemann zeta function. Furthermore, we will have an observation that almost all the approximating zeros seem to lie on the real axis in a numerical calculation. For the last, we will mention the representation theoretical background of the Meixner–Pollaczek polynomials and the future works.

- 14 矢代好克 (名大多元数理)* Cusp form に付随する L -関数の一般階導関数の零点分布と零点密度 ... 10
 Yoshikatsu Yashiro (Nagoya Univ.) Distribution of zeros and zero-density estimate for the derivatives of L -function attached to cusp form

概要 Let $L_f^{(m)}(s)$ the m -th derivative of L -function attached to cusp form. In this talk we shall give the approximate formula of the number of zeros of $L_f^{(m)}(s)$ and estimate the density of zeros of $L_f^{(m)}(s)$ in the right hand side of critical line.

- 15 若狭尊裕 (名大多元数理)* 一般 Riemann 予想の下での Dirichlet L 関数の偏角の明示的上界について 10
 Takahiro Wakasa (Nagoya Univ.) An explicit upper bound of the argument of Dirichlet L -functions on the generalized Riemann hypothesis

概要 We prove an explicit upper bound of the function $S(t, \chi)$, defined by the argument of Dirichlet L -functions. An explicit upper bound of the function $S_1(t)$, defined by the integral of the argument of the Riemann zeta-function, have already been obtained by A. Fujii. Our result is obtained by applying an idea of Fujii's result on $S_1(t)$. The constant part of the explicit upper bound of $S(t, \chi)$ in this paper does not depend on a primitive Dirichlet character $\chi \pmod{q > 1}$.

- 16 源嶋孝太 (阪大理) $SL(2, \mathbf{C})$ 上の新谷関数と Heun の微分方程式 15
 Kohta Gejima (Osaka Univ.) Shintani functions on $SL(2, \mathbf{C})$ and Heun's differential equations

概要 Shintani functions for the semisimple symmetric pair $(SL(2, \mathbf{C}), GL(1, \mathbf{C}))$ are defined for a pair (π, η) of representations. Here π is a non-unitary principal series representation of $SL(2, \mathbf{C})$ induced from its parabolic subgroup, and η a character of $GL(1, \mathbf{C})$. In this talk, we give explicit formulas of Shintani functions of type (π, η) , where π has a non-trivial minimal $SU(2)$ -type $\text{Sym}^n(\mathbf{C}^2)$ ($n = 1, 2$).

- 17 杉山真吾 (阪大理) L 値が非消滅である Hilbert 尖点形式の存在 15
 都築正男 (上智大理工)
 Shingo Sugiyama (Osaka Univ.) Existence of Hilbert cusp forms with non-vanishing L -values
 Masao Tsuzuki (Sophia Univ.)

概要 We give a derivative version of the relative trace formula on $PGL(2)$ studied in our previous work, and obtain a formula of an average of central values (derivatives) of automorphic L -functions for Hilbert cusp forms. As an application, we prove existence of Hilbert cusp forms with non-vanishing central values (derivatives) such that the absolute degrees of their Hecke fields are sufficiently large.

- 18 林 田 秀 一 (上 越 教 育 大) ジーゲル保型形式の一般化マース関係式について 15
 Shuichi Hayashida Generalizations of the Maass relation of Siegel modular forms
 (Joetsu Univ. of Edu.)

概要 This talk is related to lifts to Siegel modular forms of half-integral weight. The Maass relation is a certain relation among Fourier coefficients of a certain kind of Siegel modular forms of degree two. This relation characterizes the image of the Saito–Kurokawa lift. In particular, Siegel–Eisenstein series of degree two satisfy the Maass relation. In the end of 80’s T. Yamazaki obtained generalizations of the Maass relation for Siegel–Eisenstein series of integral weight of arbitrary degrees. In the present talk I would like to explain some refinements of Yamazaki’s results and give also a generalization of the Maass relation for Siegel cusp forms and Siegel modular forms of half-integral weight of arbitrary degrees. As an application of the generalized Maass relation we obtain lifts from pairs of two elliptic modular forms to Siegel modular forms of half-integral weight of even degrees.

- 19 兒 玉 浩 尚 (近畿大総合理工) On the mod p kernel of the theta operator 10
 長 岡 昇 勇 (近畿大理工)
 Hirotaka Kodama (Kinki Univ.) On the mod p kernel of the theta operator
 Shoyu Nagaoka (Kinki Univ.)

概要 In the previous meeting, it was reported that the image by the theta operator of Igusa cusp form χ_{35} vanishes mod 23. However, the reason why such a phenomenon exists was not clarified. Therefore, we consider whether it can be shown that there exist Siegel modular forms with a similar property. In this meeting, we report that we can construct a family of Siegel modular forms satisfying such property in the case of odd degree.

- 20 見 正 秀 彦 (東京電機大情報) 実指標 Dirichlet L 関数間の同時 d -普遍性と類数たちの多次元稠密性 ... 10
 名 越 弘 文 (群馬大理工)
 Hidehiko Mishou (Tokyo Denki Univ.) Joint d -universality for Dirichlet L -functions with real characters and
 Hirofumi Nagoshi (Gunma Univ.) multi-dimensional denseness of quadratic class numbers

概要 We establish the joint universality theorem for a set of Dirichlet L -functions $L(s, \chi_{d_1}), \dots, L(s, \chi_{d_r})$ with real primitive characters in the d -aspect. We also obtain a multi-dimensional denseness result concerning class numbers of quadratic fields.

16:30~17:30 特別講演

見 正 秀 彦 (東京電機大情報) ゼータ関数の普遍性の概要

Hidehiko Mishou (Tokyo Denki Univ.) An overview of the theory of universality for zeta functions

概要 Let X and Y be a topological space and $T_\iota : X \rightarrow Y$ ($\iota \in I$) be continuous mappings. An element $x \in X$ is called universal for $\{T_\iota\}$, if the set $\{T_\iota x \in Y \mid \iota \in I\}$ is dense in Y .

In the 1910s and later, the existence of universal objects have been confirmed for many operators in the space of entire functions. In 1975, S. M. Voronin obtained the first explicit example of universal objects. Roughly speaking he showed that any non-vanishing and holomorphic function on the strip $1/2 < \Re s < 1$ can be uniformly approximated by a suitable vertical translation $\zeta(s + i\tau)$ of the Riemann zeta function $\zeta(s)$.

In the talk, I will summarize the results and the applications in the theory of universality for arithmetic zeta functions. Also, I will talk about my recent works on the joint universality for sets of several zeta functions.

9月26日(金) 第II会場

9:30~11:5021 高橋 眞 映 (東 邦 大 理)* 実数空間 \mathbb{R} 上の順序位相半群構造について 15

小林 ゆう 治 (東 邦 大 理)

中 筋 康 夫 (放 送 大)

塚 田 眞 (東 邦 大 理)

Sin-Ei Takahasi (Toho Univ.) The structure of ordered topological semigroups on the real space \mathbb{R}

Yuji Kobayashi (Toho Univ.)

Yasuo Nakasuji (Open Univ. of Japan)

Makoto Tsukada (Toho Univ.)

概要 This is a research report about the classification problem of the structure of ordered topological semigroups on the space \mathbb{R} of all real numbers, which is an old and new problem. We determine the structure of semigroups on \mathbb{R} if they are Archimedean and satisfy a certain very weak cancellation law. Moreover we show that exactly 3 cancellative topological semigroups exist on \mathbb{R} and also exactly 8 ordered topological bands exist on \mathbb{R} up to isomorphism.

22 上 岡 修 平 (京 大 情 報) アステカダイヤモンドのタイリングと双直交多項式 15

Shuhei Kamioka (Kyoto Univ.) Tilings of the Aztec diamonds and biorthogonal polynomials

概要 On tiling problems of the Aztec diamonds Stanley exhibited a multivariate generalization of the Aztec diamond theorem by Elkies, Kuperberg, Larsen and Propp (1992). (See Ciucu (1998).) In this talk a new proof of Stabley's multivariate generalization of the Aztec diamond theorem is shown in terms of biorthogonal polynomials and determinants.

- 23 山田 裕史 (岡山大 自然) 分割恒等式と対称群の指標表 15
水川 裕司 (防衛大)
Hiro-Fumi Yamada (Okayama Univ.) A partition identity and the character table of the symmetric groups
Hiroshi Mizukawa
 (Nat. Defense Acad. of Japan)

概要 For an integer r greater than 1, an identity is presented which counts the number of parts modulo r of the r -class regular partitions. This identity is applied for computing the determinant of the regular character table of the symmetric group.

- 24 土谷 昭善 (阪大 情報) 負の係数を持つ Ehrhart 多項式 15
東谷 章弘 (京大 理)
日比 孝之 (阪大 情報)
吉田 恒太郎 (阪大 情報)
Akiyoshi Tsuchiya (Osaka Univ.) Ehrhart polynomials with negative coefficients
Akihiro Higashitani (Kyoto Univ.)
Takayuki Hibi (Osaka Univ.)
Koutarou Yoshida (Osaka Univ.)

概要 It is shown that, for each $d \geq 4$, there exists an integral convex polytope \mathcal{P} of dimension d such that each of the coefficients of n, n^2, \dots, n^{d-2} of its Ehrhart polynomial $i(\mathcal{P}, n)$ is negative.

- 25 日比 孝之 (阪大 情報) squarefree イニシャルイデアルを持つ (0,1) 凸多面体のファセット 15
東谷 章弘 (京大 理)
Takayuki Hibi (Osaka Univ.) Facets of (0,1)-polytopes with squarefree initial ideals
Akihiro Higashitani (Kyoto Univ.)

概要 Let $\mathcal{P} \subset \mathbb{R}^d$ be a (0,1)-polytope of dimension d which contains the origin of \mathbb{R}^d . Suppose that the configuration \mathcal{A} arising from \mathcal{P} satisfies $\mathbb{Z}\mathcal{A} = \mathbb{Z}^{d+1}$ and that the toric ideal $I_{\mathcal{A}}$ of \mathcal{A} possesses a squarefree initial ideal with respect to the reverse lexicographic order induced by an ordering of the variables for which the variable corresponding to the origin is the weakest. It is then proved that the equation of each facet of \mathcal{P} is of the form $\sum_{i=1}^d a_i z_i = b$, where each a_i is an integer and where $b \in \{0, 1\}$.

- 26 松田 一徳 (立教大 理)* 逆辞書式グレブナー基底と強 Koszul トーリック環 10
大杉 英史 (関西学院大 理工)
Kazunori Matsuda (Rikkyo Univ.) Reverse lexicographic Gröbner bases and strongly Koszul toric rings
Hidefumi Ohsugi
 (Kwansei Gakuin Univ.)

概要 In this talk, we give a sufficient condition for a toric ring $K[A]$ to be strongly Koszul in terms of the reverse lexicographic Gröbner bases of its toric ideal I_A . This is a partial extension of a result given by Restuccia and Rinaldo.

In addition, we show that any strongly Koszul toric ring generated by squarefree monomials is compressed. Using this fact, we show that our sufficient condition for $K[A]$ to be strongly Koszul is both necessary and sufficient when $K[A]$ is generated by squarefree monomials.

- 27 奥間 智弘 (山形大理)* 2次元特異点における good ideal と p_g -ideal 10
 渡辺 敬一 (日大文理)
 吉田 健一 (日大文理)
 Tomohiro Okuma (Yamagata Univ.) Good ideals and p_g -ideals for two-dimensional normal singularities
 Kei-ichi Watanabe (Nihon Univ.)
 Ken-ichi Yoshida (Nihon Univ.)

概要 We introduced the notion of p_g ideals for two-dimensional normal local domain. In this talk, we talk about several properties of p_g ideals, and give an existence of p_g ideals. As an application, we prove an existence theorem for good ideals for any two-dimensional Gorenstein normal local domain. Moreover, we classify Ulrich ideals for minimally (simple) elliptic singularities.

- 28 早坂 太 (北教大旭川)* 加群の2変数ブックスバウム・リム関数の計算 10
 Futoshi Hayasaka A computation of Buchsbaum–Rim functions of two variables
 (Hokkaido Univ. of Edu.)

概要 In this talk, we will compute Buchsbaum–Rim functions of two variables associated to a parameter matrix of a special form over a one-dimensional Cohen–Macaulay local ring, and will determine when the function coincides with the Buchsbaum–Rim polynomial. As a consequence, we have that there exists the case where the function does not coincide with the polynomial function, which should be contrasted with the ordinary Buchsbaum–Rim function of single variable.

- 29 張間 忠人 (新潟大教育)* Completely \mathfrak{m} -full ideals and componentwise linear ideals 10
 渡辺 純三 (東海大理)
 Tadahito Harima (Niigata Univ.) Completely \mathfrak{m} -full ideals and componentwise linear ideals
 Junzo Watanabe (Tokai Univ.)

概要 We show that the class of completely \mathfrak{m} -full ideals coincides with the class of componentwise linear ideals in a polynomial ring over an infinite field.

13:20~14:20 特別講演

- 尾角 正人 (阪市大理) 四面体方程式と量子群
 Masato Okado (Osaka City Univ.) Tetrahedron equation and quantum groups

概要 Tetrahedron equation is a 3 dimensional analogue of the Yang–Baxter equation in 2 dimensional solvable lattice models. Recently, new connections between its solutions and quantum groups have been found. They are intertwiners of the quantum coordinate ring associated to SL_3 or Sp_4 , Poincaré–Birkhoff–Witt type basis of the nilpotent part of a quantized enveloping algebra associated to finite-dimensional simple Lie algebras, and a new infinite-dimensional representation, which we call q -oscillator representation, of quantum affine algebras. In my talk I will explain what are interesting from the viewpoint of representation theory of quantum groups.

9月27日(土) 第II会場

9:30~11:25

- 30 竹ヶ原 裕元 (室蘭工大工) 交代群におけるインボリューションの個数の2進的性質について 15
 Yugen Takegahara 2-adic properties for the numbers of involutions in the alternating groups
 (Muroran Inst. of Tech.)

概要 Let $\text{Inv}(n)$ be the set consisting of the identity and all involutions in the alternating groups of degree n . D. Kim and J. S. Kim determined the largest power of 2 in $\#\text{Inv}(n)$, where $n = 4k, 4k + 2$, or $4k + 3$. They also conjectured that the largest power of 2 in $\#\text{Inv}(4k + 1)$, $k = 1, 2, \dots$, are concerned with a certain 2-adic integer. We can give an affirmative answer to this conjecture.

- 31 金久保有輝 (上智大理工)^b Cluster variables on double Bruhat cells and monomial realizations of
 中島俊樹 (上智大理工) crystal bases 15
 Yuki Kanakubo (Sophia Univ.) Cluster variables on double Bruhat cells and monomial realizations of
 Toshiki Nakashima (Sophia Univ.) crystal bases

概要 For semi simple simply connected algebraic group G and elements u, v of its Weyl group, it is known that the coordinate ring $\mathbb{C}[G^{u,v}]$ of the double Bruhat cell $G^{u,v}$ has the structure of an upper cluster algebra and the generalized minors $\Delta(k, \mathbf{i})$ are the cluster variables ([A. Berenstein, S. Fomin, A. Zelevinsky]). In this talk, for $G = SL_{r+1}(\mathbb{C})$, we consider $\Delta(k, \mathbf{i})$ as the functions on $\mathbb{C}_{\neq 0}^{l(u)+l(v)}$ by coordinate transformation. Then they become Laurent polynomials with coefficient 1. And we can express each monomial in those polynomials in terms of monomial realization of crystal.

- 32 安部利之 (愛媛大教育) A_1 型アフィン頂点作用素代数の4次巡回置換オービフォルドモデルにお
 山田裕理 (一橋大経済) ける, あるコミュタントの構造 10
 Toshiyuki Abe (Ehime Univ.) A commutant of a 4-cyclic permutation orbifold model of affine vertex
 Hiromichi Yamada (Hitotsubashi Univ.) operator algebra of type A_1

概要 I will talk on a structure of a vertex operator algebra given as a commutant in the cyclic orbifold model of $L_{sl_2}(1, 0)$ of order 4.

- 33 田邊 顕 一 朗 (北 大 理) 頂点代数上の対数項付き加群について 15
 Kenichiro Tanabe (Hokkaido Univ.) On modules with logarithmic terms over vertex algebras

概要 I will introduce a generalization of modules over vertex algebras and give some examples of such modules.

- 34 小西正秀 (名大多元数理) KLR代数の基本化 15
 Masahide Konishi (Nagoya Univ.) Basicalization of KLR algebras

概要 KLR algebras are introduced to categorify the negative part of the associated quantum group. From another point of view, they are a class of infinite dimensional algebra, defined by two data: a quiver and a weight on its vertices. It is not so pathological, therefore we know we can obtain a basic algebra Morita equivalent to a KLR algebra as a quiver with relations in principle. In this talk, I will explain an explicit algorithm for that in some special cases.

- 35 板場綾子(東京理大理) 分離条件を満たす monomial algebra のホッホシルトコホモロジー群の分解について 10
 古谷貴彦(明海大歯) 解について
 眞田克典(東京理大理)
 Ayako Itaba (Tokyo Univ. of Sci.) On the decomposition of the Hochschild cohomology group of a monomial algebra satisfying a separability condition
 Takahiko Furuya (Meikai Univ.)
 Katsunori Sanada (Tokyo Univ. of Sci.)

概要 In this talk, we consider the finite connected quiver Q having two subquivers $Q^{(1)}$ and $Q^{(2)}$ with $Q = Q^{(1)} \cup Q^{(2)} = (Q_0^{(1)} \cup Q_0^{(2)}, Q_1^{(1)} \cup Q_1^{(2)})$. Suppose that $Q^{(i)}$ is not a subquiver of $Q^{(j)}$ where $\{i, j\} = \{1, 2\}$. For a monomial algebra $\Lambda = kQ/I$ obtained by the quiver Q , when the associated sequence of paths given by I satisfies a certain separability condition, we propose the method so that we easily construct a minimal projective resolution of Λ as a right Λ^e -module and calculate the Hochschild cohomology group of Λ .

- 36 板垣智洋(東京理大理) On the cyclic homology of an algebra associated with a cyclic quiver and a monic polynomial 10
 Tomohiro Itagaki (Tokyo Univ. of Sci.) On the cyclic homology of an algebra associated with a cyclic quiver and a monic polynomial

概要 The Hochschild homology and the cyclic homology of an algebra $K[x]/(f(x))$ over a commutative ring K is given by the Buenos Aires cyclic homology group, where $f(x)$ is a monic polynomial. We consider an algebra A associated with a cyclic quiver and a monic polynomial which generalizes $K[x]/(f(x))$. In this talk, we give the module structure of the Hochschild homology and the cyclic homology of A .

11:30~12:00 代数学分科会総会

14:15~15:15 特別講演

- 源 泰幸(阪府大理)^b Derived bi-duality and DG-completion
 Hiroyuki Minamoto (Osaka Pref. Univ.) Derived bi-duality and DG-completion

概要 Duality and bi-duality are basic operations and has played prominent role in algebra and representation theory. Recently the concern with the derived duality and derived bi-duality have been growing. We show that a derived bi-duality DG-module is obtained as a tautological homotopy limit. This result gives unified proofs and generalizations of previous known results about derived bi-duality. We will also discuss derived double centralizer property, which relates to a notion of DG-completion.

15:30~17:30

- 37 宮原大樹(山梨大工) Auslander-regular フィルター環の構成とその応用 10
 Hiroki Miyahara (Univ. of Yamanashi) A construction of Auslander-regular rings and its application

- 38 星野光男 (筑波大数理物質) Group-graded and group-bigraded rings 15
 亀山統胤 (信州大総合工)
 古賀寛尚 (東京電機大情報環境)
 Mitsuo Hoshino (Univ. of Tsukuba) Group-graded and group-bigraded rings
 Noritsugu Kameyama (Shinshu Univ.)
 Hirotaka Koga (Tokyo Denki Univ.)

概要 Let I be a non-trivial finite multiplicative group with the unit element e and $A = \bigoplus_{x \in I} A_x$ an I -graded ring. We construct a Frobenius extension Λ of A and study when the ring extension A of A_e can be a Frobenius extension. Also, formulating the ring structure of Λ , we introduce the notion of I -bigraded rings and show that every I -bigraded ring is isomorphic to the I -bigraded ring Λ constructed above.

- 39 小池寿俊 (沖縄工高専) 有限三角拡大の自己双対性 10
 Kazutoshi Koike Self-duality of finite triangular extensions
 (Okinawa Nat. Coll. of Tech.)

概要 Azumaya conjectured that every exact ring has a self-duality. The conjecture is related to self-duality of finite triangular extensions. We obtain self-duality of finite triangular extensions of skew fields under certain conditions.

- 40 古賀寛尚 (東京電機大情報環境) On standard derived equivalences 15
 Hirotaka Koga (Tokyo Denki Univ.) On standard derived equivalences

概要 In this talk, we provide another sufficient condition to guarantee the existence of standard derived equivalences for derived equivalent rings.

- 41 山中聡 (岡山大自然)^b On Morita equivalence in ring extensions 10
 Satoshi Yamanaka (Okayama Univ.) On Morita equivalence in ring extensions

概要 It seems that Morita invariance judges of the important of classes of ring extensions concerned. The notion of Morita equivalence in ring extensions was introduced by Y. Miyashita, and he showed that the classes of G -Galois extensions and Frobenius extensions are Morita invariant. After that, S. Ikehata showed that the classes of symmetric extensions, QF-extensions, separable extensions, and Hirata separable extensions are Morita invariant. The purpose of this talk is to show that some classes of ring extensions are Morita invariant. Further, we will give an example of the class of ring extensions which is not Morita invariant. It may seem that the classes of almost all ring extensions are Morita invariant. But, there is a class of ring extensions which we do not know weather that is Morita invariant or not.

- 42 神田遼 (名大多元数理)^b Classification of categorical subspaces of locally noetherian schemes .. 15
 Ryo Kanda (Nagoya Univ.) Classification of categorical subspaces of locally noetherian schemes

概要 We classify the prelocalizing subcategories of the category of quasi-coherent sheaves on a locally noetherian scheme. In order to give the classification, we introduce the notion of a local filter of quasi-coherent subsheaves of the structure sheaf. We also classify the localizing subcategories and the closed subcategories in terms of filters.

- 43 土岡俊介(東大数理) 対称群の次数付一般化カルタン不変量について 15

A. Evseev
(Univ. of Birmingham)

Shunsuke Tsuchioka (Univ. of Tokyo) On graded generalized Cartan invariants of the symmetric groups
Anton Evseev (Univ. of Birmingham)

概要 We propose graded analogue of Hill's conjecture and Külshammer–Olsson–Robinson's conjecture concerning generalized Cartan invariants of the symmetric groups and discuss implications between them and their variants. As a support, we see our conjectures are compatible with certain localization and specialization of the base ring $\mathbb{Z}[v, v^{-1}]$.

- 44 柴田大樹(筑波大数理物質) On integrals for algebraic supergroups 10

Taiki Shibata (Univ. of Tsukuba) On integrals for algebraic supergroups

概要 Let k be a base field. An affine group scheme is a representable functor from the category of commutative k -algebras to the category of groups. If “algebras” are replaced with “superalgebras” (= \mathbb{Z}_2 -graded algebras), then we obtain the notion of an affine supergroup scheme (or simply, supergroup). Let A be the Hopf superalgebra corresponding to an algebraic supergroup G . A non-zero k -linear map $\phi : A \rightarrow k$ is called an integral for G if ϕ is A -colinear. It is known that the integral plays an important role in the representation theory of G . In this talk, we will discuss the existence and properties of integrals.

9月28日(日) 第II会場

9:15~12:00

- 45 岩見智宏(九州産大工)* ある三次元対数的対に対する、弱い意味での Clemens–Griffiths 成分の類似とそれに付随する有理性判定条件について 10

Tomohiro Iwami
(Kyushu Sangyo Univ.) An analogue of Clemens–Griffiths components for certain three-dimensional log pairs in the weak sense and the associated rationality criterion

概要 Succeeding to the author's previous work “A variant of Iskovskikh's rationality criterion for conic bundles in the case of polarized (log) pairs”, in which the author slightly gave a rationality criterion for \mathbb{Q} -conic bundles, as resulting datum of running log minimal model program for certain three-dimensional log pairs under several assumptions, the author will try to induce an analogue of Clemens–Griffiths components in the sense of A. Kuznetsov for rational \mathbb{Q} -conic bundles, with regards to Chow motives, contributing to such a rationality, in the induced Chow–Kunnet decomposition for such conic bundles, also by some methods related to phantom categories.

- 46 鈴木 讓(阪大理) Klein's fundamental second kind 2-form for the C_{ab} curves 15

Jo Suzuki (Osaka Univ.) Klein's fundamental second kind 2-form for the C_{ab} curves

概要 In this paper, we derive the exact formula for symplectic basis of cohomology and its associated Klein's fundamental second kind 2-form for so-called C_{ab} curves. Previously, a similar result was obtained for hyper-elliptic curves. Thus far, the symplectic bases were calculated numerically by solving equations because no such a formula was known for more general curves.

- 47 土基善文 (高知大理) 非可換ケーラー多様体 15
 Yoshifumi Tsuchimoto (Kochi Univ.) Non commutative Kähler maifolds

概要 We define non commutative counterpart of projective variety equipped with a Kähler structure.

- 48 伊藤裕貴 (名大多元数理)* Enriques 曲面上の対合の分類について 15
 大橋久範 (東京理大理工)
 Hiroki Ito (Nagoya Univ.) On the classification of involutions on Enriques surfaces
 Hisanori Ohashi (Tokyo Univ. of Sci.)

概要 We classify involutions on Enriques surfaces with the help of lattice theory due to Nikulin. We also give geometric realizations to all types by mainly Horikawa construction. This is a joint work with Hisanori Ohashi.

- 49 山本悠貴 (金沢大自然) Divisorial contractions to cDV points with discrepancy > 1 15
 Yuki Yamamoto (Kanazawa Univ.) Divisorial contractions to cDV points with discrepancy > 1

概要 I show that every divisorial contraction to cDV point with discrepancy > 1 is a weighted blow-up, except for a few cases.

- 50 那須弘和 (東海大理) Obstructions to deforming space curves lying on a smooth quartic surface 15
 Hirokazu Nasu (Tokai Univ.) Obstructions to deforming space curves lying on a smooth quartic surface

概要 We give a sufficient condition for a first order infinitesimal deformation of a curve on a 3-fold to be obstructed. As an application, we give a new example of a generically non-reduced irreducible components of the Hilbert scheme of space curves, whose general member is contained in a smooth quartic (K3) surface. Our proof is based on a calculation of the obstructions (cup products).

- 51 石川大蔵 (早大理工) Weak Fano bundles of rank 2 on cubic threefolds 15
 Daizo Ishikawa (Waseda Univ.) Weak Fano bundles of rank 2 on cubic threefolds

概要 We classified weak Fano bundles of rank 2 on nonsingular cubic hypersurfaces in projective 4-spaces.

- 52 川口良 (奈良県立医大) 3次元トーリック Fano 多様体の特徴付け 15
 Ryo Kawaguchi (Nara Medical Univ.) A characterization of toric Fano threefolds

概要 In a recent research, we obtain a lower bound for the self-intersection number of an ample divisor on a three-dimensional toric threefold. In this talk, we will see that this bound is attained if and only if our divisor is the anti-canonical divisor on a toric Fano threefolds.

- 53 東谷章弘(京大理) 5次元トーリック Fano 多様体の同値類とその高次元化 15
 Akihiro Higashitani (Kyoto Univ.) Equivalent classes for toric Fano 5-folds and higher dimensions

概要 Let \mathcal{F}_n be the set of isomorphism classes of toric Fano n -folds. We say that X and Y in \mathcal{F}_n are F-equivalent if there exists a sequence of equivariant blow-ups or blow-downs from X to Y through toric Fano n -folds. In this talk, we will present the results on F-equivalent classes and some other equivalent classes for \mathcal{F}_5 . We also consider a higher dimensional analogue of the results on equivalent classes for \mathcal{F}_5 .

14:15~15:15 特別講演

岩成勇(東北大理)^b 導来淡中双対性について

Isamu Iwanari (Tohoku Univ.) Tannaka duality for higher categories

概要 I will discuss Tannaka duality theory for stable infinity-categories; a higher categorical analogue of triangulated categories. I'll describe its backgrounds, motivations, applications to mixed motives, and prospects.

幾 何 学

9月25日(木) 第Ⅲ会場

9:40~12:00

- 1 蛭子井博孝 (Oval Research Center) 非デザルグ系の定理 (ADE Theorem) について 15
 Hirotaka Ebisui (Oval Research Center) About theorem in an-Desargues-system (ADE Theorem)

概要 We have two systems for overrapping two triangles that can be called as non Desargues and Desargues systems. Namely, in Desargues system, 3 lines connected 3 corresponding vertexs of triangle, are concurrence, and in non-Desargues system, they are not concurrence. Oppositly, we call this relation as that two triangles are in Desargues or non-Desargues system. Now we can say easily that the Desargues theorem is in Desargues system. And Hexagon theorem which was found and reported in MSJ2011 at Shinshu Univ by H. Ebisui, is defined as a theorem in non(an)-Desargues system. About this fact, we show again Hexagon theorem which is drawn or constructed in new geometry system so called non(an)-Desargues system, and some corresponding theorems.

- 2 河村彰星 (東大情報理工) 境界上の重みの釣合せ 15
 岡本吉央 (電通大情報理工)
 徳山 豪 (東北大情報)
 Akitoshi Kawamura (Univ. of Tokyo) Weight balancing on boundaries and skeletons
 Yoshio Okamoto (Univ. of Electro-Comm.)
 Takeshi Tokuyama (Tohoku Univ.)

概要 Given a polygonal region containing a target point (which we assume is the origin), it is not hard to see that there are two points on the perimeter that are antipodal, i.e., whose midpoint is the origin. We prove three generalizations of this fact. (1) For any polygon (or any bounded closed region with connected boundary) containing the origin, it is possible to place a given set of weights on the boundary so that their barycenter (center of mass) coincides with the origin, provided that the largest weight does not exceed the sum of the other weights. (2) On the boundary of any 3-dimensional bounded polyhedron containing the origin, there exist three points that form an equilateral triangle centered at the origin. (3) On the 1-skeleton of any 3-dimensional bounded convex polyhedron containing the origin, there exist three points whose center of mass coincides with the origin. This talk is partly based on work presented at the 30th Annual Symposium on Computational Geometry (SoCG 2014) jointly with 10 other authors.

- 3 泉 脩藏 (近畿大理工) 単体不等式の十分性 10
 Shuzo Izumi (Kinki Univ.) Sufficiency of simplex inequalities

概要 Let z_0, \dots, z_n be the facet areas an n -simplex. Then we have the simplex inequalities: $z_p < z_0 + \dots + z_{p-1} + z_{p+1} + \dots + z_n$ ($0 \leq p \leq n$), generalizations of triangle inequalities. Conversely, suppose that numbers $z_0, \dots, z_n > 0$ satisfy these inequalities. Does there exist an n -simplex with facet areas z_0, \dots, z_n ? Kakeya solved this problem affirmatively in the case $n = 3$ and conjectured that the assertion is affirmative also for $n \geq 4$. We assert that this is always affirmative.

- 4 緒方志保 (福岡大理) 完備 λ -超曲面のギャップ定理について 10
 Shiho Ogata (Fukuoka Univ.) Gap theorems of complete λ -hypersurfaces

概要 Since n -dimensional λ -hypersurfaces in Euclidean space are critical points of the weighted area functional for the weighted volume-preserving variations, in this talk, we talk about the rigidity properties of complete λ -hypersurfaces. We give a gap theorem of complete λ -hypersurfaces with polynomial area growth. By making use of the generalized maximum principle for λ -hypersurfaces, we prove a rigidity theorem of complete λ -hypersurfaces. This is a joint work with Professor Q.-M. Cheng and Professor G. Wei.

- 5 榊 真 (弘前大理工)* $S^3 \times R$ と $H^3 \times R$ の平均曲率ベクトル長さ一定曲面について 10
 Makoto Sakaki (Hirosaki Univ.) Surfaces with mean curvature vector of constant length in product spaces

概要 Deforming rotation surfaces with constant mean curvature in S^3 and H^3 to $S^3 \times R$ and $H^3 \times R$ respectively, we give four classes of surfaces with mean curvature vector of constant length in $S^3 \times R$ and $H^3 \times R$. Also we obtain minimal 2-tori in $S^3 \times S^1$.

- 6 入江 博 (東京電機大未来)* 非 Hamilton 体積最小な Hamilton 安定 Lagrange トーラスについて ... 15
 小野 肇 (埼玉大理工)
 Hiroshi Iriyeh (Tokyo Denki Univ.) On Hamiltonian stable Lagrangian tori which are not Hamiltonian volume minimizing
 Hajime Ono (Saitama Univ.)

概要 In 1993, Y.-G. Oh proposed a problem whether standard Lagrangian tori in C^n are volume minimizing under Hamiltonian isotopies of C^n . We prove that most of them do not have such property if the dimension n is greater than two.

- 7 江尻典雄 (名城大理工) 3重周期的極小曲面の退化極限 15
 藤森祥一 (岡山大自然)
 庄田敏宏 (佐賀大文化教育)
 Norio Ejiri (Meijo Univ.) Limits of triply periodic minimal surfaces
 Shoichi Fujimori (Okayama Univ.)
 Toshihiro Shoda (Saga Univ.)

概要 We consider limits of triply periodic minimal surfaces in \mathbb{R}^3 . We prove that some important examples of doubly periodic minimal surfaces can be obtained as limits of triply periodic minimal surfaces.

- 8 小池直之(東京理大理) ホロノミー不変な非等方的表面エネルギー汎関数 15
Naoyuki Koike (Tokyo Univ. of Sci.) A holonomy invariant anisotropic surface energy functional

概要 In this paper, we investigate a holonomy invariant anisotropic surface energy (functional) in a complete Riemannian manifold, where “holonomy invariant” means that the parametric Lagrangian (on the tangent bundle of the Riemannian manifold) used to define the anisotropic surface energy is constant along each holonomy subbundle of the tangent bundle. First we obtain the first variational formula for the functional. Next we shall introduce the notions of an anisotropic convex hypersurface, an anisotropic equifocal hypersurface and an anisotropic isoparametric hypersurface for the functional, and give examples of these hypersurfaces in the case where the ambient space is a symmetric space. Also, we prove that the anisotropic equifocality is equivalent to the anisotropic isoparametricness in the case where the ambient space is a symmetric space of non-negative curvature.

- 9 小磯憲史(阪大理) 2重調和部分多様体 15
浦川 肇(東北大国際教育)
Norihito Koiso (Osaka Univ.) Bi-harmonic submanifold
Hajime Urakawa (Tohoku Univ.)

概要 A map $f : (M, g) \rightarrow (X, h)$ is called a bi-harmonic map if it is a solution of the variational problem defined by the square integral of the tension field τ . A submanifold Y of (X, h) is called a bi-harmonic submanifold if the inclusion map ι is a bi-harmonic map with respect to the induced metric ι^*h . The equation of bi-harmonic submanifold is over-determined, but has many solutions: all minimal submanifolds are bi-harmonic submanifolds. B. Y. Chen conjectured that there are no bi-harmonic submanifolds in Euclidean spaces except minimal submanifolds. In this talk, we show that the conjecture is true for hypersurfaces under a certain generic condition.

14:15~16:10

- 10 正谷優典(名大多元数理) On unbounded diameter of the space of Lagrangian submanifolds in bi-disks 15
Yusuke Masatani (Nagoya Univ.) On unbounded diameter of the space of Lagrangian submanifolds in bi-disks

概要 By using S. Seyfaddini’s idea, we prove unboundedness of Lagrangian Hofer metric associated with uncountably many Lagrangian submanifolds in standard bi-disks $(D^2 \times D^2, \bar{\omega}_0)$.

- 11 今田充洋(慶大理工) \mathbf{C}^{2n} の複素超曲面における複素概接触計量構造について 10
Mitsuhiro Imada (Keio Univ.) On complex almost contact metric structures on complex hypersurfaces of \mathbf{C}^{2n}

概要 Complex contact manifolds, whose definition is analogous to that of real contact manifolds, are also expected to be important, but few examples of complex contact manifolds are known so far. In this report, we show that the (standard) hyperkähler structures on \mathbf{C}^{2n} induce complex almost contact metric structures on some complex hypersurfaces of \mathbf{C}^{2n} . Also, we discuss whether those structures satisfy the normality defined by Ishihara and Konishi.

- 12 古賀 勇 (九大数理)* 複素グラスマン多様体のある種の部分多様体の分類 10
 Isami Koga (Kyushu Univ.) Classification of some submanifolds of the complex Grassmannian manifold

概要 Let $M \rightarrow Gr_p(\mathbb{C}^n)$ be a holomorphic isometric immersion of a compact Kähler manifold into the complex Grassmannian manifold. We assume that the pull-back bundle of the universal quotient bundle $Q \rightarrow Gr_p(\mathbb{C}^n)$ over the complex Grassmannian manifold is projectively flat. In this talk, we classify such submanifolds with parallel second fundamental form.

- 13 前田 定 廣 (佐賀大理工)* 非平坦複素空間形内の自然還元等質実超曲面 15
 田丸 博 士 (広島大理)
 Sadahiro Maeda (Saga Univ.) Naturally reductive homogeneous real hypersurfaces in a nonflat complex space form
 Hiroshi Tamaru (Hiroshima Univ.)

概要 In this talk, we give some geometric characterizations of naturally reductive homogeneous real hypersurfaces in a nonflat complex space form.

- 14 塚 田 和 美 (お茶の水女大理) Totally complex submanifolds of a complex Grassmann manifold of 2-planes 15
 Kazumi Tsukada (Ochanomizu Univ.) Totally complex submanifolds of a complex Grassmann manifold of 2-planes

概要 A complex Grassmann manifold $G_2(\mathbb{C}^{m+2})$ of all 2-dimensional complex subspaces in \mathbb{C}^{m+2} has two nice geometric structures —the Kähler structure and the quaternionic Kähler structure. We study totally complex submanifolds of $G_2(\mathbb{C}^{m+2})$ with respect to the quaternionic Kähler structure. We show that the projective cotangent bundle $P(T^*\mathbb{C}P^{m+1})$ of a complex projective space $\mathbb{C}P^{m+1}$ is a twistor space of the quaternionic Kähler manifold $G_2(\mathbb{C}^{m+2})$. Applying the twistor theory, we construct maximal totally complex submanifolds of $G_2(\mathbb{C}^{m+2})$ from complex submanifolds of $\mathbb{C}P^{m+1}$. Then we obtain many interesting examples.

- 15 入 江 博 (東京電機大未来) 複素旗多様体内の四元数旗多様体の交叉の構造 10
 酒 井 高 司 (首都大東京理工)
 田 崎 博 之 (筑波大数理物質)
 Hiroshi Iriyeh (Tokyo Denki Univ.) On the structure of the intersection of quaternionic flag manifolds in a complex flag manifold
 Takashi Sakai (Tokyo Metro. Univ.)
 Hiroyuki Tasaki (Univ. of Tsukuba)

概要 We study antipodal structure of the intersections of two real forms in complex flag manifolds. In particular, in the complex flag manifold consisting of sequences of complex subspaces in a complex vector space we describe the intersection of real forms consisting of sequences of quaternionic subspaces.

- 16 田崎博之 (筑波大数理物質) 対蹠的部分集合の系列と評価 10
 Hiroyuki Tasaki (Univ. of Tsukuba) Sequences and estimates of antipodal subsets

概要 Let $P_k(n)$ be the set consisting of all subsets of cardinalities k in $\{1, \dots, n\}$. A subset A of $P_k(n)$ is *antipodal*, if for any α and β in A the cardinality of $\alpha - \beta = \{i \in \alpha \mid i \notin \beta\}$ is even. An antipodal subset of $P_k(n)$ is related to an antipodal set of the oriented real Grassmann manifold $\tilde{G}_k(\mathbf{R}^n)$, which was introduced by Chen–Nagano. We have already obtained the classification of maximal antipodal subsets of $P_k(n)$ for $k \leq 4$. We construct sequences of antipodal subsets of $P_k(n)$. Using this we show estimates of the cardinalities of antipodal subsets of $P_5(n)$.

- 17 清原一吉 (岡山大自然)* 楕円体上の共役跡と D_4^+ ラグランジュ特異点 15
 伊藤仁一 (熊本大教育)
 Kazuyoshi Kiyohara (Okayama Univ.) The conjugate locus on ellipsoid and D_4^+ Lagrangean singularity
 Jin-ichi Itoh (Kumamoto Univ.)

概要 In the 2008 annual meeting of MSJ we talked that, on ellipsoids with distinct axes (or more generally on certain Liouville manifolds), the conjugate locus of a “general point” has three components of singularities and they are cuspidal edges. In this talk we show that the end points of those cuspidal edges are D_4^+ Lagrangean singularities (of Arnold) and no other type of singularities do appear. The D_4^+ singularities appear exactly at the points where the multiplicity of the conjugacy is greater than one.

16:30~17:30 特別講演

- 井川治 (京都工繊大工芸)* 対称三対の基礎と応用
 Osamu Ikawa (Kyoto Inst. Tech.) Foundations and applications of symmetric triads

概要 The notion of a symmetric triad with multiplicities is a generalization of that of an irreducible root system with multiplicities. We shall construct symmetric triads with multiplicities from compact symmetric triads with certain conditions. We apply symmetric triads into two directions:

- (1) The orbits of Hermann actions. A Hermann action is a generalization of the isotropy action of a compact symmetric space. We describe the orbit spaces of Hermann actions using symmetric triads with multiplicities, and we also study the properties of each orbit such as regular, minimal, austere and totally geodesic orbit.
- (2) The intersection of two real forms in an irreducible Hermitian symmetric space of compact type, which is a joint work with M. S. Tanaka and H. Tasaki. They proved that the intersection is an antipodal set if it is discrete. We will describe the necessary and sufficient condition that two real forms intersect discretely using a symmetric triad.

9月26日(金) 第三会場

10:30~10:45 2014年度幾何学賞授賞式

10:50~11:50 2014年度幾何学賞受賞特別講演(トポロジー分科会と合同) —倉西正武氏の受賞を記念して—

後藤 竜 司(阪大 理) 倉西数学 —変形理論および CR 幾何について—

宮嶋 公 夫(鹿児島大*)

Ryushi Goto (Osaka Univ.) Kuranishi's masterpieces and their developments in deformation theory

Kimio Miyajima (Kagoshima Univ.*) and CR-geometry

概要 In this talk, we discuss Kuranishi's works and their developments focussing on deformation theory and CR-geometry.

13:15~14:15 特別講演

横田 巧(京大数理研) 半径の小さい CAT(1)-空間上の凸関数と確率測度の重心

Takumi Yokota (Kyoto Univ.) Convex functions and barycenter on CAT(1)-spaces of small radii

概要 CAT(1)-spaces are metric spaces of curvature at most 1 in the sense of Alexandrov defined in terms of geodesic triangles. CAT(0)-spaces are a generalization of simply-connected complete Riemannian manifolds of non-positive curvature and convexity of their distance functions yields many interesting results.

In this talk, we discuss convexity of certain functions on small metric balls in CAT(1)-spaces. This function is discovered by W. Kendall and is similar to the one used by W. Jäger and H. Kaul in the proof of uniqueness of harmonic maps between Riemannian manifolds.

As an application, we prove that any Borel probability measure on a complete CAT(1)-space of small radius admits a unique Karcher mean which is also a unique barycenter. We also see that our Karcher mean on CAT(1)-spaces shares various properties, such as Jensen's inequality, with barycenter on CAT(0)-spaces. This is a generalization of preceding results on CAT(0)-spaces and CAT(1)-spaces of small diameters by K.-T. Sturm, K. Kuwae, S.-i. Ohta, etc.

Finally, we use our barycenter to formulate an analogue of Banach–Saks property for general metric spaces. We prove that complete CAT(1)-spaces with small radii enjoy this property. This can be regarded as an extension of Kakutani's theorem on uniformly convex Banach spaces.

9月27日(土) 第三会場

9:40~12:00

18 奥村 和 浩(旭川工高専) 非平坦複素空間形内の ϕ -不変な Ricci tensor を持つ実超曲面について

Kazuhiro Okumura

(Asahikawa Nat. Coll. of Tech.)

Real hypersurfaces admitting ϕ -invariant Ricci tensors in a nonflat complex space form

10

概要 In this talk, we consider real hypersurfaces with ϕ -invariant Ricci tensors in a non-flat complex space form $\widetilde{M}_n(c)$. In particular, we classify Hopf hypersurfaces having weakly ϕ -invariant Ricci tensor in $\widetilde{M}_n(c)$.

- 19 佐々木東容 (早大理工) 自由群上のサブセットカレント空間上の交叉汎関数 15
 Dounnu Sasaki (Waseda Univ.) An intersection functional on the space of subset currents on a free group

概要 Kapovich and Nagnibeda introduced the space $\mathcal{SCurr}(F_N)$ of subset currents on a free group F_N of rank $N \geq 2$, which can be thought of as a measure-theoretic completion of the set of all conjugacy classes of finitely generated subgroups of F_N . We define a product $\mathcal{N}(H, K)$ of two finitely generated subgroups H and K of the free group F_N by the sum of the reduced rank $\overline{\text{rk}}(H \cap gKg^{-1})$ over all double cosets HgK ($g \in F_N$), and extend the product \mathcal{N} to a continuous symmetric $\mathbb{R}_{\geq 0}$ -bilinear functional on $\mathcal{SCurr}(F_N) \times \mathcal{SCurr}(F_N)$.

- 20 O. Goertsches (Univ. Hamburg) Riemann 葉層の Chern–Simons 型不変量の局所化 15
 野沢啓 (立命館大理工)
 D. Töben
 (Fed. Univ. of São Carlos)
 Oliver Goertsches (Univ. Hamburg) On localization of Chern–Simons type invariants of Riemannian folia-
 Hiraku Nozawa (Ritsumeikan Univ.) tions
 Dirk Töben (Fed. Univ. of São Carlos)

概要 We will present an Atiyah–Bott–Berline–Vergne type localization formula in the context of equivariant basic cohomology for certain Riemannian foliations on compact manifolds. It localizes some Chern–Simons type invariants, for example the volume of Sasakian manifolds or secondary characteristic classes of Riemannian foliations, to the union of closed leaves. We give various examples to illustrate our method.

- 21 金澤知世 (東京理大理) An S^1 -reduction of non-formal star product 10
 吉岡朗 (東京理大理)
 Tomoyo Kanazawa An S^1 -reduction of non-formal star product
 (Tokyo Univ. of Sci.)
 Akira Yoshioka (Tokyo Univ. of Sci.)

概要 We studied and talked about a star product and besides its characteristic equation to obtain the energy-eigenspace of the MIC-Kepler problem. At present, we are going to find a shape in the star product which should produce its characteristic equation for the Hamiltonian of the MIC-Kepler problem, because we reached our solutions by means of reduction of Hamiltonian systems. Since we used the Moyal product on the cotangent bundle of the principal $U(1)$ -bundle whose base space is the three-dimensional Euclidean space except the origin, we tried to research into the reduction of star products; we would like to make a report about a reduced algebra which may be endowed with a star product derived from the Moyal product.

- 22 多羅間大輔 (京大 理) Stability analysis for the free rigid body dynamics on $U(n)$ 15
 Daisuke Tarama (Kyoto Univ.) Stability analysis for the free rigid body dynamics on $U(n)$

概要 The free rigid body dynamics, which is the geodesic flow on $SO(3)$ with respect to a left-invariant metric, is a typical solvable example in analytical mechanics. The stability of its equilibria is well known. In this talk, a natural extension of the free rigid body dynamics to the unitary group $U(n)$ is considered. The dynamics is described by Euler equation on the Lie algebra $\mathfrak{u}(n)$, which has a bi-Hamiltonian structure, and it can be restricted to the adjoint orbit. The complete integrability and the stability of the equilibria on the generic orbits are considered by using the results of Bolsinov and Oshemkov. In particular, it is shown that all the equilibria on generic orbits are Lyapunov stable.

- 23 川村昌也 (首都大東京理工) 超対称性と Gauduchon 予想に関わる Monge–Ampère 方程式の a priori 評価 15
 Masaya Kawamura A priori estimates for the Monge–Ampère equation related to supersymmetry and Gauduchon conjecture
 (Tokyo Metro. Univ.)

概要 Supersymmetry is the symmetry between Bosons and Fermions in superstring theory. Yau and others showed that supersymmetry requires that there exists a balanced metric (i.e., Hermitian metrics satisfying $d(\omega^{n-1}) = 0$) whose Chern–Ricci curvature (Bismut curvature) vanishes on compact complex manifolds. Gauduchon conjecture is Calabi conjecture for Gauduchon metrics (i.e., Hermitian metrics satisfying $\partial\bar{\partial}(\omega^{n-1}) = 0$) on compact complex manifolds. We study on a priori estimates for the Monge–Ampère equation closely related to supersymmetry and Gauduchon conjecture under an estimate of Laplacian of a smooth solution. These estimates are important for applying continuity method to obtain a solution.

- 24 本間泰史 (早大 理工) Twisted Dirac operators and generalized gradients 15
 Yasushi Homma (Waseda Univ.) Twisted Dirac operators and generalized gradients

概要 Generalized gradients are the first order differential operators naturally defined on Riemannian or spin manifolds. It is known that there exist commutation relations for gradients called Weitzenböck formulas. In this talk, we show the other commutation relations by using twisted Dirac operators and PRV theorem in representation theory. We also give some applications.

- 25 十鳥健太 (東北大 理)* Calabi’s gradient metric on the space of Kähler metrics 10
 Kenta Tottori (Tohoku Univ.) Calabi’s gradient metric on the space of Kähler metrics

概要 Calabi’s gradient metric is a Riemannian metric on the space of Kähler metrics \mathcal{H} . Since \mathcal{H} is an infinite dimensional space, the existence of a geodesic is not trivial. In this talk, we will show the Cauchy problem for the geodesic equation has a short time solution. Furthermore, we will show Calabi’s gradient metric defines a distance function on \mathcal{H} .

- 26 納谷 信 (名大多元数理) CR 多様体上の 1 形式に対する Rumin–Bochner 公式 15
 Shin Nayatani (Nagoya Univ.) Rumin–Bochner formular for 1-forms on a CR manifold

概要 Around 1990, Michel Rumin defined a differential complex for a contact manifold, and developed the associated harmonic theory for a strongly pseudoconvex CR manifold with a fixed contact form. Rumin proved a Bochner-type formula for a harmonic one-form on such a manifold. In this talk, I discuss the sharpness of Rumin’s formula with an explicit example and an application of the formula to the eigenvalue estimate of the subelliptic Laplacian. I’ll also discuss a generalization of the formula to the case of two-forms.

14:15~16:00

- 27 高橋 良輔 (名大多元数理) Modified Kähler–Ricci flow on projective bundles 15
 Ryoosuke Takahashi (Nagoya Univ.) Modified Kähler–Ricci flow on projective bundles

概要 In this talk, we propose a method of studying the modified Kähler–Ricci flow on special projective bundles, called admissible bundles, from the view point of symplectic geometry. As a result, we can reduce the modified Kähler–Ricci flow to a simple PDE for a time-dependent function on the interval $[-1, 1]$. Moreover, we show that the solution of this evolution equation converges uniformly to the function corresponding to a GQE metric in exponential order under some assumptions.

- 28 只野 誉 (阪大理) A lower diameter bound for closed domain manifolds of shrinking Ricci-harmonic solitons 15
 Homare Tadano (Osaka Univ.) A lower diameter bound for closed domain manifolds of shrinking Ricci-harmonic solitons

概要 In this talk, we remark that the arguments by Futaki–Sano (Asian J. Math. 17, 17–31, 2013) and Futaki *et al* (Ann. Global Anal. Geom. 44, 105–114, 2013) also work well for closed domain manifolds of shrinking Ricci-harmonic solitons and the arguments also give a lower diameter bound for the domain manifolds.

- 29 櫻井 陽平 (筑波大数理物質) リッチ曲率が下に有界な境界付き多様体の剛性 15
 Yohei Sakurai (Univ. of Tsukuba) Rigidity of manifolds with boundary under a lower Ricci curvature bound

概要 We study Riemannian manifolds with boundary under a lower Ricci curvature bound, and a lower mean curvature bound for the boundary. We prove a volume comparison theorem of Bishop–Gromov type concerning the volumes of the metric neighborhoods of the boundaries. We conclude several rigidity theorems. As one of them, we obtain a volume growth rigidity theorem. We also obtain a splitting theorem of Cheeger–Gromoll type under the assumption of the existence of a single ray.

- 30 竹内 司 (東京理大理) Berger metric の測地流における可積分性について 10
 Tsukasa Takeuchi (Tokyo Univ. of Sci.) Integrability in the geodesic flow for the Berger metric

概要 The $(1,1)$ -tensor field on symplectic manifold that satisfies some integrability conditions is called a recursion operator. It is known the recursion operator is a characterization for integrable systems, and gives constants of motion for integrable systems. Our aim is to construct recursion operators for the geodesic flow for the Berger metric.

- 31 北 別 府 悠 (京 大 理) A finite generation of the fundamental groups on metric measure spaces
 S. Lakzian (HCM) with small linear diameter growth 10
 Yu Kitabeppu (Kyoto Univ.) A finite generation of the fundamental groups on metric measure spaces
 Sajjad Lakzian (HCM) with small linear diameter growth

概要 We prove a finite generation of the fundamental groups on metric measure spaces with small linear diameter growth. This is done by using the Abresch–Gromoll inequality on RCD spaces. Another key idea is the local to global property of nonnegative Ricci curvature bound in the non-smooth setting.

- 32 棕 野 純 一 (名大多元数理)* On the fundamental group of a complete globally hyperbolic Lorentzian
 manifold with a lower bound for the curvature tensor 10
 Junichi Mukuno (Nagoya Univ.) On the fundamental group of a complete globally hyperbolic Lorentzian
 manifold with a lower bound for the curvature tensor

概要 In this talk, we prove the finiteness of the fundamental group of a certain class of complete globally hyperbolic Lorentzian manifolds with the positivity of the curvature tensor.

- 33 宮 地 秀 樹 (阪 大 理)* タイヒミュラー空間上の等長写像のある粗化と剛性 15
 Hideki Miyachi (Osaka Univ.) Rigidity of a coarsification of isometries on Teichmuller space

概要 In this talk, I will give a coarsification (discretization) of isometries on Teichmuller space. I will discuss a rigidity property of the coarsification. If time permits, I will also give a motivation on this work.

16:15~17:15 特別講演

尾 國 新 一 (愛 媛 大 理)* 粗 Baum–Connes 予想と粗代数的トポロジー

Shin-ichi Oguni (Ehime Univ.) The coarse Baum–Connes conjecture and coarse algebraic topology

概要 Coarse geometry studies unbounded proper metric spaces, for example, non-compact complete riemannian manifolds and finitely generated infinite groups with word metrics, from a large-scale geometric viewpoint.

In general, a Dirac-type operator on a non-compact complete riemannian manifold is not a Fredholm operator and thus we cannot take the Fredholm index. John Roe defined an ‘index’ in the K -group of the Roe algebra. The K -group of the Roe algebra depends only on the coarse structure of the manifold and thus can be treated in coarse geometry. Moreover, by regarding the operator as a cycle of the coarse K -homology, John Roe encoded taking the coarse geometric index for the operator as the coarse assembly map. The coarse assembly map from the coarse K -homology to the K -group of the Roe algebra is defined for any proper metric space, and then the coarse Baum–Connes conjecture claims that the coarse assembly map is an isomorphism for any ‘nice’ proper metric space.

In this talk, first I will explain why the coarse Baum–Connes conjecture is interesting. Indeed I will introduce some applications to differential topology and explain that the conjecture is natural in view of coarse algebraic topology. Next I will report known results on the conjecture. Also I will confirm that the conjecture is true for some simple examples by using coarse algebraic topology as one of main tools. Finally I will introduce a result by our joint work with Tomohiro Fukaya (Tohoku University), which claims that the coarse Baum–Connes conjecture is true for the product of finitely many groups which consist of some of $CAT(0)$ groups, hyperbolic groups, polycyclic groups and appropriate relatively hyperbolic groups. Also I will explain ideas for its proof.

函数論

9月25日(木) 第I会場

9:00~12:00

- 1 尾 和 重 義 (大 和 大 教 育)* ムービウス変換に関連したカラテオドリ関数について 15
Shigeyoshi Owa (Yamato Univ.) Notes on Carathéodory functions involving Möbius transformations

概要 Let \mathcal{A} be the class of functions $p(z)$ which are analytic in the open unit disk \mathbb{U} with $p(0) = 1$. If $p(z) \in \mathcal{A}$ satisfies $\operatorname{Re} p(z) > 0$ ($z \in \mathbb{U}$), then $p(z)$ is said to be Carathéodory function in \mathbb{U} . For $p(z) \in \mathcal{A}$, we introduce three subclasses $\mathcal{P}(\alpha)$ for $\alpha < 1$, $\mathcal{P}(\beta)$ for $\beta > 1$, and $\mathcal{P}(\alpha, \beta)$ for $\alpha < 1$ and $\beta > 1$ of \mathcal{A} . Applying Möbius transformations $w(\zeta)$ for $p(z)$, we discuss some properties for Carathéodory functions $p(z)$.

- 2 天 野 政 紀 (東 工 大 理 工) Jenkins–Strebel 測地線間の Teichmüller 距離の極限值について 15
Masanori Amano (Tokyo Tech) On the limit value of the Teichmüller distance between Jenkins–Strebel rays

概要 We consider the asymptotic behavior of two Teichmüller geodesic rays determined by Jenkins–Strebel differentials. We give the limit value of the Teichmüller distance between the rays. In particular, we obtain a condition under which the rays are asymptotic.

- 3 四 之 宮 佳 彦 (早 大 教 育) Veech 曲面の周期点の個数について 15
Yoshihiko Shinomiya (Waseda Univ.) On the numbers of periodic points on Veech surfaces

概要 A Veech surface is a flat surface whose Veech group is a lattice in $\operatorname{PSL}(2, \mathbb{R})$. A Veech surface is arithmetic if the Veech group is commensurable with $\operatorname{PSL}(2, \mathbb{Z})$. Otherwise, the Veech surface is called a non-arithmetic Veech surface. It is known that the number of periodic points on a non-arithmetic Veech surface is finite. We will give upper bounds of the numbers of periodic points on non-arithmetic Veech surfaces.

- 4 柳 下 剛 広 (早 大 理 工) 2乗可積分タイヒミュラー空間上の Weil–Petersson 計量について 15
Masahiro Yanagishita (Waseda Univ.) Weil–Petersson metric on square integrable Teichmüller spaces

概要 The square integrable Teichmüller space is the metric subspace of the Teichmüller space composed of the Teichmüller equivalence classes with square integrable Beltrami coefficient. Since this subspace has a complex structure modeled on a Hilbert space, we can introduce a Hermitian metric, which is called the Weil–Petersson metric. In this talk, we discuss its Kählerity.

- 5 志 賀 啓 成 (東 工 大 理 工) Teichmüller 曲線の剛性と有限性について 15
Hiroshige Shiga (Tokyo Tech) On rigidity and finiteness for Teichmüller curves

概要 Let f be a holomorphic map from a Riemann surface C of finite type to the moduli space of Riemann surfaces. If the map f is a locally isometry, then the pair (C, f) is called a Teichmüller curve. We show a rigidity theorem and a finiteness theorem for Teichmüller curves.

- 6 志賀啓成 (東工大理工) Conformal invariants defined by harmonic functions on Riemann surfaces 15
 Hiroshige Shiga (Tokyo Tech) Conformal invariants defined by harmonic functions on Riemann surfaces

概要 We consider conformal invariants defined by various spaces of harmonic functions on Riemann surfaces. The Harnack distance is a typical one. We give sharp inequalities comparing those invariants with the hyperbolic metric on the Riemann surface and we determine when equalities hold. We also describe the Harnack distance in terms of the Martin compactification and discuss some properties of the distance.

- 7 志賀啓成 (東工大理工) Deformation spaces of Kleinian groups 10
 Hiroshige Shiga (Tokyo Tech) Deformation spaces of Kleinian groups

概要 Let G_0 be a non-elementary Kleinian group. We consider the deformation space of $D(G_0)$, the space of quasiconformal deformations of G_0 , and their complex analytic properties. We show some analytic structures of $D(G_0)$ which are improvements of results by Bers, Kra, Maskit and McMullen. In particular, we clarify that the structures for Kleinian groups with non-simply connected components are different from those for Kleinian groups without non-simply connected components.

- 8 堀田一敬 (東工大理工) L^d -Loewner chains with quasiconformal extensions 15
 Ikkei Hotta (Tokyo Tech) L^d -Loewner chains with quasiconformal extensions

概要 Recently, a new approach in Loewner theory has been proposed which gives a unified treatment of both the radial and chordal version of the Loewner equations. The notion of Loewner chains are generalized in this framework, called a Loewner chain of order d . In this talk we discuss a sufficient condition that a Loewner chain of order d has a quasiconformal extension to the Riemann sphere. It is provided as a generalization of Becker's quasiconformal extension theorem.

- 9 R. M. Porter (CINVESTAV) 線形系による離散擬等角写像 15
 島内宏和 (東北大情報)
 R. Michael Porter (CINVESTAV) Discrete quasiconformal maps via a linear system
 Hirokazu Shimauchi (Tohoku Univ.)

概要 We propose a discretization method for the quasiconformal mappings from the unit disk to itself. The disk is triangulated in a simple way and the quasiconformal mappings are approximated by piecewise linear mappings; the images of the vertices of the triangles are defined by an overdetermined system of linear equations. The linear system is sparse and its solution is obtained by standard least-squares. The least-squares solution is unique and its mapping sequence converges to the true solution in some certain cases.

- 10 宮地秀樹 (阪大理工)* 擬等角写像の無限小空間に関する力学系的考察 15
 Hideki Miyachi (Osaka Univ.) A dynamical approach to the theory of infinitesimal spaces of quasiconformal mappings

概要 In this talk, I will study the infinitesimal spaces of quasiconformal mappings on the plane via continuous flows on the space of quasiconformal mappings.

- 11 柴 雅和 (広島大*) 開リーマン面の閉リーマン面への等角的埋め込み — 周期行列の極值的性
 山口博史 (滋賀大*) 質 — 15
 Masakazu Shiba (Hiroshima Univ.*) Conformal embeddings of an open Riemann surface into closed ones
 Hiroshi Yamaguchi (Shiga Univ.*) —Extremal property of period matrices—

概要 Let R be an open Riemann surface of positive finite genus g and $\{A_j, B_j\}_{j=1}^g$ be a canonical homology basis modulo dividing cycles. For each $t \in (-1, 1]$ and each j there exists a unique holomorphic differential φ_j^t on R such that $\exp(-\pi it/2)\varphi_j^t$ is canonical semiexact in the sense of Kusunoki and the A_k period is equal to δ_{jk} . Let τ_{jk}^t be the B_k period of φ_j^t and fix any real g vector $\mathbf{a} = (a_1, a_2, \dots, a_g)$. Then we show that $\tau_{\mathbf{a}}^t := \sum_{j,k=1}^g a_j a_k \tau_{jk}^t$ describes a circle when t moves in $(-1, 1]$, and that the circle has an interesting extremal property in the set of compact continuations of R .

14:10~16:50

- 12 田所勇樹 (木更津工高専) The period matrix of the hyperelliptic curve $w^2 = z^{2g+1} - 1$ 15
 Yuuki Tadokoro The period matrix of the hyperelliptic curve $w^2 = z^{2g+1} - 1$
 (Kisarazu Nat. Coll. of Tech.)

概要 Let C_g be the hyperelliptic curve defined by the affine equation $w^2 = z^{2g+1} - 1$ for genus $g \geq 2$. We explicitly obtain the period matrix of this curve whose all entries are elements of the $(2g+1)$ -st cyclotomic field.

- 13 李 正勳 (名大多元数理) J -stability of immediately expanding polynomial maps in p -adic dynamics 15
 Junghun Lee (Nagoya Univ.) J -stability of immediately expanding polynomial maps in p -adic dynamics

概要 Given a family $\{f_\lambda\}_{\lambda \in \Lambda}$ of polynomial maps of degree d where Λ is the set of parameters, a polynomial map f_{λ_0} is called J -stable in Λ if there exists a neighborhood of λ_0 in Λ such that for any element λ in the neighborhood, there exists a conjugacy between the dynamics on the Julia sets of f_λ and f_{λ_0} . The aim of this paper is to show that a polynomial map f_{λ_0} over the field \mathbb{C}_p of p -adic complex numbers is J -stable in the set of polynomial maps over \mathbb{C}_p if f_{λ_0} is *immediately expanding*.

- 14 木坂正史 (京大人間環境) Transcendental entire function of slow growth with prescribed polynomial dynamics 15
 Masashi Kisaka (Kyoto Univ.) Transcendental entire function of slow growth with prescribed polynomial dynamics

概要 We construct a transcendental entire functions of arbitrarily slow growth which has a given polynomial dynamics as its subdynamics. We also show several applications of the result, one of which is the following: there exists a transcendental entire function which has a Cremer point but its Julia set is locally connected.

- 15 稲生 啓行 (京大 理) An implosion arising from saddle connection in 2D complex dynamics
 中根 静男 (東京工芸大工) 15
 Hiroyuki Inou (Kyoto Univ.) An implosion arising from saddle connection in 2D complex dynamics
 Shizuo Nakane
 (Tokyo Polytechnic Univ.)

概要 It is known that the fiber Julia set of a skew product is discontinuous if there exist two saddle fixed points one of whose stable manifold intersects the unstable manifold of the other. We will explain this discontinuity by an analogous argument as in the parabolic implosion.

- 16 濱野 佐知子 (福島大人間発達文化) 円環における半完全正則微分のなす空間の再生核 15
 Sachiko Hamano (Fukushima Univ.) Reproducing kernels for the spaces of holomorphic semiexact differentials on annuli

概要 We give a precise representation of a reproducing kernel for the Hilbert space of all holomorphic semiexact differentials on an annulus by using the calculation of the L_1 -constants with two logarithmic poles. We give an example such that the pseudoconvex variation of the annulus with complex parameter t yields the subharmonicity of the L_1 -constant with t .

- 17 足立 真訓 (名大多元数理) 弱擬凸領域の Diederich–Fornaess 指数の大域的評価 15
 J. Brinkschulte (Univ. Leipzig)
 Masanori Adachi (Nagoya Univ.) A global estimate for the Diederich–Fornaess index of weakly pseudoconvex domains
 Judith Brinkschulte (Univ. Leipzig)

概要 A uniform upper bound for the Diederich–Fornaess index is given for weakly pseudoconvex domains whose Levi-form of the boundary vanishes in ℓ -directions everywhere.

- 18 清水 悟 (東北大 理)^b 原点を含むある種の非有界ラインハルト領域に関する正則同値問題 15
 Satoru Shimizu (Tohoku Univ.) Holomorphic equivalence problem for a class of unbounded Reinhardt domains containing the origin

概要 In this talk, I talk about the holomorphic equivalence problem for (essentially) unbounded pseudoconvex Reinhardt domains. When the domains contain no coordinate hyperplanes, an affirmative answer was already given. As an opposite case to such a case, we discuss a class of unbounded Reinhardt domains containing the origin, which are said to be of broadened type. The main purpose of this talk is to give an affirmative answer to the holomorphic equivalence problem for them by making use of the way of a Liouville foliation.

- 19 相川 弘明 (北大 理) Averaging property of capacity 15
 伊藤 翼 (東工大 理工)
 Hiroaki Aikawa (Hokkaido Univ.) Averaging property of capacity
 Tsubasa Itoh (Tokyo Tech)

概要 We study the distribution of a subset of \mathbb{R}^n with respect to the p -capacity. Stegenga investigated the distribution with respect to the logarithmic capacity and he proved that the averaging property holds for logarithmic capacity. We show that the same averaging property holds for the p -capacity.

- 20 大野 貴雄 (大分大教育福祉)* Trudinger's exponential integrability for Riesz potentials of functions in
水田 義弘 (広島工大工) generalized grand Morrey spaces 15
Takao Ohno (Oita Univ.) Trudinger's exponential integrability for Riesz potentials of functions in
Yoshihiro Mizuta generalized grand Morrey spaces
(Hiroshima Inst. of Tech.)

概要 Our aim in this talk is to discuss Trudinger's exponential integrability for Riesz potentials of functions in generalized grand Morrey spaces. Our result will imply the boundedness of the Riesz potential operator from a grand Morrey space to a Morrey space.

- 21 大野 貴雄 (大分大教育福祉)* Trudinger's inequality for Riesz potentials of functions in Musielak-
下村 哲 (広島大教育) Orlicz spaces 10
Takao Ohno (Oita Univ.) Trudinger's inequality for Riesz potentials of functions in Musielak-
Tetsu Shimomura (Hiroshima Univ.) Orlicz spaces

概要 In this talk we are concerned with Trudinger's inequality for Riesz potentials of functions in Musielak-Orlicz spaces.

17:00~18:00 特別講演

- 水田 義弘 (広島工大工) 変動指数をもつ関数空間におけるソボレフ型定理
Yoshihiro Mizuta Function spaces of variable exponent and Sobolev's theorem
(Hiroshima Inst. of Tech.)

概要 First we show the well-known Sobolev type inequality, and then extend it to the variable exponent settings.

9月26日(金) 第I会場

9:00~10:30

- 22 足立 幸信 * \mathbf{C}^2 から \mathbf{C}^2 への非退化超越正則写像のジュリアの方向について 15
Yukinobu Adachi On the Julia directions of the value distribution of nondegenerate transcendental holomorphic maps of \mathbf{C}^2 to \mathbf{C}^2

概要 We prove that for a nondegenerate holomorphic map $F = (f(x, y), g(x, y))$ where f and g are entire functions and f is a transcendental one, there exists a ray $J(\theta) = \{(x, y); x = t \exp(i\theta), y = kt \exp(i\theta) (0 \leq t < \infty)\}$ where k is an arbitrarily fixed complex number except Lebesgue measure zero set and θ is some real number depending on value k , such that $F(x, y)$, in any open cone in \mathbf{C}^2 with vertex $(0, 0)$ containing $J(\theta)$, does not omit any algebraic curve with three irreducible components in a general position.

- 23 足立 幸信 * 高次元のリーマンの除去可能定理について 15
Yukinobu Adachi On a high dimensional Riemann's removability theorem

概要 Let M be a (connected) complex manifold and E be a closed capacity zero set. Let X be a (connected) complex compact Kobayashi hyperbolic space whose universal covering is Stein and let f be a holomorphic map of $M - E$ to X . Then f be extended holomorphically to a map of M to X .

- 24 足立 幸信 * 高次元のリーマンの写像定理について 10
 Yukinobu Adachi On a high dimensional Riemann's mapping theorem

概要 We prove that the domain D in $\Gamma \times \mathbf{C}_z$ where Γ is a polydisk centered at (0) and the fiber of D over every point of Γ is a simply connected domain in \mathbf{C}_z which contains a small disk $\{|z| \leq \varepsilon\}$, where ε is independent of every point of Γ , is biholomorphic to some complete Hartogs domain. And we give applications of the uniformization of some fiber spaces.

- 25 泊 昌孝 (日大文理)* 2次元次数付き特異点, および星型特異点の極大イデアルサイクルと基本
 都丸 正 (群馬大医) サイクルについて 15
 Masataka Tomari (Nihon Univ.) On maximal ideal cycle and fundamental cycle of normal two-dimensional
 Tadashi Tomaru (Gunma Univ.) singularities with star-shaped resolution, and graded singularities

概要 We study the maximal ideal cycle and the fundamental cycle for normal two-dimensional singularities with star-shaped resolution. Our interest is the identification of these. If the cordinated ring of singularity have a homogeneous reduced element in the minimal degree, then the identification can be regarded at the central curve. We can show that this condition is satisfied for Brieskorn complete intersection singularity.

- 26 本田 竜広 (広島工大工) Growth and distortion theorems for pluriharmonic mappings 15
 濱田 英隆 (九州産大工)
 G. Kohr (Babeş-Bolyai Univ.)
 Tatsuhiro Honda Growth and distortion theorems for pluriharmonic mappings
 (Hiroshima Inst. of Tech.)
 Hidetaka Hamada
 (Kyushu Sangyo Univ.)
 Gabriela Kohr (Babeş-Bolyai Univ.)

概要 In this talk, we give growth and distortion theorems for affine and linearly invariant families of pluriharmonic mappings of the unit ball B into \mathbf{C}^n . Also, we obtain two-point distortion theorems.

- 27 本田 竜広 (広島工大工) Strongly starlike mappings in several complex variables 10
 濱田 英隆 (九州産大工)
 G. Kohr (Babeş-Bolyai Univ.)
 Kwang Ho Shon (Pusan Nat. Univ.)
 Tatsuhiro Honda Strongly starlike mappings in several complex variables
 (Hiroshima Inst. of Tech.)
 Hidetaka Hamada
 (Kyushu Sangyo Univ.)
 Gabriela Kohr (Babeş-Bolyai Univ.)
 Kwang Ho Shon (Pusan Nat. Univ.)

概要 Let f be a normalized biholomorphic mapping on the Euclidean unit ball in \mathbf{C}^n and let $\alpha \in (0, 1)$. In this talk, we will show that if f is strongly starlike of order α in the sense of Liczberski and Starkov, then it is also strongly starlike of order α in the sense of Kohr and Liczberski. We also give an example which shows that the converse of the above result does not hold in dimension $n \geq 2$.

13:20~14:20 特別講演

阿部 幸隆 (富山大理工)* Analytic study of singular curves

Yukitaka Abe (Univ. of Toyama) Analytic study of singular curves

概要 We study singular curves from analytic point of view. We reformulate the Serre duality, a generalized Abel's theorem etc. in quite natural way from our view point, and give completely analytic proofs of them. We also reconsider Picard varieties, Albanese varieties and generalized Jacobi varieties of singular curves analytically. We call an Albanese variety considered as a complex Lie group an analytic Albanese variety. We investigate them in detail and construct several examples. We think that an analytic Albanese variety is suitable for a generalized Jacobi variety because it has similar properties to those of the Jacobi variety for a compact Riemann surface.

函数方程式論

9月25日(木) 第V会場

9:00~12:00

- 1 小川原 弘士 (熊本大 自然) 一階線形 q 差分方程式を満たす形式ローラン級数の微分超越性 10
 Hiroshi Ogawara (Kumamoto Univ.) Differential transcendency of a formal Laurent series satisfying a rational linear q -difference equation

概要 We report that a formal Laurent series satisfying a rational linear q -difference equation of first order does not satisfy any nontrivial algebraic differential equation over the rational function field unless it represents a rational function. This also provides an algebraic proof of Ishizaki's theorem on differential transcendency for a meromorphic function satisfying a linear q -difference equation, and then slightly relaxes the assumption of the theorem. We also report differential transcendency of non-rational solutions of a rational q -difference Riccati equation and a rational linear homogeneous q -difference equation of second order that have rational solutions.

- 2 西口 純矢 (京大 理) 遅延フィードバック制御による不安定な定常解の安定化: Lambert W 関数によるアプローチ 10
 Junya Nishiguchi (Kyoto Univ.) Stabilization of unstable steady solutions by delayed feedback control: Approach by Lambert W function

概要 The delayed feedback control is a method for changing the structure and nature of solutions of original differential systems by adding a delay term. In this talk, I consider stabilization of unstable steady solutions of an ordinary differential equation $\dot{x}(t) = f(x(t))$ on the Euclidean space by the delayed feedback control. Here a delayed feedback term is represented by $K(x(t-\tau) - x(t))$ using a constant matrix K and a positive number τ . Adding this term, the original ODE becomes a delay differential equation, and we can expect the change of dynamics of the differential equation. For this purpose, I use the Lambert W function, which is the multi-valued inverse of the complex function ze^z . A main theorem shows that the stabilization is always possible if unstable eigenvalues of the steady solution consist of a unique pair of complex conjugates.

- 3 廣 惠 一 希 (城西大 理) 局所 Fourier 変換とブローアップ 10
 Kazuki Hiroe (Josai Univ.) Local Fourier transform and blowing up

概要 We consider a resolution of ramified irregular singularity of formal meromorphic connections on disk via local Fourier transforms as an analogy of the blowing up of the singularity of plane curve germs. A necessary and sufficient condition for an irreducible connection to have a resolution of singularity will be determined. A relationship between Komatsu-Malgrange irregularity of connections and some curve invariants will also be discussed.

- 4 岩木 耕平 (京大数理研) Stokes segment 上における Painlevé 函数の WKB 解析的変換論について 10
 Kohei Iwaki (Kyoto Univ.) On WKB theoretic transformations for Painlevé transcendents on degenerate Stokes segments

概要 We show that the second Painlevé equation (PII) and the third Painlevé equation (PIII'(D7)) of type D7 give a normal form of Painlevé equations on a degenerate Stokes segments connecting two different simple turning points and on a degenerate Stokes segment of loop-type, respectively. That is, any 2-parameter formal solution of a Painlevé equation is reduced to a 2-parameter formal solution of (PII) or (PIII'(D7)) on these degenerate Stokes segments by the transformation.

- 5 山澤 浩司 (芝浦工大デザイン工) q -アナログにおける Briot–Bouquet 型方程式の正則解と特異解の存在について 10
 Hiroshi Yamazawa Existence of holomorphic and singular solutions of q -analogue of Briot–Bouquet type difference-differential equations
 (Shibaura Inst. of Tech.)

概要 In 1990, Gérard–Tahara introduced the Briot–Bouquet type partial differential equation $t\partial_t u = F(t, x, u, \partial_x u)$, and they determined the structure of holomorphic and singular solutions provided that the characteristic exponent $\rho(x)$ satisfies $\rho(0) \notin \mathbf{N}^*$. In this talk we consider holomorphic and singular solutions of the following type of difference-differential equations $tD_q u = F(t, x, u, \partial_x u)$.

- 6 山澤 浩司 (芝浦工大デザイン工) q -アナログにおける線形差分-微分方程式に対する形式解の総和法について 10
 田原 秀敏 (上智大理工)
 Hiroshi Yamazawa q -Analogue of summability of formal solutions of some linear q -difference-differential equations
 (Shibaura Inst. of Tech.)
 Hidetoshi Tahara (Sophia Univ.)

概要 Let $q > 1$. In this talk we considers a linear q -difference-differential equation: it is a q -difference equation in the time variable t , and a partial differential equation in the space variable z . Under suitable conditions and by using q -Borel and q -Laplace transforms, we consider that if it has a formal power series solution $\hat{X}(t, z)$ one can construct an actual holomorphic solution which admits $\hat{X}(t, z)$ as a q -Gevrey asymptotic expansion of order 1.

- 7 柴田 徹太郎 (広島大工)* Asymptotic behavior of the bifurcation diagrams for semilinear problems with cubic-like nonlinearity 10
 Tetsutaro Shibata (Hiroshima Univ.) Asymptotic behavior of the bifurcation diagrams for semilinear problems with cubic-like nonlinearity

概要 We consider the nonlinear ordinary differential equation with a parameter λ and cubic-like nonlinear term $f(u)$. It is known that under the suitable conditions on $f(u)$, there are three bifurcation curves $\lambda = \lambda_j(\xi)$ ($j = 1, 2, 3$) which are parameterized by the maximum norm ξ of the solution u_λ associated with λ . The purpose of this talk is to study the precise global structures of $\lambda_j(\xi)$ ($j = 1, 2, 3$).

- 8 森 竜 樹 (龍谷大理工) Global bifurcation structure of stationary solutions to a cell polarization model 10
 久 藤 衡 介 (電通大情報理工)
 辻 川 亨 (宮 崎 大 工)
 四 ツ 谷 晶 二 (龍 谷 大 理 工)
Tatsuki Mori (Ryukoku Univ.) Global bifurcation structure of stationary solutions to a cell polarization model
 Kousuke Kuto (Univ. of Electro-Comm.)
 Tohru Tsujikawa (Univ. of Miyazaki)
 Shoji Yotsutani (Ryukoku Univ.)

概要 We are interesting in wave-pinning in a reaction-diffusion model for cell polarization proposed by Y. Mori, A. Jilkine and L. Edelstein-Keshet in SIAM J. Appl. Math (2011). Wave-pinning means a phenomenon that a wave of activation of one of the species is initiated at one end of the domain, moves into the domain, decelerates, and eventually stops inside the domain, forming a stationary front. Several mathematical bifurcation results of stationary solutions are obtained by Kuto and Tsujikawa in DCDS Supplement (2013). We propose a method to represent a bifurcation sheet of a shadow-system. It determines the global bifurcation structure of stationary solutions of the shadow-system completely including even secondary bifurcation branches.

- 9 山 崎 貴 士 (島根大総合理工) 重み付き減衰に基づく漸近安定性に対する Smith 型判定基準 10
 杉 江 実 郎 (島根大総合理工)
Takasi Yamasaki (Shimane Univ.) Smith-type criterion for the asymptotic stability based on the weighted damping
 Jitsuro Sugie (Shimane Univ.)

概要 The equations considered in this talk are $x'' + h(t)x' + \omega^2 x = 0$ and its generalization $x'' + h(t)|x'|^{q-2}x' + \omega^2 x = 0$, $q \geq 2$, where the prime denotes d/dt , the damping coefficient $h(t)$ is continuous and nonnegative for $t \geq 0$, and the restoring coefficient ω is positive. A necessary and sufficient condition for the asymptotic stability of the equilibrium is obtained in consideration of the weighted damping. Our result includes many previous ones. Some examples are presented to illustrate the main result.

- 10 柴 山 允 瑠 (阪大基礎工) 変分法による4体問題の超8の字解の存在証明 10
 Mitsuru Shibayama (Osaka Univ.) Variational proof of the existence of the super-eight solution in the four-body problem

概要 Using the variational method, Chenciner and Montgomery proved the existence of an eight-shaped periodic solution of the planar three-body problem with equal masses. Just after the discovery, Gerver have numerically found a similar periodic solution called “super-eight” in the planar four-body problem with equal mass.

In this talk I will prove the existence of the super-eight orbit by using the variational method. The difficulty of the proof is to eliminate the possibility of collisions. In order to solve it, we apply the scaling technique established by Tanaka and investigate the asymptotic behavior of a binary collision.

- 11 谷川 智幸 (熊本大教育) 3階 Emden–Fowler 型微分方程式の正值解の漸近挙動について 10
Tomoyuki Tanigawa (Kumamoto Univ.) Asymptotic behavior of positive solutions of third order Emden–Fowler differential equations

概要 In this talk we devote to the asymptotic analysis of third order Emden–Fowler differential equation

$$x''' + q(t)|x|^\gamma \operatorname{sgn} x = 0, \quad q(t) > 0, \quad 0 < \gamma < 1$$

in the framework of regular variation. It is shown that in case function $q(t)$ is nearly regularly varying accurate information can be acquired about the existence of possible positive solutions of the equation and their asymptotic behavior at infinity.

- 12 宇佐美 広介 (岐阜大工)* 常微分方程式の双曲型方程式への応用 10
Hiroyuki Usami (Gifu Univ.) Applications of ordinary differential equations to hyperbolic equations

概要 We introduce applications of asymptotic theory of ODEs to that of one-dimensional hyperbolic equations via Fourier analysis. In particular, we give sufficient conditions which ensure the existence of limit functions of solutions to some hyperbolic problems.

- 13 宇佐美 広介 (岐阜大工)^b 逆爆発問題の大域解 10
上村 豊 (東京海洋大海洋)
Hiroyuki Usami (Gifu Univ.) Global solution of an inverse blow-up problem
Yutaka Kamimura
(Tokyo Univ. of Marine Sci. and Tech.)

概要 An inverse blow-up problem is discussed. We establish a global continuation result showing that a nonlinearity realizing a blow-up time for large initial data can be continued in the direction of smaller initial data as long as the blow-up time is Lipschitz continuous.

- 14 竹内 慎吾 完全 p 楕円積分と π_3 の計算公式 10
(芝浦工大システム理工)
Shingo Takeuchi Complete p -elliptic integrals and computation of π_3
(Shibaura Inst. of Tech.)

概要 Complete p -elliptic integrals are introduced and applied to compute π_3 , a generalization of π . The strategy is same as those of Salamin (1976) and Brent (1976), but the Legendre relation and the Landen transformation are required for these new integrals.

- 15 西本勝之 (デカルト出版)* The solutions to the Laplace's homogeneous ordinary differential equations by means of the N-fractional calculus 4
 Katsuyuki Nishimoto The solutions to the Laplace's homogeneous ordinary differential equations by means of the N-fractional calculus
 (Descartes Press Co.)

概要 In this article, the solutions to the homogeneous ordinary differential equations with linear coefficients, $\varphi_2 \cdot (az + b) + \varphi_1 \cdot (gz + h) + \varphi \cdot (pz + q) = 0$, ($\varphi\nu = d^\nu \varphi / dz^\nu$ for $\nu > 0$, $\varphi_0 = \varphi = \varphi(z)$, a, b, g, h, p, q ; constants, $agp \neq 0$), which are called as “the Laplace's ordinary differential equations”, are discussed by means of the N (Nishimoto's)-fractional calculus (NFC-Method) (The calculus in the 21th Century).

14:15~16:15

- 16 塚本一郎 (東洋大理工)* $x'' = t^{\alpha\lambda-2}x^{1+\alpha}$ ($\alpha = \lambda_0, \lambda > 0$) のある正值解の漸近的表示について 10
 Ichiro Tsukamoto (Toyo Univ.) On an asymptotic expression of a positive solution of $x'' = t^{\alpha\lambda-2}x^{1+\alpha}$ ($\alpha = \lambda_0, \lambda > 0$)

概要 We state an asymptotic expression of a positive solution of the differential equation denoted in the title. This expression is valid as t tends to $+0$ and improves the expression stated in the meeting of the Mathematical Society of Japan at Kyoto University in 2013.

- 17 難波時永 (東大数理) On cell problems for Hamilton–Jacobi equations with non-coercive Hamiltonians and its application to homogenization problems 10
 中安淳 (東大数理)
 浜向直 (早大教育)
 Tokinaga Namba (Univ. of Tokyo) On cell problems for Hamilton–Jacobi equations with non-coercive Hamiltonians and its application to homogenization problems
 Atsushi Nakayasu (Univ. of Tokyo)
 Nao Hamamuki (Waseda Univ.)

概要 We study the solvability of a cell problem arising in a homogenization problem for a Hamilton–Jacobi equation whose Hamiltonian is not coercive because of its boundedness. By using a certain function, a necessary and sufficient condition to solve the cell problem is given. As its application, we also give a homogenization result.

- 18 水谷治哉 (阪大理工) Strichartz estimates for non-elliptic Schrödinger equations 10
 N. Tzvetkov
 (Univ. Cergy-Pontoise)
 Haruya Mizutani (Osaka Univ.) Strichartz estimates for non-elliptic Schrödinger equations
 Nikolay Tzvetkov
 (Univ. Cergy-Pontoise)

概要 We consider the Schrödinger equation on compact manifolds equipped with possibly degenerate metrics. We prove Strichartz estimates with a loss of derivatives. The rate of loss of derivatives depends on the degeneracy of metrics. For the non-degenerate case we obtain, as an application of the main result, the same Strichartz estimates as that in the elliptic case. This extends Strichartz estimates for the Riemannian Laplacian. We also investigate the optimality of the result for the case on a product of compact Lie groups.

- 19 鈴木拓也 (東大数理) C^1 領域上の有界関数空間で高階楕円型作用素が生成する半群の解析性
 10
 Takuya Suzuki (Univ. of Tokyo) Analyticity of semigroups generated by higher order elliptic operators
 in spaces of bounded functions on C^1 domains

概要 Our goal is to establish the analyticity of semigroups generated by higher order divergence type elliptic operators in spaces of bounded functions with the Dirichlet condition when the domain is uniformly C^1 domain. We mainly consider resolvent estimates in L^∞ type spaces and also consider the existence and uniqueness of the resolvent equations in order to show the generation of analytic semigroups. Masuda–Stewart’s method is a well known method to show the analyticity of semigroups, but this method may need the assumption that the boundary of a domain is uniformly C^{2m} . We relax the smoothness assumption of boundaries to uniformly C^1 by applying resolvent estimates in L^∞ type spaces obtained by a contradiction argument and a blow up argument.

- 20 原 宇 信 (首都大東京理工) 1 階の項をもつ楕円型方程式におけるポテンシャル評価について 10
 Takanobu Hara (Tokyo Metro. Univ.) Potential estimates for elliptic equations with drift terms

概要 We consider divergence form elliptic equations with a strongly singular drift term $-\nabla \cdot (A\nabla u) + \vec{b} \cdot \nabla u = \mu$ in a domain $\Omega \subset \mathbb{R}^n$ ($n \geq 3$). We give a weak-type $L^1 - L^{n/(n-2), \infty}$ estimate for a solution to the Dirichlet problem with the homogeneous boundary condition. Moreover, we give a two-sided pointwise potential estimate for a weak solution.

- 21 加藤伸幸 (日本工大工)* 放物型方程式系の近似解の一致 Hölder 連続性 10
 Nobuyuki Kato (Nippon Inst. of Tech.) Uniform Hölder continuity of approximate solutions to parabolic systems

概要 By means of the discrete Morse flow method we construct approximate solutions to parabolic systems and generate a Cauchy–Euler polygon. From Campanato-type estimates for the approximate solutions, we derive Hölder continuity of the Cauchy–Euler polygon uniformly with respect to the approximation.

- 22 川上竜樹 (阪府大工)* 坂口 茂 (東北大情報) When does the heat equation have a solution with a sequence of similar level sets? 10
 Tatsuki Kawakami (Osaka Pref. Univ.) When does the heat equation have a solution with a sequence of similar level sets?
 Shigeru Sakaguchi (Tohoku Univ.) level sets?

概要 We consider an overdetermined Cauchy problem for the heat equation. We prove that if the problem has a non-trivial non-negative solution with a certain sequence of similar level sets, then the solution must be radially symmetric.

- 23 小川卓克 (東北大理)* 放物型方程式の初期値問題の最大 L^1 正則性について 10
 清水扇丈 (静岡大理)
 Takayoshi Ogawa (Tohoku Univ.) Maximal L^1 -regularity for a Cauchy problem to parabolic equations
 Senjo Shimizu (Shizuoka Univ.)

概要 We consider maximal L^1 -regularity of the Cauchy problem for parabolic equations in the Besov space $\dot{B}_{p,1}^0(\mathbb{R}^n)$ with $1 \leq p \leq \infty$. The estimate obtained here is not available by abstract theory for the class of UMD since the end-point Besov space is included. We consider the endpoint estimate obtained by Danchin and show that the estimate is indeed, optimal. Besides, we discuss the optimality of maximal regularity in L^1 for the linear parabolic equation for the variable coefficient case.

- 24 神保秀一 (北大理)* Eigenvalues of 2nd order elliptic operators in a domain with a thin tubular hole 10
 Shuichi Jimbo (Hokkaido Univ.) Eigenvalues of 2nd order elliptic operators in a domain with a thin tubular hole

概要 In this talk I consider an eigenvalue problem of a 2nd order elliptic operator in a domain with a thin tubular hole. I present an asymptotic formula for the behavior of the eigenvalue when the hole shrinks to a lower dimensional manifold.

16:30~17:30 特別講演

- 塩路直樹 (横浜国大工) 一般化 Pohozaev 関数と楕円型方程式の正值球対称解の一意性について
 Naoki Sioji (Yokohama Nat. Univ.) A generalized Pohozaev identity and uniqueness of positive radial solutions for an elliptic equation

概要 We introduce a generalized Pohozaev identity related to an elliptic equation. By using it, we study the uniqueness of positive radial solutions of the equation in a ball (or in R^n) under the Dirichlet boundary condition. We also study its nondegeneracy in a space of radial functions and in a space which involves nonradial functions. We apply our results to various examples like the scalar field equation, the Brezis–Nirenberg problem on a spherical cap, etc.

9月26日(金) 第V会場

9:00~12:00

- 25 側島基宏 (Univ. of Salento) Weighted Calderón–Zygmund and Rellich inequalities 10
 G. Metafune (Univ. of Salento)
 C. Spina (Univ. of Salento)
 Motohiro Sobajima (Univ. of Salento) Weighted Calderón–Zygmund and Rellich inequalities
 Giorgio Metafune (Univ. of Salento)
 Chiara Spina (Univ. of Salento)

概要 We find necessary and sufficient conditions for the validity of weighted Rellich and Calderón–Zygmund inequalities with respect to L^p -norm ($1 \leq p \leq \infty$) for functions in the whole space. Moreover, weighted Rellich and Calderón–Zygmund inequalities with respect to the operators $L = \Delta + c|x|^{-2}x \cdot \nabla - b|x|^{-2}$ ($c, b \in \mathbb{R}$) are also considered.

- 26 三村与士文 (東北大理) 多角形領域における Keller–Segel 系の定常解の先験的有界性 …………… 10
 Yoshifumi Mimura (Tohoku Univ.) A priori bounds of stationary solutions of two dimensional Keller–Segel system on polygonal domains

概要 We consider stationary solutions of two dimensional Keller–Segel system on polygonal domains and prove that a priori bounds of the solutions fail for specific parameters determined by the position of singular points and angles of domains. The similar results are already proved by Senba and Suzuki in 2000 when domains are open sets in \mathbb{R}^2 with smooth boundaries. The novelty is that foregoing specific parameters depend on angles of domains.

- 27 猪奥倫左 (愛媛大理) Existence, non-existence, and unconditional uniqueness for a heat equation with exponential nonlinearity in \mathbb{R}^2 …………… 10
 B. Ruf (Univ. di Milano)
 E. Terraneo (Univ. di Milano)
 Norisuke Ioku (Ehime Univ.) Existence, non-existence, and unconditional uniqueness for a heat equation with exponential nonlinearity in \mathbb{R}^2
 Bernhard Ruf (Univ. di Milano)
 Elide Terraneo (Univ. di Milano)

概要 In this talk, a heat equation with exponential nonlinearity in \mathbb{R}^2 is discussed. We prove a non-existence result for some initial data in an Orlicz space $\exp L^2(\mathbb{R}^2)$. In order to consider the existence of local solutions and unconditional uniqueness, we introduce the closure of $C_0^\infty(\mathbb{R}^2)$ in $\exp L^2(\mathbb{R}^2)$.

- 28 長谷川翔一 (東北大理) 双曲空間における Hénon 型方程式に対する Liouville の定理 …………… 10
 Shoichi Hasegawa (Tohoku Univ.) Liouville theorem for Hénon type equation on the hyperbolic space

概要 In this talk, we study the Liouville result of nontrivial stable solutions of the equation
 (H) $-\Delta_{\mathbb{H}} u = (\sinh d_{\mathbb{H}}(0, x))^\alpha |u|^{p-1} u$ in \mathbb{B}^N ,
 where $\alpha > 0$, $p > 1$, $N \geq 3$. Here the operator $\Delta_{\mathbb{H}}$ denotes the Laplace–Beltrami operator on the hyperbolic space \mathbb{B}^N and $d_{\mathbb{H}}(0, x)$ is the hyperbolic distance from the origin to $x \in \mathbb{B}^N$. We show that there exists an exponent $p_c(\alpha, N)$ such that the equation (H) has no nontrivial stable solutions for $1 < p < p_c$. We can expect that p_c will be critical from the asymptotic behavior of the radial solutions of (H) for $p \geq p_c$ large enough.

- 29 石関 彩 (埼玉大理工)* 分解されたメビウス・エネルギーの変分公式とその評価 …………… 10
 長澤 壯之 (埼玉大理工)
 Aya Ishizeki (Saitama Univ.) Variational formulae of decomposed Möbius energy and estimates
 Takeyuki Nagasawa (Saitama Univ.)

概要 The Möbius energy can be decomposed into three parts, each of which is invariant under Möbius transformations. The third part is the absolute constant. Hence when we consider the energy as a variational problem, we may deal with first two parts only. We find some similarities between these parts and relatively easily get variational formulae and applicable estimates.

- 30 久藤 衡介 (電通大情報理工) Limiting structure of shrinking solutions to the stationary SKT model with large cross-diffusion 10
 Kousuke Kuto (Univ. of Electro-Comm.) Limiting structure of shrinking solutions to the stationary SKT model with large cross-diffusion

概要 We are concerned with the limiting behavior of the coexistence steady-states to the Lotka–Volterra competition model as one of the cross-diffusion terms tends to infinity. Under the Neumann boundary condition, Lou–Ni (1999) derived a couple of shadow systems which characterize the limiting behaviors of the coexistence steady-states. One of the shadow system characterizing the segregation of the competing species has been studied by Lou–Ni–Yotsutani (2004) in detail. This talk focuses on the other shadow system characterizing the shrinking of the species not endowed with the cross-diffusion effect. The bifurcation structure of nonconstant solutions to the shadow system will be stated. In particular for the 1D case, we obtain an unbounded connected set of nonconstant solutions.

- 31 宮本 安人 (東大数理) 優臨界準線形楕円型方程式の正值球対称解の交点数と分岐図式について 10
 Yasuhito Miyamoto (Univ. of Tokyo) Intersection properties of radial solutions and global bifurcation diagrams for supercritical quasilinear elliptic equations

概要 We study the positive solution $u(r, \rho)$ of the quasilinear elliptic equation

$$\begin{cases} r^{-\gamma}(r^\alpha|u'|^\beta u')' + |u|^{p-1}u = 0, & 0 < r < \infty, \\ u(0) = \rho > 0, \quad u'(0) = 0. \end{cases}$$

This class of differential operators includes the usual Laplace, m -Laplace, and k -Hessian operators in the space of radial functions. The equation has a singular positive solution $u^*(r)$. A generalized Joseph–Lundgren exponent, which we denote by p_{JL}^* , is obtained. We study the intersection numbers between $u(r, \rho)$ and $u^*(r)$ and between $u(r, \rho_0)$ and $u(r, \rho_1)$, and see that p_{JL}^* plays an important role.

- 32 田中 視英子 (東京理大理)* Generalized eigenvalue problem for (p, q) -Laplacian with indefinite weight 10
 D. Motreanu (Univ. de Perpignan)
 Mieko Tanaka (Tokyo Univ. of Sci.) Generalized eigenvalue problem for (p, q) -Laplacian with indefinite weight
 Dumitru Motreanu (Univ. de Perpignan)

概要 We provide existence and non-existence results of a positive solution for the (p, q) -Laplace equation under the Dirichlet boundary condition, with $1 < q < p$.

- 33 田中 視英子 (東京理大理)* Bifurcation of positive solutions for the one dimensional (p, q) -Laplace equation 10
 梶木 屋龍治 (佐賀大理工)
 田中 敏 (岡山理大理)
 Mieko Tanaka (Tokyo Univ. of Sci.) Bifurcation of positive solutions for the one dimensional (p, q) -Laplace equation
 Ryuji Kajikiya (Saga Univ.)
 Satoshi Tanaka (Okayama Univ. of Sci.)

概要 In this talk, we consider the bifurcation of positive solutions for the one dimensional (p, q) -Laplace equation under the Dirichlet boundary condition. We investigate the shape of the bifurcation diagram near the bifurcation point and near infinity. As a consequence, we prove the existence of multiple positive solutions and show the uniqueness of positive solutions for a bifurcation parameter in a certain range.

- 34 内 免 大 輔 (阪 市 大 理) Dirichlet 積分量を持つ非線形楕円型方程式の多重解の存在について ... 10
 Daisuke Naimen (Osaka City Univ.) On the multiple solutions of a nonlinear elliptic problem with the Dirichlet energy

概要 In this talk, we consider a nonlinear elliptic equation involving the Dirichlet energy. We introduce a new approach which enables us to get multiple (possibly infinitely many) solutions. In addition, we put a positive answer to the solvability of the critical problem in high dimension which has never been solved.

- 35 橋 詰 雅 斗 (阪 市 大 理) A minimization problem with a sign changing condition 10
 Masato Hashizume (Osaka City Univ.) A minimization problem with a sign changing condition

概要 In this talk, we consider a minimization problem with a sign changing condition. This problem is related with “ L^p Lyapunov inequality”. The author proves the existence of the minimizer of the minimization problem.

- 36 藤 田 安 啓 (富 山 大 理)* 局所 Lipschitz 連続函数に対する対数型ソボレフの不等式 10
 Yasuhiro Fujita (Univ. of Toyama) Log-Sobolev inequality for locally Lipschitz continuous functions

概要 We discuss a logarithmic Sobolev inequality for locally Lipschitz continuous functions on \mathbb{R}^n .

- 37 小坂 篤志 (阪市大数学研) \mathbb{S}^N 上の測地球における Laplace–Beltrami 作用素の固有値の摂動問題
 壁谷 喜継 (阪府大工) 10
 川上 竜樹 (阪府大工)
 二宮 広和 (明大先端数理)
Atsushi Kosaka (Osaka City Univ.) Asymptotic behavior of eigenvalues to the Laplace–Beltrami operator
Yoshitsugu Kabeya (Osaka Pref. Univ.) on a spherical cap in \mathbb{S}^N
Tatsuki Kawakami (Osaka Pref. Univ.)
Hirokazu Ninomiya (Meiji Univ.)

概要 In this conference, we speak of the linear eigenvalue problem of the Laplace–Beltrami operator on a large spherical cap. If a spherical cap is sufficiently large, then we can treat the eigenvalue problem as the perturbation of the eigenvalue problem on a unit sphere.

In this study, we obtained asymptotic formulas of eigenvalues. By using the asymptotic formulas, we can exactly see the multiplicity of eigenvalues of the Laplace–Beltrami operator on a large spherical cap. This result is applied for, e.g., the investigation of a local bifurcation structure of semilinear elliptic problems on a spherical cap.

- 38 Soohyun Bae (Hanbat Nat. Univ.) Critical phenomena in the separation property for semilinear elliptic
 内藤 雄基 (愛媛大理) equations 10
 Soohyun Bae (Hanbat Nat. Univ.) Critical phenomena in the separation property for semilinear elliptic
Yūki Naito (Ehime Univ.) equations

概要 We consider the semilinear elliptic equation, and study separation phenomena of positive radial solutions. With respect to intersection and separation, we establish a classification of the solution structures, and investigate the structures of intersection, partial separation and separation.

- 39 高橋 太 (阪市大理) 2次元円環領域上の平均場方程式の2点爆発点の位置について 10
 M. Grossi
 (Univ. di Roma “La Sapienza”)
Futoshi Takahashi (Osaka City Univ.) On the location of two blow up points on an annulus for the mean field
 Massimo Grossi equation
 (Univ. di Roma “La Sapienza”)

概要 We consider the mean field equation on two-dimensional annular domains, and prove that if P_1 and P_2 are two blow up points of a blowing-up solution sequence of the equation, then we must have $P_1 + P_2 = 0$.

13:15~14:15 特別講演

中澤 秀夫 (日本医大)^b 摩擦項を伴う波動方程式の散乱問題とその周辺

Hideo Nakazawa (Nippon Medical School) Scattering problems for wave equations with dissipation and related topics

概要 In this talk, we shall overview several results related to the scattering problem for wave equations with dissipation. Our topics are given as followings. The problem of decay and non-decay of the total energy of the solution, the existence of a scattering state and the principle of limiting absorption, non-uniform decay with linear dissipation of spatial anisotropy, traveling wave solution and exponential decay of the total energy with special Coulomb-type dissipation, spectral analysis of the Schrödinger and wave equation with non-self-adjoint perturbation of rank 1, uniform resolvent estimate for stationary equation in 2D exterior domain and their applications to smoothing estimate and the principle of limiting amplitude. We shall also discuss several unsolved problems.

9月27日(土) 第V会場

9:00~12:00

40 三竹大寿 (広島大 ISSD)* 非線形随伴法を用いた長時間挙動に関する解析: 障害問題 10
Hung V. Tran (Univ. of Chicago)

Hiroyoshi Mitake (Hiroshima Univ.) Analysis on the large-time behavior by the nonlinear adjoint method:
Hung V. Tran (Univ. of Chicago) obstacle problems

概要 Cagnetti, Gomes, Mitake and Tran (2013) introduced a new idea to study the large time behavior for degenerate viscous Hamilton–Jacobi equations. In this talk, we apply the method to study the large-time behavior of the solution to the obstacle problem for degenerate viscous Hamilton–Jacobi equations. We establish the convergence result under rather general assumptions.

41 三竹大寿 (広島大 ISSD)* 無限大ラプラス方程式の弱結合型連立方程式: 存在, 一意性, 一般化された角錐による比較原理 10
Hung V. Tran (Univ. of Chicago)

Hiroyoshi Mitake (Hiroshima Univ.) Weakly coupled systems of the infinity Laplace equations: existence, uniqueness, comparison with generalized cones
Hung V. Tran (Univ. of Chicago)

概要 We derive the weakly coupled systems of the infinity Laplace equations via a tug-of-war game introduced by Peres, Schramm, Sheffield, and Wilson (2009). We establish existence, uniqueness results of the solutions, and introduce a new notion of “generalized cones” for systems. By using “generalized cones” we analyze blow-up limits of solutions.

42 相木雅次 (東京理大理工) Motion of a vortex filament in an external flow 10
井口達雄 (慶大理工)

Masashi Aiki (Tokyo Univ. of Sci.) Motion of a vortex filament in an external flow
Tatsuo Iguchi (Keio Univ.)

概要 We consider a nonlinear model equation describing the motion of a vortex filament immersed in an incompressible and inviscid fluid. In the present problem setting, we also take into account the effect of external flow. We prove the unique solvability, locally in time, of an initial value problem posed on the one-dimensional torus. The problem describes the motion of a closed vortex filament.

- 43 澤田 宙 広 (岐 阜 大 工)* オイラー方程式のシェアフローについて 10
Okihiro Sawada (Gifu Univ.) On the shear flows of the Euler equations

概要 The nonstationary incompressible inviscid flow of ideal fluids in the whole space is considered. This is described as the Cauchy problem of the Euler equations. The mathematical meaning of the shear flow and its application are discussed. It is proved that the shear flow is a unique global-in-time strong solution in certain class due to the argument of renormalization solutions. The proof is based on the boundedness of the solutions to the linear transport equation, component-wisely.

- 44 米 田 剛 (東 工 大 理 工) オイラー方程式の $B_{\infty,1}^1$ クラスにおける局所非適切性について 10
G. Misiolek (Univ. of Notre Dame)
Tsuyoshi Yoneda (Tokyo Tech) Local ill-posedness of the Euler equations in $B_{\infty,1}^1$
Gerard Misiolek (Univ. of Notre Dame)

概要 We show that the incompressible Euler equations are not locally well-posed in the sense of Hadamard in the Besov space $B_{\infty,1}^1$. Our approach relies on the technique of Lagrangian deformations of Bourgain and Li. We show that the assumption that the data-to-solution map is continuous in $B_{\infty,1}^1$ leads to a contradiction with a well-posedness result in $W^{1,p}$ of Kato and Ponce.

- 45 牛 越 惠 理 佳 (玉 川 大 工)* ストークス作用素の固有値に対するアダマール変分公式について 10
神 保 秀 一 (北 大 理)
Erika Ushikoshi (Tamagawa Univ.) Hadamard variational formula for the eigenvalue of the Stokes equations
Shuichi Jimbo (Hokkaido Univ.) with the Dirichlet boundary conditions

概要 We consider the eigenvalue problem of the Stokes equations in a bounded domain. Under the domain perturbation keeping its topological type, we establish the Hadamard variational formula for the eigenvalues of the Stokes equations with the Dirichlet boundary condition. Our result covers the case when the eigenvalue has multiplicity.

- 46 阿 部 健 (名 大 多 元 数 理)* 有界関数空間上のストークス流の評価について 10
Ken Abe (Nagoya Univ.) On estimates for the Stokes flow in a space of bounded functions

概要 We consider the composition operator $S(t)\mathbb{P}\partial$ for the Stokes semigroup $S(t)$ and the Helmholtz projection \mathbb{P} in a space of bounded functions. It is well known that this composition plays a fundamental role for studying the nonlinear Navier–Stokes equations on L^p . The situation is different for the case $p = \infty$ since in this case the Helmholtz projection does not acts as a bounded operator anymore. In this talk, we show that some a priori estimate is valid for the composition operator on a space of bounded functions.

- 47 清水扇丈 (静岡大理) On local well-posedness of incompressible two-phase flows with phase transitions 10
 八木真太郎 (静岡大自然) transitions 10
 Senjo Shimizu (Shizuoka Univ.) On local well-posedness of incompressible two-phase flows with phase transitions
 Shintaro Yagi (Shizuoka Univ.) transitions

概要 The basic model for incompressible two-phase flows with phase transitions where the interface is nearly flat in the case of non-equal densities is considered. The local well-posedness of the model in L_p -setting was proved by Prüss and Shimizu. We prove local well-posedness of the model L_p in time L_q in space setting.

- 48 齋藤平和 (早大理工) Global well-posedness of a free boundary problem for the Navier–Stokes equations in the L_p - L_q framework 10
 柴田良弘 (早大理工) equations in the L_p - L_q framework 10
 Hirokazu Saito (Waseda Univ.) Global well-posedness of a free boundary problem for the Navier–Stokes equations in the L_p - L_q framework
 Yoshihiro Shibata (Waseda Univ.) equations in the L_p - L_q framework

概要 In this talk, we show the global well-posedness of a free boundary problem for the Navier–Stokes equations in the L_p - L_q framework, and large time behavior of solutions. There are many results about free boundary problems for the Navier–Stokes equations in the L_2 - L_2 framework, and also in the L_p - L_p framework more generally. On the other hand, the L_p - L_q framework will be used in our argument. We note that it is essential to control integrability of solutions with respect to time t .

- 49 久保隆徹 (筑波大数理物質) Maximal L_p - L_q regularity of the compressible-incompressible two phase problem, without surface tension and phase transition case 10
 柴田良弘 (早大理工) problem, without surface tension and phase transition case 10
 Takayuki Kubo (Univ. of Tsukuba) Maximal L_p - L_q regularity of the compressible-incompressible two phase problem, without surface tension and phase transition case
 Yoshihiro Shibata (Waseda Univ.) problem, without surface tension and phase transition case

概要 We consider the compressible-incompressible viscous two phase problem without surface tension and phase transition case. In this talk, we would like to report Maximal L_p - L_q regularity of our problem. The essential point of our proof is to show the existence of \mathcal{R} -boundedness of solution operator to the resolvent problem corresponding to our problem.

- 50 久保隆徹 (筑波大数理物質) Local and global well-posedness of the compressible-incompressible two phase problem, without surface tension and phase transition case 10
 柴田良弘 (早大理工) phase problem, without surface tension and phase transition case 10
 Takayuki Kubo (Univ. of Tsukuba) Local and global well-posedness of the compressible-incompressible two phase problem, without surface tension and phase transition case
 Yoshihiro Shibata (Waseda Univ.) phase problem, without surface tension and phase transition case

概要 We consider the compressible-incompressible viscous two phase problem without surface tension and phase transition case. In this talk, we would like to report local and global well-posedness of our problem. Our proof is based on Banach’s fixed point theorem and maximal $L_p - L_q$ estimate of linearized problem corresponding to our problem.

- 51 村田 美帆 (早大理工) 圧縮性粘性流体に対する時間大域解の一意存在性 10
 柴田 良弘 (早大理工)
 Miho Murata (Waseda Univ.) On the global well-posedness for a compressible viscous fluid flow
 Yoshihiro Shibata (Waseda Univ.)

概要 We show a global in time unique existence theorem for a compressible viscous fluid flow with a slip boundary condition in the L_p in time and L_q in space framework with $2 < p < \infty$ and $N < q < \infty$ under the assumption that the domain is bounded and initial data are small. Kobayashi and Zajaczkowski proved the global well-posedness for a compressible viscous fluid flow with slip boundary condition in the L_2 framework. One of the merits of our approach is less regularity on initial data.

- 52 前川 泰則 (東北大理) Large time asymptotics for two-dimensional exterior flows with small circulation at infinity 10
 Yasunori Maekawa (Tohoku Univ.) Large time asymptotics for two-dimensional exterior flows with small circulation at infinity

概要 We consider the incompressible Navier–Stokes equations in a two-dimensional exterior domain Ω , with no-slip boundary conditions. Our initial data are of the form $u_0 = \alpha\Theta_0 + v_0$, where $\alpha \in \mathbb{R}$ and Θ_0 is the Lamb–Oseen vortex with unit circulation at infinity, and v_0 is a solenoidal perturbation belonging to $L^2(\Omega)^2$. When $|\alpha|$ is sufficiently small we show that the solution behaves asymptotically in time like the self-similar Lamb–Oseen vortex with circulation α . This is a global stability result, in the sense that the perturbation v_0 can be arbitrarily large.

- 53 菱田 俊明 (名大多元数理)^b Stability of time-dependent Navier–Stokes flow and algebraic energy decay 10
 M. Schonbek
 (Univ. California, Santa Cruz)
 Toshiaki Hishida (Nagoya Univ.) Stability of time-dependent Navier–Stokes flow and algebraic energy decay
 Maria Schonbek
 (Univ. California, Santa Cruz)

概要 Specific decay rates of energy of disturbance around time-dependent NS flow are discussed.

14:15~16:15

- 54 橋本 伊都子 (富山高専) 空間多次元バーガス方程式に対する球対称解の希薄波の漸近形について 10
 Itsuko Hashimoto
 (Toyama Nat. Coll. of Tech.) Asymptotic stability of rarefaction wave of radially symmetric solutions for Burgers equation in several space dimensions

概要 We present recent results on large-time behavior of the radially symmetric solution for Burgers equation on the exterior of a ball in multi-dimensional space, where boundary data at the far field are prescribed.

In this talk, we prove that for radially symmetric solution to the Burgers equation on multi-dimensional space, the asymptotic states are classified into four cases.

The proof is given by a standard L^2 energy method.

- 55 榎本 翔太 (九大数理) 周期層状領域における圧縮性 Navier–Stokes 方程式の定常解の線形化安
隠居 良行 (九大数理) 定性について 10
Shouta Enomoto (Kyushu Univ.) On linearized stability of stationary solutions to the compressible Navier–
Yoshiyuki Kagei (Kyushu Univ.) Stokes equation in a periodic layer

概要 We consider the stability of stationary solutions to the compressible Navier–Stokes equation in a periodic layer of \mathbb{R}^3 . There exists a spatially periodic stationary solution if the external force is spatially periodic and is sufficiently small in some Sobolev space. We show that the linearized semigroup decays in the L^2 -norm in the order $t^{-\frac{1}{2}}$ if the initial data belongs to L^1 and L^2 . Furthermore, it is shown that the asymptotic leading part of the semigroup is given by a 2-D heat semigroup.

- 56 森 直文 (九大数理) Decay property for the Timoshenko system with thermal effects: Cat-
川島 秀一 (九大数理) taneo versus Fourier’s law 10
Naofumi Mori (Kyushu Univ.) Decay property for the Timoshenko system with thermal effects: Cat-
Shuichi Kawashima (Kyushu Univ.) taneo versus Fourier’s law

概要 We study the Timoshenko system with thermal effects in the one-dimensional whole space. We investigate the dissipative structure of the system and derive the optimal L^2 decay estimate of the solutions in a general situation. Our decay estimate is based on the detailed pointwise estimate of the solutions in the Fourier space and we observe that the decay property is of the regularity-loss type.

- 57 森 直文 (九大数理) Global existence and energy decay of solutions of the nonlinear Timo-
shenko system with memory 10
Naofumi Mori (Kyushu Univ.) Global existence and energy decay of solutions of the nonlinear Timo-
shenko system with memory

概要 We consider the initial value problem for the nonlinear Timoshenko system with a memory term. The main purpose is to investigate the global existence and the optimal decay of the solutions to the nonlinear problem by employing L^2 energy method. Y. Liu and S. Kawashima (2013) showed them by employing a time-weighted L^2 energy method. We show that they can be proved without employing a time-weighted L^2 energy method under less assumptions on the initial data than those which were needed in Y. Liu and S. Kawashima (2013).

- 58 藤江 健太郎 (東京理大理) ロジスティック項と感応性関数をもつ放物・楕円型 Keller–Segel 系の時
M. Winkler (Univ. Paderborn) 間大域解の存在および有界性 10
横田 智巳 (東京理大理)
Kentarou Fujie (Tokyo Univ. of Sci.) Blow-up prevention by logistic sources in a parabolic-elliptic Keller–
Michael Winkler (Univ. Paderborn) Segel system with singular sensitivity
Tomomi Yokota (Tokyo Univ. of Sci.)

概要 This talk is concerned with the parabolic-elliptic Keller–Segel system with singular sensitivity $\frac{\chi_0}{v}$ and logistic source $ru - \mu u^2$ in a smoothly bounded domain $\Omega \subset \mathbb{R}^2$, where $\chi > 0, r \in \mathbb{R}, \mu > 0$. It is shown that in this two-dimensional setting, the absorptive character of the logistic kinetics is sufficient to enforce global existence of classical solutions even for arbitrarily large $\chi_0 > 0$ and any $\mu > 0$ and $r \in \mathbb{R}$. It is moreover shown that if in addition $r > 0$ is sufficiently large then all these solutions are uniformly bounded.

- 59 藤江健太郎 (東京理大理) シグナル依存型感応性関数をもつ放物・放物型 Keller–Segel 系の解の有界性 10
 Kentarou Fujie (Tokyo Univ. of Sci.) Boundedness in a fully parabolic chemotaxis system with singular sensitivity

概要 This talk deals with a fully parabolic chemotaxis system with singular sensitivity $\frac{\chi_0}{\nu}$. The main result solves the open problem that the solution is uniform-in-time bounded for sufficiently small $\chi_0 > 0$, which was conjectured by Winkler (2011).

- 60 石田祥子 (東京理大理) Global existence for a 2D quasilinear chemotaxis-Navier–Stokes system with rotation 10
 Sachiko Ishida (Tokyo Univ. of Sci.) Global existence for a 2D quasilinear chemotaxis-Navier–Stokes system with rotation

概要 We deal with a degenerate chemotaxis-Navier–Stokes system with rotation on a bounded domain in 2 dimension. As to the known result, Tao–Winkler (2012) proved global existence and boundedness in a chemotaxis-Stokes system with arbitrary porous medium diffusion. This talk extends the result for global existence of weak solutions to that for a chemotaxis-Navier–Stokes system with rotation.

- 61 三浦正成 (九大数理) On uniqueness theorem on weak solutions to the parabolic-parabolic
 杉山由恵 (九大数理) Keller–Segel system of degenerate and singular types 10
 Masanari Miura (Kyushu Univ.) On uniqueness theorem on weak solutions to the parabolic-parabolic
 Yoshie Sugiyama (Kyushu Univ.) Keller–Segel system of degenerate and singular types

概要 In this talk, we shall bring a focus onto the parabolic-parabolic and parabolic-elliptic Keller–Segel systems of the singular and degenerate types and show uniqueness of weak solutions in the class of Hoelder continuous functions.

- 62 溝口紀子 (東京学大教育) A new proof to finite-time blowup in the parabolic-parabolic Keller–
 Segel system 10
 Noriko Mizoguchi A new proof to finite-time blowup in the parabolic-parabolic Keller–
 (Tokyo Gakugei Univ.) Segel system

概要 It has remained open for several decades whether finite-time blowup is a generic feature in the parabolic-parabolic Keller–Segel system in two dimensional domains. An affirmative answer was recently given by the speaker and Winkler. However their proof cannot work in the degenerate Keller–Segel system in the higher dimensions. In this talk, we give a new proof to their theorem which can be applied also to such a case.

- 63 溝口紀子 (東京学大教育) Finite-time blowup for the parabolic-parabolic Keller–Segel system with critical diffusion 10
 P. Laurençot
 (Univ. de Toulouse • CNRS)
- Noriko Mizoguchi Finite-time blowup for the parabolic-parabolic Keller–Segel system with critical diffusion
 (Tokyo Gakugei Univ.)
- Philippe Laurençot
 (Univ. de Toulouse/CNRS)

概要 It is known that a similar phenomenon to that in the two-dimensional Keller–Segel system appears in the parabolic-elliptic simplification with degenerate critical diffusion in the higher dimensions. One can take the same approach for the global existence of solution even in the parabolic-parabolic system. However there have been no results on the existence of solutions blowing up in finite time in the parabolic-parabolic system. In this talk, we give a criterion for finite-time blowup to initial data and then show that the criterion is satisfied by a large class of initial data.

16:30~17:30 特別講演

- 津川光太郎 (名大多元数理) 5階の非線形分散型方程式の局所適切性
 Kotaro Tsugawa (Nagoya Univ.) Local well-posedness for fifth-order nonlinear dispersive equations

概要 We consider the Cauchy problem for fifth order dispersive equations on the torus. We assume that the nonlinear part is a polynomial of $u, \partial_x u, \partial_x^2 u$ and $\partial_x^3 u$ and does not include any constants and linear terms. The equations include the fifth order KdV equation, the fifth order modified KdV equation and some equations in physics. In this talk, we discuss on conditions to the nonlinear terms. We first give a necessary and sufficient condition for the local well-posedness when the initial data is sufficiently smooth. Next, we show the local well-posedness with low regularity initial data when the nonlinear term satisfy a cancellation property. In both studies, the main difficulty comes from the derivative loss in the nonlinear terms. Remark that the linear part does not have any smoothing effect since the torus is compact. It seems difficult to show the well-posedness by the standard energy method or the standard fixed point argument when the nonlinear part include $\partial_x^2 u$ or $\partial_x^3 u$. To avoid the difficulty, we combine the energy method and the normal form reduction in the first result and combine the fixed point argument and the normal form reduction in the second result.

9月28日(日) 第V会場

9:00~12:00

- 64 可香谷隆 (北大理) 自由境界の準線形放物型方程式に対する局所存在性 10
 Takashi Kagaya (Hokkaido Univ.) A local existence on a free boundary problem for quasilinear parabolic equation

概要 We prove a local existence on a free boundary problem for quasilinear parabolic equation whose spatial dimension is one. The parabolic equation has Dirichlet and Neumann boundary conditions at each boundary points. This free boundary problem arises in various applications such as the heat equation and the curvature flow. If the equation has no lower order term, the local existence on this free boundary problem is well known. We consider the case for the equation with lower order terms, and examine how the maximal existence time depends on initial data and lower order terms.

- 65 蛭子くるみ (東北大理) 前立腺癌の間欠的内分泌療法を記述するハイブリッドシステムの動的様相
 Kurumi Hiruko (Tohoku Univ.) A dynamical aspect of hybrid system describing intermittent androgen
 suppression therapy of prostate cancer 10

概要 Since a prostate tumor is influenced by androgen, continuous androgen suppression (CAS) therapy is the most famous therapy of prostate cancer in Japan. However the relapse of tumors often occurs in spite of under CAS therapy. Recently, clinical studies suggested that intermittent androgen suppression (IAS) therapy may delay or prevent the relapse. In the IAS therapy, medication is stopped when the size of tumor decreases less than a lower threshold, and resumed when the size exceeds an upper threshold. We deal with a hybrid system describing IAS therapy and prove mathematically that the size of tumor remains in some bounded interval for any time under IAS therapy.

- 66 側島基宏 (Univ. of Salento) Existence of solutions to heat equations with singular lower order terms
 岡沢登 (東京理大理) 10
 横田智巳 (東京理大理)
 Motohiro Sobajima (Univ. of Salento) Existence of solutions to heat equations with singular lower order terms
 Noboru Okazawa (Tokyo Univ. of Sci.)
 Tomomi Yokota (Tokyo Univ. of Sci.)

概要 In this talk we consider the solvability for the Cauchy problem of heat equations in \mathbb{R}^N ($N \geq 2$) with singular lower order terms $\partial u / \partial t = \Delta u - \beta |x|^{-2} x \cdot \nabla u + c |x|^{-2} u + f$, where $\beta \in \mathbb{R}$, $c \leq (N - 2 - \beta)^2 / 4$, u_0 is an initial value and f is an inhomogeneous term. We give a result which generalizes the one of Baras and Goldstein (1984) from $\beta = 0$ to $\beta \neq 0$ and improves the one of Arendt, Goldstein and Goldstein (2006).

- 67 原田潤一 (秋田大教育文化)* ある非線形熱方程式系の爆発点について 8
 Junichi Harada (Akita Univ.) Blow-up set for a parabolic system equation

概要 We discuss blow-up problems for a semilinear parabolic system equation. We characterize the blow-up set by using the zero of solutions.

- 68 浅井智朗 (東大数理)* On self-similar solutions to the surface diffusion flow equations with
 儀我美一 (東大数理) contact angle boundary conditions 10
 Tomoro Asai (Univ. of Tokyo) On self-similar solutions to the surface diffusion flow equations with
 Yoshikazu Giga (Univ. of Tokyo) contact angle boundary conditions

概要 We consider the surface diffusion flow equation on the half line when the curve is represented by the graph of a smooth function. We impose two boundary conditions. The first boundary condition is the contact angle condition and the second boundary condition is the linearized version of the no-flux condition. We construct a bounded self-similar solution to this problem. We further prove the stability of this self-similar solution.

- 69 水野 将司 (日大理工)* Neumann 境界条件付 Allen–Cahn 方程式の特異極限問題 …… 10
利根川 吉廣 (北大理)
Masashi Mizuno (Nihon Univ.) A singular limit problem of the Allen–Cahn equation with Neumann
Yoshihiro Tonegawa (Hokkaido Univ.) boundary conditions

概要 We study a singular limit problem of the Allen–Cahn equation with Neumann boundary conditions and general initial data of uniformly bounded energy. We prove that the time-parametrized family of limit energy measures is Brakke’s mean curvature flow with a generalized right angle condition on the boundary.

- 70 松家 敬介 (東大数理)^b 離散半線形熱方程式の爆発解の存在について …… 10
時弘 哲治 (東大数理)
Keisuke Matsuya (Univ. of Tokyo) Existence of blow-up solutions for a discrete semilinear heat equation
Tetsuji Tokihiro (Univ. of Tokyo)

概要 Existence of blow-up solutions to initial value problems for a discrete analogue of a d -dimensional semilinear heat equation is investigated. We prove that a parameter α in the partial difference equation plays exactly the same role as the parameter of nonlinearity does in the semilinear heat equation. That is, we prove non-existence of non-blow-up solutions for $0 < \alpha \leq 2/d$, and, for $\alpha > 2/d$, existence of non-blow-up solutions for sufficiently small initial data.

- 71 松澤 寛 (沼津工高専)* ある非線形拡散方程式の自由境界問題における spreading speed の評価と
Yihong Du 解の漸近的形状について …… 10
(Univ. of New England)
Maolin Zhou (東大数理)
Hiroshi Matsuzawa Spreading speed and sharp asymptotic profiles of solutions in free bound-
(Numazu Nat. Coll. of Tech.) ary problems for nonlinear diffusion equations
Yihong Du (Univ. of New England)
Maolin Zhou (Univ. of Tokyo)

概要 We study free boundary problems of nonlinear diffusion equations. For monostable, bistable, and combustion types of nonlinearities, Du and Lou (to appear in J. Eur. Math. Soc.) obtained a rather complete description of the long-time dynamical behavior of the problem and revealed sharp transition phenomena between spreading ($u(t, x) \rightarrow 1$ as $t \rightarrow \infty$) and vanishing ($u(t, x) \rightarrow 0$ as $t \rightarrow \infty$). They also determined the asymptotic spreading speed of the fronts when spreading happens. In this talk, we give a much sharper estimate for the spreading speed, and we show that the asymptotic profile of solution depends only on the nonlinearity when spreading happens.

- 72 高橋 仁 (東工大理工) 吸収項付き半線形熱方程式に対する動的特異点を持つ解について …… 10
柳田 英二 (東工大理工)
Jin Takahashi (Tokyo Tech) Solutions with time-dependent singularities for a semilinear heat equa-
Eiji Yanagida (Tokyo Tech) tion with absorption

概要 In this talk, we consider a semilinear heat equation with a nonlinear absorption term. It is shown that if the power of the nonlinearity is in some range, there is no time-dependent singular solution. On the other hand, in other range, two types of time-dependent singular solutions exist, and any singularity is removable if it is weaker than the order of the fundamental solution of the Laplace equation.

- 73 山本 征法 (弘前大理工) 臨界拡散を持つ移流拡散方程式の解の挙動について 10
 杉山 裕介 (東京理大理)
 加藤 圭一 (東京理大理)
 Masakazu Yamamoto (Hirosaki Univ.) Asymptotic expansion of solutions to the drift-diffusion equation with
 Yuusuke Sugiyama critical dissipation
 (Tokyo Univ. of Sci.)
 Keiichi Kato (Tokyo Univ. of Sci.)

概要 The initial value problem for the drift-diffusion equation in the whole space is studied. The dissipation on this equation is given by the half Laplacian. It is proved that the solution to this equation decays as time variable tends to infinity. Moreover the asymptotic expansion of the solution is established.

- 74 中川 和重 (福島大理工)* Global behavior of solutions to degenerate drift diffusion system in be-
 小川 卓克 (東北大理) tween two critical exponents 10
 君島 敦史 (東北大理)
 Kazushige Nakagawa Global behavior of solutions to degenerate drift diffusion system in be-
 (Fukushima Univ.) tween two critical exponents
 Takayoshi Ogawa (Tohoku Univ.)
 Atsushi Kimijima (Tohoku Univ.)

概要 We are concerned with the time global behavior of solutions to the degenerate drift-diffusion system. Our aim is to classify the global existence of the weak solution and the finite time blow up of the solution by the initial condition in between two critical exponents.

- 75 黒木場 正城 (室蘭工大)^b Two dimensional drift-diffusion system in a critical weighted space ... 10
 Masaki Kurokiba Two dimensional drift-diffusion system in a critical weighted space
 (Muroran Inst. of Tech.)

概要 We establish the time local well-posedness of a multi-component parabolic-elliptic drift-diffusion model in two space dimensions in a scaling critical weighted space. In order to show the critical well-posedness, we introduce a refined version of Brezis-Gallouet inequality in term of the Besov space and apply it to the some weighted estimate in two dimensions.

- 76 谷口 雅治 (岡山大自然) Convex compact sets in \mathbb{R}^{N-1} give traveling fronts of cooperation-
 diffusion systems in \mathbb{R}^N 10
 Masaharu Taniguchi (Okayama Univ.) Convex compact sets in \mathbb{R}^{N-1} give traveling fronts of cooperation-
 diffusion systems in \mathbb{R}^N

概要 This paper studies traveling fronts to cooperation-diffusion systems in \mathbb{R}^N for $N \geq 3$. We consider $(N-2)$ -dimensional smooth surfaces as boundaries of strictly convex compact sets in \mathbb{R}^{N-1} , and define an equivalence relation between them. We prove that there exists a traveling front associated with a given surface and that it is asymptotically stable for given initial perturbation. The associated traveling fronts coincide up to phase transition if and only if the given surfaces satisfy the equivalence relation.

- 77 鈴木 貴 (阪大基礎工)^b 2D normalized Ricci flow 軌道の compact 性 —Hamilton の定理の解析的証明— 10
 Takashi Suzuki (Osaka Univ.) Compactness of 2D normalized Ricci flow orbit —an analytic proof of Hamilton’s theorem—

概要 We show compactness of 2D normalized Ricci flow orbit for any compact Riemannian surface without boundary. Hence if the stationary state admits only the trivial solution, then the non-stationary solution converges to a constant. This is an improvement of our previous result in 2010.

14:15~16:15

- 78 岡本 葵 (信州大工) 空間 1 次元 Chern–Simons–Dirac 方程式の初期値問題の非適切性に対する注意 10
 町原 秀二 (埼玉大理工)
 Mamoru Okamoto (Shinshu Univ.) Remarks on ill-posedness of the Cauchy problem for the Chern–Simons–Dirac system in one dimension
 Shuji Machihara (Saitama Univ.)

概要 We consider ill-posedness of the Cauchy problem for the Chern–Simons–Dirac system in one spatial dimension. Bournaveas, Candy, and Machihara proved that local in time well-posedness in $H^s(\mathbb{R}) \times H^r(\mathbb{R})$ with $-1/2 < r \leq s \leq r + 1$. We showed well-posedness holds if s and r satisfy $|s| < 1/2$, $r = -1/2$ or $s = 0$, $-1 \leq r \leq -1/2$. In this talk, we show that these results are optimal in the sense that the Cauchy problem is ill-posed in $H^s(\mathbb{R}) \times H^r(\mathbb{R})$ if s and r do not satisfy these conditions. We deal with both massless and massive cases for the ill-posedness.

- 79 岸本 展 (京大数理研) Normal form reduction for the unconditional uniqueness of periodic nonlinear dispersive equations 10
 Nobu Kishimoto (Kyoto Univ.) Normal form reduction for the unconditional uniqueness of periodic nonlinear dispersive equations

概要 We study the unconditional uniqueness property for periodic nonlinear dispersive equations by successive applications of integration by parts in the time variable, which we call Normal Form Reduction (NFR). We first give an abstract framework of infinite NFR scheme, and then apply it to the nonlinear Schrödinger equations (NLS) with power-type nonlinearities of odd order, a cubic NLS with a derivative in the nonlinearity, and the modified Benjamin–Ono equation.

- 80 佐々木浩宣 (千葉大理)* Remark on the scattering operator for the cubic nonlinear Dirac equation in three space dimensions 10
 Hironobu Sasaki (Chiba Univ.) Remark on the scattering operator for the cubic nonlinear Dirac equation in three space dimensions

概要 We study the scattering operator S for the three dimensional Dirac equation with a cubic nonlinearity. It has been proved that S can be defined on a neighborhood of 0 in the Sobolev space $H^{s_0}(\mathbb{R}^3; \mathbb{C}^4)$ for any $s_0 > 1$. We prove that for any $M \in \mathbb{N}$ and $s \geq \max\{s_0, M\}$, there exists some neighborhood U of 0 in the weighted Sobolev space $H^{s,M}(\mathbb{R}^3; \mathbb{C}^4)$ such that $S(U) \subset H^{s,M}(\mathbb{R}^3; \mathbb{C}^4)$.

- 81 成 亥 隆 恭 (京 大 理) 絶対値べき乗型非線形シュレディンガー方程式の解のライフスパンと局
池 田 正 弘 (京 大 理) 所解の非存在について 10
Takahisa Inui (Kyoto Univ.) Remark on the lifespan of solutions and non-existence of local solution
Masahiro Ikeda (Kyoto Univ.) for a nonlinear Schrödinger equation

概要 We consider the nonlinear Schrödinger equation with p -th power absolute value nonlinearity. Let the initial value λf belong to $H^s(\mathbb{R}^d)$ where $\lambda > 0$ is a parameter and $s = 0$ or 1 . It is well known that this is locally well-posed in H^s when $1 < p \leq 1 + 4/(d - 2s)$. In the case $1 < p \leq 1 + 4/(d - 2s)$, we prove that if λ is sufficiently large and $f \in H^s$ has a singularity near the origin, then the H^s -solution blows up in finite time and the lifespan has the upper bound by a polynomial order of λ . In the opposite case $p > 1 + 4/(d - 2s)$, we also prove that if $f \in H^s$ has a singularity near the origin, then the local H^s -solution does not exist.

- 82 瓜 屋 航 太 (東 北 大 理)* 三波相互作用をもつ非線形 Schrödinger 方程式系に対する終値問題 10
Kota Uriya (Tohoku Univ.) Final state problem for a system of nonlinear Schrödinger equations
with three wave interaction

概要 In this talk, we consider the final state problem for a system of nonlinear Schrödinger equations with three wave interaction in two dimensions. In our previous study, we constructed a solution of a two component system which describes the mass transition phenomenon by using the hyperbolic functions. We show the existence of a solution to a three component system describing the mass transition phenomenon periodically in time by using the Jacobi elliptic functions.

- 83 中 村 誠 (山 形 大 理)* On the Cauchy problem for nonlinear Schrödinger equations in de Sitter
spacetime 10
Makoto Nakamura (Yamagata Univ.) On the Cauchy problem for nonlinear Schrödinger equations in de Sitter
spacetime

概要 The Cauchy problem for Schrödinger equations with weighted nonlinear terms is considered. Local and global solutions are shown in Sobolev spaces.

- 84 若 杉 勇 太 (阪 大 理)* Critical exponent for the Cauchy problem to the weakly coupled damped
西 原 健 二 (早 大 政 経) wave system 10
Yuta Wakasugi (Osaka Univ.) Critical exponent for the Cauchy problem to the weakly coupled damped
Kenji Nishihara (Waseda Univ.) wave system

概要 In this talk, we consider a system of weakly coupled semilinear damped wave equations. We determine the critical exponent for any space dimensions. We also give estimates of the lifespan of solutions from above for subcritical nonlinearities.

- 85 三好啓也 (早大理工) Convergence of hydrodynamical limits for generalized Carleman models
堤正義 (早大理工) 10
Hironari Miyoshi (Waseda Univ.) Convergence of hydrodynamical limits for generalized Carleman models
Masayoshi Tsutsumi (Waseda Univ.)

概要 We consider a generalized Carleman's model describing the time evolution of one dimensional gas composed of two kinds of particles that move parallel to x -axis with constant and equal speeds, either in the positive x -direction with density u , or in the negative x -direction with density v with initial and homogeneous boundary conditions. The existence of time global solutions in $L^1(0,1) \times L^1(0,1)$ and a nonlinear diffusion limit under the usual parabolic scaling is established for this problem.

- 86 若狭恭平 (北大理)* 1次元空間における重みつき非線形項をもつ波動方程式の解の最大存在時間 10
Kyouhei Wakasa (Hokkaido Univ.) The lifespan of solutions to nonlinear wave equations with weighted functions in 1D

概要 We consider the initial value problem for nonlinear wave equations with weighted functions in one space dimension. Kubo & Osaka & Yazici (2013) showed that the solution exists globally in time if the initial data are odd functions. On the other hand, they showed that the solution blows up in finite time if the initial data are not odd functions. Also, they have obtained the estimate of the lifespan which is the maximal existence time of solutions, however the sharpness was not clarified. Our aim in this talk is to get the sharp upper and lower bounds of the lifespan in such case.

- 87 谷口晃一 (中大理工)* 外部領域におけるポテンシャル項を持つ半線形波動方程式の散乱問題 .. 10
岩渕司 (中大理工)
松山登喜夫 (中大理工)
Koichi Taniguchi (Chuo Univ.) Scattering problem for semilinear wave equation with a potential in an
Tsukasa Iwabuchi (Chuo Univ.) exterior domain
Tokio Matsuyama (Chuo Univ.)

概要 We consider the scattering problem for the initial-boundary value problem to semilinear wave equation with a potential in an exterior domain Ω . Here, Ω is a compliment of a convex and compact obstacle \mathcal{O} in \mathbb{R}^3 . First, we will inform global existence theorems, and then, wave operators and scattering operators will be constructed.

16:30~17:30 特別講演

岡部 真也 (東北大理) 四階放物型方程式に対する障害物問題

Shinya Okabe (Tohoku Univ.) A fourth order parabolic obstacle problem

概要 In this talk, we show a recent joint work with M. Novaga (Pisa University) on the obstacle problem for the parabolic biharmonic equation. We let $\Omega \subset \mathbb{R}^N$ be a bounded domain with smooth boundary, and we let $f : \Omega \rightarrow \mathbb{R}$ be the obstacle function satisfying

$$f \in C^2(\bar{\Omega}), \quad f < 0 \quad \text{on} \quad \partial\Omega.$$

For an initial datum $u_0 : \Omega \rightarrow \mathbb{R}$ such that

$$u_0 \in H_0^2(\Omega), \quad u_0 \geq f \quad \text{a.e. in } \Omega,$$

we shall consider the following fourth order parabolic obstacle problem:

$$\begin{aligned} u_t(x, t) + \Delta^2 u(x, t) &\geq 0 && \text{in } \Omega \times \mathbb{R}_+, \\ u_t(x, t) + \Delta^2 u(x, t) &= 0 && \text{in } \{(x, t) \in \Omega \times \mathbb{R}_+ \mid u(x, t) > f(x)\}, \\ u(x, t) = 0, \quad \nabla u(x, t) \cdot \nu^\Omega(x) &= 0 && \text{on } \partial\Omega \times \mathbb{R}_+, \\ u(x, t) &\geq f(x) && \text{in } \Omega \times \mathbb{R}_+, \\ u(x, 0) &= u_0(x) && \text{in } \Omega, \end{aligned}$$

where ν^Ω denotes the unit outer normal of $\partial\Omega$.

The purpose of this talk is to investigate the regularity properties of solutions to the obstacle problem via an implicit time discretization. Since our motivation for the problem rose from a geometric obstacle problem, we wish to mention the geometric obstacle problem.

実函数論

9月27日(土) 第I会場

9:00~11:45

- 1 渡辺 俊一 (日大理工非常勤) 順序集合における拡張された縮小写像タイプの不動点定理について 15
豊田 昌史 (玉川大工)
Toshikazu Watanabe (Nihon Univ.) On fixed point theorems for generalized contractive type mappings in
Masashi Toyoda (Tamagawa Univ.) partially ordered sets

概要 Nieto and Lopez consider fixed point theorems for contractive mappings in partially ordered sets. On the other hand, Weissinger proved the contractive mapping which include the Banach contraction mappings. In this paper, we consider fixed point theorems for Weissinger type contractive mappings and Kannan mappings in partially ordered sets.

- 2 木村 泰紀 (東邦大理) 完備測地距離空間上の写像族に対する共通不動点近似 15
Yasunori Kimura (Toho Univ.) Approximation of a common fixed point of mappings on a complete
geodesic space

概要 The shrinking projection method is an approximation scheme of a common fixed point of mappings, and it is known that this method has a kind of rigidity toward the calculation errors. In this talk, we consider the shrinking projection method with errors in the setting of complete geodesic spaces and discuss the convergence property of this scheme.

- 3 松下 慎也 (秋田県立大) Douglas–Rachford 法について 15
徐 粒 (秋田県立大)
Shin-ya Matsushita (Akita Pref. Univ.) On Douglas–Rachford method
Li Xu (Akita Pref. Univ.)

概要 In this talk, we are concerned with the Douglas–Rachford method for solving the convex feasibility problem.

- 4 厚 芝 幸子 (山梨大教育人間) Convergence theorems for nonlinear mappings by Halpern’s type iterations 15
Sachiko Atsushiba Convergence theorems for nonlinear mappings by Halpern’s type iterations
(Univ. of Yamanashi) tions

概要 In this talk, we study Halpern’s type iterations for nonlinear mappings in Banach spaces. Then, we prove strong convergence theorems for uniformly asymptotically regular nonexpansive semigroups in Banach spaces. Furthermore, we give some convergence theorems for the nonlinear mappings.

- 5 加藤 幹雄 (九工大*) On the uniform non- ℓ_1^n -ness of direct sums of Banach spaces 15
 田村 高幸 (千葉大人文社会)
 Mikio Kato (Kyushu Inst. of Tech.*) On the uniform non- ℓ_1^n -ness of direct sums of Banach spaces
 Takayuki Tamura (Chiba Univ.)

概要 We shall characterize the uniform non- ℓ_1^n -ness of direct sums of Banach spaces.

- 6 徳永 清久 (福岡工大情報科学研) Riemann 積分の定義に基づく部分積分の証明 15
 Kiyohisa Tokunaga Proof of integral by parts based on the definition of Riemann integral
 (Fukuoka Inst. of Tech.)

概要 The definite integral is defined as the limit at infinity of sum of sequences. The well-known theorem of single integral by parts for a 1-variable function is derived from this definition.

- 7 河邊 淳 (信州大工) 分布関数で定まる非線形積分汎関数の有界収束定理 15
 Jun Kawabe (Shinshu Univ.) Bounded convergence theorem for distribution-based nonlinear integral
 functionals

概要 We introduce a new notion of the perturbation of nonlinear integral functionals to formulate a functional form of the convergence theorems for nonlinear integrals in nonadditive measure theory. As its direct consequences, we obtain the bounded convergence theorems for typical nonlinear integrals, which show that the autocontinuity of a nonadditive measure is equivalent to the validity of the bounded convergence theorems for the Choquet, the Sugeno, and the Shilkret integrals as well as their symmetric and asymmetric extensions.

- 8 森藤 紳哉 (奈良女大理) オストロフスキーの不等式とその離散化 10
 Shinya Moritoh (Nara Women's Univ.) Ostrowski's inequality and its discretization

概要 A comparison of integral and discrete Ostrowski's inequalities in the plane is considered. An integral inequality is described by Legendre's elliptic integrals. A natural discrete analogue of the inequality is also given. The main point is to find a suitable decomposition of the radius in polar coordinates.

- 9 伊東 由文 (徳島大*)^b 自然統計物理学の法則 15
 Yoshifumi Ito (Univ. of Tokushima*) Laws of natural statistical physics

概要 In this paper we propose the laws of natural statistical physics and give the new formulations in the following three cases :

- (1) The case where the Schrödinger operator has only the discrete spectrum.
- (2) The case where the Schrödinger operator has only the continuous spectrum.
- (3) The case where the physical system is composed of particles moving periodically.

- 10 伊東由文 (徳島大*) 自然確率の概念 15
 Yoshifumi Ito (Univ. of Tokushima*) Concept of natural probability

概要 In this paper, we study the concepts of natural probability and natural random variable and give the new formulations.

For a given normalized L^2 -function ψ on \mathbf{R}^n , ($n \geq 1$), we define the orthogonal probability measure $\psi_A = \chi_A \psi$ on the probability space $(\mathbf{R}^n, \mathcal{M}_n, \mu)$ to be a natural probability measure. Here \mathcal{M}_n is the σ -additive family of all Lebesgue measurable sets on \mathbf{R}^n and μ is the σ -additive measure

$$\mu(A) = \int_A |\psi(\mathbf{r})|^2 d\mathbf{r}, \quad (A \in \mathcal{M}_n).$$

We define the vector-valued natural random variable $\mathbf{r} = \mathbf{r}(\omega)$ on a certain probability space $\Omega(\mathcal{B}, P)$ whose probability distribution law is defined by the L^2 -density ψ so that the fundamental relation $P(\{\rho \in \Omega; \mathbf{r}(\rho) \in A\}) = \mu(A)$ holds. Further we study the fundamental properties of these concepts.

Further we study two other cases of these concepts.

12:10~12:30 2014年度解析学賞授賞式

14:15~16:20

- 11 齋藤洋樹 (埼玉大理) Takeya maximal operator (no dilation) with radial weights on the plane 15
 Hiroki Saito (Saitama Univ.) Takeya maximal operator (no dilation) with radial weights on the plane

概要 We consider the weighted “small” Takeya maximal operator is defined to be

$$K_{N,w}^a f(x) := \sup_{x \in R} \frac{1}{w(R)} \int_R |f(y)| w(y) dy$$

where the supremum is taken over all rectangles with sizes $a \times aN$ containing x . In this talk, we shall prove

$$\|K_{N,w}^a f\|_{L^2(w)} \leq C \sqrt{\log N} \|f\|_{L^2(w)},$$

for a certain radial weight w .

- 12 飯田毅士 (福島工高専) Weighted estimates of higher order commutators generated by BMO -functions and the fractional integral operator on Morrey spaces 15
 Takeshi Iida Weighted estimates of higher order commutators generated by BMO -
 (Fukushima Nat. Coll. of Tech.) functions and the fractional integral operator on Morrey spaces

概要 The purpose of this paper is to investigate the weighted estimates of commutators generated by BMO -functions and the fractional integral operator on Morrey spaces. In particular, the conditions of indices is the Olsen type conditions. The main result is generalized the Sawano Sugano and Tanaka result to a weighted setting.

- 13 菅野 聡子 (神戸工高専)* 高階のシュレーディンガー型のカルデロン・ジグムント作用素について 15
 Satoko Sugano On a Calderón–Zygmund operator of higher order Schrödinger type
 (Kobe City Coll. of Tech.)

概要 We consider higher order Schrödinger type operators with nonnegative potentials. We assume that the potential belongs to the reverse Hölder class which includes nonnegative polynomials. We show that an operator of higher order Schrödinger type is a Calderón–Zygmund operator.

- 14 松岡 勝男 (日大経済) On the generalization of fractional integrals and λ -CMO spaces 10
 Katsuo Matsuoka (Nihon Univ.) On the generalization of fractional integrals and λ -CMO spaces

概要 In this talk, we will extend the boundedness of fractional integrals I_α for the non-homogeneous central Morrey spaces $B^{p,\lambda}(\mathbb{R}^n)$ over the whole of λ such that $-n/p \leq \lambda < \infty$. In order to show the above, we will introduce the generalized λ -CMO spaces $\Lambda_{p,\lambda}^{(d)}(\mathbb{R}^n)$ and the generalized fractional integrals $\tilde{I}_{\alpha,d}$.

- 15 澤野 嘉宏 (首都大東京理工)^b Atomic decomposition for Morrey spaces 15
 飯田 毅士 (福島工高専)
 田中 仁 (東大数理)
 Yoshihiro Sawano (Tokyo Metro. Univ.) Atomic decomposition for Morrey spaces
 Takeshi Iida
 (Fukushima Nat. Coll. of Tech.)
 Hitoshi Tanaka (Univ. of Tokyo)

概要 We propose here a non-smooth decomposition of Morrey spaces. We content ourselves with stating two main decomposition results.

- 16 澤野 嘉宏 (首都大東京理工)^b Wavelet characterization and modular inequalities for weighted Lebesgue
 出来 光夫 (岡山大教育) spaces with variable exponent 15
 中井 英一 (茨城大理)
 Yoshihiro Sawano (Tokyo Metro. Univ.) Wavelet characterization and modular inequalities for weighted Lebesgue
 Mitsuo Izuki (Okayama Univ.) spaces with variable exponent
 Eiichi Nakai (Ibaraki Univ.)

概要 We consider the wavelet characterization of the space $L^{p(\cdot)}(w)$ with $w \in A_{p(\cdot)}$.

- 17 前田文之(広島大)* Grand Musielak–Orlicz–Morrey 空間の関数のポテンシャルに関する Sobolev 型と Trudinger 型の不等式 15
 水田義弘(広島工大工)
 大野貴雄(大分大教育福祉)
 下村哲(広島大教育)
 Fumi-Yuki Maeda (Hiroshima Univ.*) Sobolev and Trudinger type inequalities for potentials of functions in grand Musielak–Orlicz–Morrey spaces
 Yoshihiro Mizuta (Hiroshima Inst. of Tech.)
 Takao Ohno (Oita Univ.)
 Tetsu Shimomura (Hiroshima Univ.)

概要 We define (generalized) grand Musielak–Orlicz–Morrey space on a bounded open set in \mathbf{R}^N and give a Sobolev type inequality as well as a Trudinger type inequality for potentials of functions in such spaces.

- 18 前田文之(広島大)* Growth properties of Musielak–Orlicz integral means for Riesz potentials 15
 水田義弘(広島工大工)
 下村哲(広島大教育)
 Fumi-Yuki Maeda (Hiroshima Univ.*) Growth properties of Musielak–Orlicz integral means for Riesz potentials
 Yoshihiro Mizuta (Hiroshima Inst. of Tech.)
 Tetsu Shimomura (Hiroshima Univ.)

概要 In this talk we are concerned with growth properties of integral means for Riesz potentials of functions in Musielak–Orlicz spaces.

16:40~17:40 特別講演

- 三谷健一(岡山県立大情報工) バナッハ空間の幾何学的定数について
 Kenichi Mitani (Okayama Pref. Univ.) On geometrical constants of Banach spaces

概要 In this talk, we describe some recent results on James constant, von Neumann–Jordan constant and the skewness of Banach spaces. These constants play an important role in the description of various geometrical structures of Banach spaces. We first discuss James and von Neumann–Jordan constants for absolute normalized norms. We also exhibit some connections between the skewness and James constant for general Banach spaces.

9月28日(日) 第I会場

9:00~11:50

- 19 伊藤昭夫(近畿大工) 終了時刻が未知関数に依存する日本酒醸造過程モデルに対する最適制御
 剣持信幸(佛教大教育) 問題について 15
 村瀬勇介(名城大理工)

Akio Ito (Kinki Univ.) Optimal control problems for mathematical model for the process of
 Nobuyuki Kenmochi (Bukkyo Univ.) brewing Japanese Sake with unknown terminal time
 Yusuke Murase (Meijo Univ.)

概要 Our mathematical model for the process of brewing Japanese Sake is a system of differential equations with a constraint condition, and whose terminal time is depend upon the solution self. The system was solved by Y. Murase and A. Ito in 2013 with putting homogeneous Neumann boundary conditions and Robin boundary condition for heat equation. Our next objective is optimal control problem for the model. In this talk, we talk about existence theorem of optimal control and optimal solution characterized by certain cost function which is corresponding to phenomena.

- 20 藤江健太郎(東京理大理) 癌浸潤現象に関するある数理モデルの時間大域解の存在と漸近挙動 15
 伊藤昭夫(近畿大工)
 M. Winkler (Univ. Paderborn)
 横田智巳(東京理大理)

Kentarou Fujie (Tokyo Univ. of Sci.) Global existence and asymptotic behavior of solutions to a model for
 Akio Ito (Kinki Univ.) tumor invasion
 Michael Winkler (Univ. Paderborn)
 Tomomi Yokota (Tokyo Univ. of Sci.)

概要 This talk deals with a chemotaxis system modeling tumor invasion in a smoothly bounded domain $\Omega \subset \mathbb{R}^N$, $N \leq 3$. This system was recently proposed as a modified tumor invasion model in which the role of an active extracellular matrix is taken into consideration. Moreover, the existence and uniqueness of local-in-time classical solutions to the model was proved in the general case $N \in \mathbb{N}$. This talk clarifies boundedness and asymptotic behavior of solutions to the model in the physically relevant case when $N \leq 3$.

- 21 横田智巳(東京理大理) 増殖項をもつ準線形非退化放物・楕円型 Keller–Segel 系への作用素論的
 吉野徳晃(東京理大理) アプローチ 15
 Tomomi Yokota (Tokyo Univ. of Sci.) Operator-theoretic approach to a quasilinear nondegenerate parabolic-
 Noriaki Yoshino (Tokyo Univ. of Sci.) elliptic Keller–Segel system with growth term

概要 This talk deals with the chemotaxis system with nonlinear diffusion and superlinear growth term $f(b) = |b|^{\alpha-1}b$ when $n \leq 3$. It is shown that if $\alpha \leq 4$ ($n = 1$), $\alpha < 1 + \frac{4}{n}$ ($n = 2, 3$), then there exists a local solution to this system for any large data. In the case of Lipschitz growth, Marinoschi (*J. Math. Anal. Appl.* 2013; 402: 415–439) established the existence of local solutions to this system with sufficiently small initial data and showed that under a stronger assumption on the chemotactic sensitivity there exists a global solution with large initial data. This talk develops the local solvability with Lipschitz growth to the one with superlinear growth and allows the system to have a local solution with large initial data without any stronger assumption. The key to including the superlinear growth lies in the Yosida approximation of f .

- 22 清水 翔司 (早大理工) The solvability of complex Ginzburg–Landau equation focusing on parabolicity
大谷 光春 (早大理工) 15
Shoji Shimizu (Waseda Univ.) The solvability of complex Ginzburg–Landau equation focusing on parabolicity
Mitsuharu Ôtani (Waseda Univ.) icity

概要 The main purpose of this talk is to study Cauchy problem for complex Ginzburg–Landau equation (CGL) by focusing on its parabolicity. In former studies, (CGL) is always treated as a single equation over complex spaces. However in this approach, one can not make most use of the parabolicity of (CGL). So here we regard the real part of (CGL) as the principal part governed by the subdifferential operator and the pure imaginary part as its perturbations. By this approach we can show a new smoothing effect of (CGL) which ameliorates former results.

- 23 内田 俊 (早大理工) Global attractor of some autonomous double-diffusive convection system
大谷 光春 (早大理工) 15
Shun Uchida (Waseda Univ.) Global attractor of some autonomous double-diffusive convection system
Mitsuharu Ôtani (Waseda Univ.)

概要 We consider the large time behavior of solutions of some system which describes double-diffusive convection in some porous medium. In our previous works, it has already been showed that there exist a unique global solution of this system under the homogeneous Dirichlet or Neumann boundary condition. In this talk, we discuss the existence of global attractor of the system. Especially, for the homogeneous Neumann boundary condition case, we need some restriction on the external forces and mass of the initial data. We focus on the differences between two cases, i.e. Dirichlet and Neumann boundary condition cases.

- 24 都築 寛 (東京理大理) Solvability of heat equations with hysteresis coupled with Navier–Stokes equations in 2D domains 15
Yutaka Tsuzuki (Tokyo Univ. of Sci.) Solvability of heat equations with hysteresis coupled with Navier–Stokes equations in 2D domains

概要 This talk is concerned with a system of heat equations with hysteresis and Navier–Stokes equations in two-dimensional domains. There are some studies on the system of heat equations and Navier–Stokes equations in which the temperature (the solution to heat equations) is controlled by obstacle functions. In this talk, the control of the temperature is precisely described by introducing hysteresis. This talk provides the existence of solutions along the abstract theory for evolution equations with subdifferential operators.

- 25 赤木 剛朗 (神戸大システム情報) Doubly nonlinear evolution equations in variable exponent Lebesgue spaces 15
G. Schimperna (Univ. di Pavia) spaces
Goro Akagi (Kobe Univ.) Doubly nonlinear evolution equations in variable exponent Lebesgue spaces
Giulio Schimperna (Univ. di Pavia) spaces

概要 This talk is concerned with the Cauchy–Dirichlet problem for a doubly nonlinear parabolic equation involving variable exponents and provides some theorems on existence and regularity of strong solutions. In the proof of these results, we also analyze the relations occurring between Lebesgue spaces of space-time variables and Lebesgue–Bochner spaces of vector-valued functions, with a special emphasis on measurability issues and particularly referring to the case of space-dependent variable exponents.

- 26 松本敏隆 (広島大理) 弱連続作用素に対する抽象的 Cauchy 問題について 15
 田中直樹 (静岡大理)
 Toshitaka Matsumoto Abstract Cauchy problem for weakly continuous operators
 (Hiroshima Univ.)
 Naoki Tanaka (Shizuoka Univ.)

概要 The abstract Cauchy problem for weakly continuous operators is discussed in a general Banach space. A class of weakly continuous operators is introduced by using sets of functionals. An existence theorem for weakly differentiable solutions and its applications to concrete PDEs are given.

- 27 登口大 (早大教育) 確率保存型方程式に対する非斉次 Dirichlet 問題の解の存在定理 15
 小林和夫 (早大教育)
 Dai Noboriguchi (Waseda Univ.) An existence theorem for a nonhomogeneous Dirichlet problem for a
 Kazuo Kobayasi (Waseda Univ.) stochastic scalar conservation law

概要 We consider the initial-boundary value problem for a randomly forced scalar conservation law with a multiplicative noise on a bounded convex domain D in R^d : $du + \operatorname{div}(A(u)) dt = \Phi(u) dW(t)$ in $(0, T) \times D$, $u(0, \cdot) = u_0(\cdot)$ on D , $u \cong u_b$ on $(0, T) \times \partial D$. We introduce a notion of kinetic formulations in which the kinetic defect measures on the boundary of a domain are truncated. In such a kinetic formulation we give a result of uniqueness and existence.

- 28 側島基宏 (Univ. of Salento) L^p -theory for second-order elliptic operators with unbounded coefficients in an endpoint class 15
 Motohiro Sobajima (Univ. of Salento) L^p -theory for second-order elliptic operators with unbounded coefficients in an endpoint class

概要 The m -accretivity and m -sectoriality of the minimal and maximal realizations of second-order elliptic operators of the form $Au = -\operatorname{div}(a\nabla u) + F \cdot \nabla u + Vu$ in $L^p(\mathbb{R}^N)$ are shown, where the coefficients a , F and V are unbounded. The result may be regarded as an endpoint assertion of my previous result in 2012 and an improvement of that in Metafuno et al. (2010). Moreover, an L^p -generalization of a self-adjoint problem posed by T. Kato in 1981 is also discussed.

14:15~15:40

- 29 渡邊 紘 (サレジオ工高専) Allen-Cahn 型方程式と結晶粒界モデルによる連立系の時刻無限大における解挙動 15
 白川 健 (千葉大教育)
 Hiroshi Watanabe (Salesian Polytech.) Large time behavior for mathematical models of grain boundary motions
 Ken Shirakawa (Chiba Univ.) involving isothermal solidifications

概要 We consider coupled systems of Allen-Cahn type equations and PDE models of grain boundary motions. In this light, these systems can be regarded as interactive mathematical models of the grain boundaries motions under isothermal solidifications. In this talk, we discuss the large time behavior of solutions to our systems. Moreover, we also mention about some improvements to approximating arguments, associated with the Γ -convergence of governing energies.

- 30 白川 健 (千葉大教育) 凝固現象と結晶粒界現象の等温系カップリングモデルにおけるエネルギー
渡邊 紘 (サレジオ工高専) 消散解 15
Ken Shirakawa (Chiba Univ.) Energy-dissipative solutions to models of grain boundary motions under
Hiroshi Watanabe (Salesian Polytech.) isothermal solidifications

概要 In this talk, we consider parabolic type systems of the ϕ - η - θ model, that is known as a phase-field model of grain boundary motion under an isothermal solidification. The main objective is to confirm the reproduction ability of the smoothing effect and energy-dissipation in our systems. With this view, the main results will be concerned with the existence of special kind of solution, called energy-dissipative solution. Although each of our systems has specific difficulties in mathematics, the main results are proved on the basis of the time-discretization for a common approximating problem. As a consequence, we provide a uniform finding method for energy-dissipative solutions to wide scope of nonstandard parabolic systems, associated with the ϕ - η - θ model.

- 31 山崎 教昭 (神奈川大工) Singular limit of Allen–Cahn equation with constraints and its Lagrange
深尾 武史 (京都教育大教育) multiplier 15
M. Hassan Farshbaf-Shaker
(WIAS)
Noriaki Yamazaki (Kanagawa Univ.) Singular limit of Allen–Cahn equation with constraints and its Lagrange
Takeshi Fukao (Kyoto Univ. of Edu.) multiplier
Mohammad Hassan Farshbaf-Shaker
(WIAS)

概要 We consider the Allen–Cahn equation with constraint. Our constraint is the subdifferential of the indicator function on the closed interval, which is the multivalued function. In this talk we give the characterization of the Lagrange multiplier to our equation. Moreover, we consider the singular limit of our system and clarify the limit of the solution and the Lagrange multiplier to our problem.

- 32 深尾 武史 (京都教育大教育) Cahn–Hilliard equation with dynamic boundary conditions and mass
P. Colli (Univ. di Pavia) constraints on the boundary 15
Takeshi Fukao (Kyoto Univ. of Edu.) Cahn–Hilliard equation with dynamic boundary conditions and mass
Pierluigi Colli (Univ. di Pavia) constraints on the boundary

概要 The well-known Cahn–Hilliard equation has the structure of the mass conservation under the suitable boundary condition. In the case when the equation is treated with the dynamic boundary condition, the total mass on the inside of the domain and its trace on the boundary should be conserved. The new issue is the setting of the mass constraint on the boundary. The effect of this additional constraint is characterized by the Lagrange multiplier, that is, there are two Lagrange multipliers, one is for the inside of the domain, the other is for the boundary.

- 33 愛木 豊彦 (日本女大理) 水分吸着過程を記述する自由境界問題の解の時間無限大の挙動について
 村瀬 勇介 (名城大理工) 15
 Toyohiko Aiki (Japan Women's Univ.) Large time behavior of a solution to the free boundary problem describ-
 Yusuke Murase (Meijo Univ.) ing adsorption phenomena

概要 In this talk we consider the free boundary problem which is a mathematical model for adsorption phenomena in porous media. On this problem the modeling process was proposed and the well-posedness was already discussed. Here, under some condition for the growth rate of wetting area we show a result on large time behavior of a solution to the problem. Also, we can provide some conjectures on the rate of convergence of the solution for large time from observations for our numerical results.

16:00~17:00 特別講演

- 吉井 健太郎 (東京理大理) 双曲型線形発展方程式について
 Kentarou Yoshii (Tokyo Univ. of Sci.) On the hyperbolic type linear evolution equations

概要 Set $I := [0, T]$. Let $\{A(t); t \in I\}$ be a family of closed linear operators in a complex Hilbert space X . Then we consider the abstract Cauchy problem for linear evolution equations of the form

$$(ACP) \quad \begin{cases} (d/dt)u(t) + A(t)u(t) = f(t), & t \in I, \\ u(0) = u_0 \in Y. \end{cases}$$

Here Y is another Hilbert space, embedded densely and continuously in X .

One of those methods to solve (ACP) depends on the unique existence of *evolution operator* for (ACP)

$$\{U(t, s); 0 \leq s \leq t \leq T\}.$$

If the evolution operator for (ACP) is known, then it is expected that the function $u(\cdot)$ given by the Duhamel formula

$$u(t) := U(t, 0)u_0 + \int_0^t U(t, s)f(s) ds$$

is a unique solution of (ACP).

Our purpose in this talk is to obtain the sufficient condition guaranteeing the existence of evolution operator.

函数解析学

9月25日(木) 第IV会場

14:15~16:35

- 1 新國裕昭(前橋工科大)* Spectral band structure of periodic Schrödinger operators on generalized degenerate zigzag nanotubes 15
 Hiroaki Niikuni Spectral band structure of periodic Schrödinger operators on generalized degenerate zigzag nanotubes
 (Maebashi Inst. of Tech.)

概要 In this talk, we consider periodic Schrödinger operators on the dumbbell-like metric graph, which is a periodic graph consisting of lines and rings. Let one line and two rings be in the basic period cell. We see the relationship between the structure of graph and the band-gap spectrum.

- 2 佐々木浩宣(千葉大理) 平均場シュレーディンガー作用素のスペクトル解析 10
 清水翔之(東工大理工)
 鈴木章斗(信州大工)
 Hironobu Sasaki (Chiba Univ.) Spectral analysis for mean-field Schrödinger operators
 Shoji Shimizu (Tokyo Tech)
 Akito Suzuki (Shinshu Univ.)

概要 We consider a time-dependent Schrödinger operator H_t which appears in recent study for mean-field analysis of quantum many-body systems. We show some basic spectral properties of H_t .

- 3 伊藤将吾(信州大理工) 遠方で減衰しないポテンシャルをもつ離散シュレーディンガー作用素の
 鈴木章斗(信州大工) スペクトル 15
 Shougo Ito (Shinshu Univ.) The spectrum of a discrete Schrödinger operator with a non-decaying
 Akito Suzuki (Shinshu Univ.) potential

概要 In this talk, we study the spectrum of a discrete Schrödinger operator $H = H_0 + V$ on $\mathbb{Z}^d (d \geq 2)$ with a potential defined by $V = v \sum_{x_n=(n,0,\dots,0), n \in \mathbb{Z}} |x_n\rangle\langle x_n| (v > 0)$. In the case of $d = 2, 3$, there exists a positive γ_d^+ such that: (1) if $v \leq \gamma_d^+$, then the spectrum $\sigma(H)$ equals $[-2d, \lambda^+]$ with some $\lambda^+ > 2d$; (2) if $v > \gamma_d^+$, then $\sigma(H) = [-2d, 2d] \cup [\lambda^-, \lambda^+]$ with some $\lambda^+ > \lambda^- > 2d$. For $d \geq 4$, there exists a positive γ_d^- such that $\sigma(H) = [-2d, 2d]$ holds if $v \leq \gamma_d^-$.

- 4 伊藤 宏(愛媛大理工) 遠方で発散するポテンシャルをもつ Dirac 作用素の非相対論的極限について 15
 Hiroshi Ito (Ehime Univ.) The nonrelativistic limit for Dirac operators with a potential diverging at infinity

概要 We consider Dirac operators with a bounded magnetic potential and an unbounded electric potential diverging at infinity. Under the assumption that the potentials are dilation analytic, resonances of the operator are defined as eigenvalues of the dilated operator. In this talk we show that if the speed of light goes to infinity (the nonrelativistic limit), resonances of the Dirac operator exist near resonances of two Pauli operators.

- 5 門脇光輝 (愛媛大理工) 二層媒質中の波動伝播に対するレゾルベントの空間遠方での漸近挙動 .. 15
磯崎洋 (筑波大数理物質)
渡辺道之 (新潟大教育)
Mitsuteru Kadowaki (Ehime Univ.) Asymptotic behavior in far field of the resolvent for wave propagation
Hiroshi Isozaki (Univ. of Tsukuba) in two-layered media
Michiyuki Watanabe (Niigata Univ.)

概要 In this talk, we report a result for asymptotic behavior in far field of the resolvent for wave propagation in two-layered media of three dimensions. It is well-known that asymptotic behavior in far field of the resolvent for wave propagation in homogeneous medium of whole space. Moreover that is very useful to study obstacle or Schrödinger scattering and inverse scattering problem. However, it was not established that sharp study for asymptotic behavior in far field of the resolvent for wave propagation in two-layered media because of the existence of refracted (transmitted) waves.

- 6 二口伸一郎 (北大理) 非有界作用素に対する複素平面上の Time-ordered exponential と Gell-
白井耕太 (北大理) Mann-Low 公式 10
Shinichiro Futakuchi (Hokkaido Univ.) Time-ordered exponential on the complex plane for unbounded opera-
Kouta Usui (Hokkaido Univ.) tors and Gell-Mann-Low formula

概要 The time-ordered exponential plays a fundamental role in perturbative calculation. We introduce a method for constructing the time-ordered exponential on the complex plane for unbounded operators, and its applications to quantum field theory.

- 7 船川大樹 (北大理) Existence of ground states for a Wess-Zumino model 10
Daiju Funakawa (Hokkaido Univ.) Existence of ground states for a Wess-Zumino model

概要 Supersymmetry is a symmetry corresponding to replacing bosons by fermions. We investigate the Wess-Zumino model which is one of the simplest example of supersymmetric quantum field theory. We show the existence of the ground state in two different cases: where the particles are massive and where these are massless.

- 8 長谷部高広 (北大理) 自由 Lie 代数におけるキュムラントと Campbell-Hausdorff 公式 15
F. Lehner (Graz Univ. of Tech.)
Takahiro Hasebe (Hokkaido Univ.) Cumulants for free Lie algebras and Campbell-Hausdorff formula
Franz Lehner (Graz Univ. of Tech.)

概要 We find a relation between cumulants in noncommutative probability and the Campbell-Hausdorff formula in free Lie algebras. Combinatorics of set compositions (or ordered set partitions) is the main tool in the result.

- 9 柳研二郎 (山口大理工) 非エルミート型の不確定性関係 15
Kenjiro Yanagi (Yamaguchi Univ.) Uncertainty relations for non-hermitian type

概要 Recently Dou and Du obtained several uncertainty relations for non-hermitian type. We show that their results can be given as corollaries of our uncertainty relations for generalized metric adjusted skew informations.

17:00~18:00 特別講演

水町 徹 (九大数理) KP-II 方程式の line soliton 解の安定性
 Tetsu Mizumachi (Kyushu Univ.) Stability of line solitons for the KP-II equation

概要 In this talk, we discuss nonlinear stability of line soliton solutions of the KP-II equation. The KP-II equation is a two dimensional generalization of the KdV equation which takes into account of slow variations of waves in the transversal direction.

If the perturbation to line solitons are periodic in the transversal direction, the line soliton solution is stable in the same manner as 1-soliton solutions of the KdV equation.

However, with respect to localized transversal perturbations in \mathbb{R}^2 , the modulating speed and the phase shift cannot be uniform in the transversal direction. Modulation equations of the local amplitude and the local phase shift of the crest of the line solitons are described by a system of 1D wave equations with diffraction terms.

9月26日(金) 第IV会場

10:00~11:50

10 田端 亮 (広島大理) $n \rightarrow \infty$ のときの $n \times n$ Immanants の挙動 15
 Ryo Tabata (Hiroshima Univ.) The behavior of $n \times n$ Immanants as $n \rightarrow \infty$

概要 When λ is a Young diagram, $T_\lambda = \max T_\lambda(A) = \max_{A \geq 0} (\bar{d}_\lambda(A) - \det A) / (\text{per} A - \det A)$ measures how large $\bar{d}_\lambda(A)$ can be on the determinant-permanent line segment. The celebrated permanental dominance conjecture asserts that $T_\lambda \leq 1$ for all λ . Conjecturally the maximum value is attained by $Y_n = (n/(n-1)\delta_{ij} - 1/(n-1))$ and we study the behavior of $T_\lambda(Y_n)$ as $|\lambda| \rightarrow \infty$. We give precise values when λ 's are hooks, random young diagrams, and $\lim_{n \rightarrow \infty} T_{(k^n)}(Y_n)$. We also discuss the relation with Pierce's conjecture, which says that the maximum permanent of singular correlation matrices is attained by Y_n .

11 中島 秀斗 (九大数理) 等質開凸錐の基本相対不変式の明示的公式 15
 Hideto Nakashima (Kyushu Univ.) Explicit formula of the basic relative invariants of homogeneous cones

概要 In a study of homogeneous convex cones, the basic relative invariants play an important role. It is known that they are obtained inductively as the irreducible factors of Vinberg polynomials (Ishi 2001). In this talk, we give an explicit formula which calculates the basic relative invariants all at once with Vinberg polynomials by using data related to the homogeneous cones.

12 松木 敏彦 (龍谷大文) 奇数次直交群の有限型多重旗多様体の分類 15
 Toshihiko Matsuki (Ryukoku Univ.) Classification of orthogonal multiple flag varieties of finite type

概要 Let G be the split orthogonal group of degree $2n+1$ over an arbitrary field \mathbb{F} of characteristic not 2. In this talk, we classify multiple flag varieties $G/P_1 \times \cdots \times G/P_k$ of finite type. Here a multiple flag variety is called of finite type if it has a finite number of G -orbits with respect to the diagonal action of G when $|\mathbb{F}| = \infty$.

- 13 松木敏彦(龍谷大文) 奇数次直交群の3重旗多様体の軌道分解 15
Toshihiko Matsuki (Ryukoku Univ.) Orbits on orthogonal triple flag varieties

概要 Let G be the split orthogonal group of degree $2n + 1$ over a field \mathbb{F} of characteristic not 2. In this talk, we describe G -orbits on the triple flag variety $\mathcal{M} = \text{Fl}_{(\alpha)} \times \text{Fl}_{(\beta)} \times \text{Fl}_{(n)}$. Here $\text{Fl}_{(\alpha)}$ denote the flag variety consisting of α -dimensional isotropic subspaces in \mathbb{F}^{2n+1} .

- 14 笹木集夢(東海大理) A Cartan decomposition for Cayley type homogeneous spaces 15
Atsumu Sasaki (Tokai Univ.) A Cartan decomposition for Cayley type homogeneous spaces

概要 We explain how to find an abelian B satisfying $G = KBH$ for Cayley type homogeneous spaces G/H . Here, K is a maximal compact subgroup of G . We note that G/H is non-symmetric. Then, our result gives a kind of generalizations of Cartan decomposition for symmetric spaces to non-symmetric ones.

- 15 伊師英之(名大多元数理)* 正則凸錐上のジーゲル型積分 15
Hideyuki Ishi (Nagoya Univ.) Siegel type integral on a regular convex cone

概要 The Laplace transform of a power of the determinant over the cone of positive definite symmetric matrices is called the Siegel integral. The formula is generalized to integrals over general homogeneous cones by Gindikin, while the analogous integrals are considered over regular cones of symmetric matrices with prescribed zeros by Roverato. In this talk, we introduce a new class of regular convex cones and consider the Siegel type integral over the cones, which unifies Gindikin's and Roverato's formulas.

13:10~14:10 特別講演

織田 寛(拓殖大工) 実簡約 Lie 群と次数 Hecke 環の表現論のつながり

Hiroshi Oda (Takushoku Univ.) Connections between representation theories for real reductive Lie groups and graded Hecke algebras

概要 The representation theory for real reductive Lie groups G and that for graded Hecke algebras \mathbf{H} are very similar. For example, the Langlands classification for G versus Evens' classification of irreducible \mathbf{H} -modules, and the Helgason–Fourier transform versus the Opdam–Cherednik transform. In addition Ciubotaru and Trapa constructed for various classical G functors sending G -modules to modules of the corresponding \mathbf{H} . We unify many parallel objects and phenomenons in two representation theories by introducing the notion of *radial pairs*. A radial pair is a pair of a $(\mathfrak{g}_{\mathbb{C}}, K)$ -module and an \mathbf{H} -module satisfying axioms which are formally the same as a generalized Chevalley restriction theorem and a generalized radial part formula. Related to this we construct three functors sending \mathbf{H} -modules to $(\mathfrak{g}_{\mathbb{C}}, K)$ -modules.

9月27日(土) 第IV会場

9:00~12:00

- 16 三宅啓道 On compactness in L^1 15
 Hiromichi Miyake On compactness in L^1

概要 Recently, we discussed a method of constructing a separated locally convex topology τ on L^1 with the assumption that the reference measure is σ -finite. In this talk, we show a characterization of compactness for the weak topology of L^1 associated with the topology τ and apply similar arguments to study some conditions for strong and weak compactness in L^1 , respectively. We also discuss (weak) almost periodicity of linear contractions on L^1 .

- 17 和田和幸(北大理) 空間切断の入った複素スカラー場の自己相互作用モデルにおける基底状態の存在 15
 Kazuyuki Wada (Hokkaido Univ.) Existence of a ground state for a self-interaction model of a complex scalar field with spacial cut-off

概要 We consider the Hamiltonian H of a charged scalar field with self-interaction. Here, we assume that the mass of boson is just zero. By introducing a spacial cut-off function, H is realized as a linear operator on a boson Fock space. Under certain conditions, H is bounded-below, self-adjoint. Moreover, H has a ground state for arbitrary coupling constants under an infrared regularly condition.

- 18 泉池敬司(新潟大自然)* Topological structure of the space of weighted composition operators
 大野修一(日本工大) between different Hardy spaces 15
 Kei Ji Izuchi (Niigata Univ.) Topological structure of the space of weighted composition operators
 Shūichi Ohno (Nippon Inst. of Tech.) between different Hardy spaces

概要 We consider properties related to weighted composition operators boundedly acting from the classical Hardy space H^p to H^q for $1 \leq q < p < \infty$. Especially, we shall completely determine path connected components in the set of weighted composition operators and explicitly characterize by function-theoretic properties of analytic self-maps.

- 19 大野修一(日本工大)* Composition operators related to the Dirichlet space 10
 Shūichi Ohno (Nippon Inst. of Tech.) Composition operators related to the Dirichlet space

概要 The Hilbert-Schmidtness of composition operators acting between the classical Hilbert Hardy space and the Dirichlet space is known. We here consider boundedness and compactness of composition operators acting between their spaces.

- 20 泉池 耕平 (山口大教育) Ranks of backward shift invariant subspaces of Hardy space over the bidisk 15
 Kouhei Izuchi (Yamaguchi Univ.) Ranks of backward shift invariant subspaces of Hardy space over the bidisk

概要 Let $\{\varphi_n(z)\}_{n \geq 0}$ be a sequence of inner functions satisfying that $\zeta_n(z) := \varphi_n(z)/\varphi_{n+1}(z) \in H^\infty(z)$ for every $n \geq 0$ and $\{\varphi_n(z)\}_{n \geq 0}$ has no nonconstant common inner divisors. Associated with it, we have a Rudin type invariant subspace \mathcal{M} of $H^2(\mathbb{D}^2)$. The ranks of $\mathcal{M} \ominus w\mathcal{M}$ for \mathcal{F}_z and \mathcal{F}_z^* respectively are determined, where \mathcal{F}_z is the fringe operator on $\mathcal{M} \ominus w\mathcal{M}$. Let $\mathcal{N} = H^2(\mathbb{D}^2) \ominus \mathcal{M}$. It is also proved that the rank of $\mathcal{M} \ominus w\mathcal{M}$ for \mathcal{F}_z^* equals to the rank of \mathcal{N} for T_z^* and T_w^* .

- 21 岡 康之 (釧路工高専) A characterization of the tempered distributions supported by a regular closed set on the Heisenberg group 15
 Yasuyuki Oka A characterization of the tempered distributions supported by a regular closed set on the Heisenberg group
 (Kushiro Nat. Coll. of Tech.)

概要 In this talk, we will give the characterization of the tempered distributions supported by a regular closed set on the Heisenberg group by means of the heat kernel method.

- 22 古市 茂 (日大文理) 相対作用素エントロピーの上界と下界について 10
 Shigeru Furuichi (Nihon Univ.) On bounds for relative operator entropies

概要 Recently, Zou obtained the generalized results on the bounds for Tsallis relative operator entropy. In this talk, we give precise bounds for Tsallis relative operator entropy. We also give precise bounds of relative operator entropy.

- 23 藤井 淳一 (大阪教育大教育) 基本的な作用素エントロピーについて 15
 Junichi Fujii (Osaka Kyoiku Univ.) On basic operator entropies

概要 As a tangent vector for the geodesics of a certain geometry of positive-definite matrices, relative entropies for operators were introduced and have been discussed. So I introduce basic entropies for operators (or matrices) corresponding to those in the classical information theory. We observe some good properties for these entropies.

- 24 瀬尾 祐貴 (大阪教育大教育) $1 < t < 2$ に対する Lawson–Lim–Pálfia による行列べき平均について 10
 Yuki Seo (Osaka Kyoiku Univ.) Matrix power means due to Lawson–Lim–Pálfia for $1 < t < 2$

概要 For $-1 \leq t \leq 1$, Lim–Pálfia defined a new family of operator power means of positive definite matrices and subsequently by Lawson–Lim their notion and most of their results extend to the setting of positive invertible operators on a Hilbert space. Each of these means except $t \neq 0$ arises as a unique positive invertible solution of a non-linear operator equation and satisfies all desirable properties of power arithmetic means of positive real numbers. The purpose of this talk is to extend the range in which operator power means due to Lawson–Lim–Pálfia are defined. We investigate some properties of operator power means for $t \in (-2, 2) \setminus [-1, 1]$.

- 25 遠山 宏明 (前橋工科大) Operator valued α -divergence and noncommutative ratio 15
 伊佐 浩史 (前橋工科大)
 伊藤 公智 (前橋工科大)
 亀井 栄三郎
 渡邊 雅之 (前橋工科大)
 Hiroaki Tohyama Operator valued α -divergence and noncommutative ratio
 (Maebashi Inst. of Tech.)
 Hiroshi Isa (Maebashi Inst. of Tech.)
 Masatoshi Ito (Maebashi Inst. of Tech.)
 Eizaburo Kamei
 Masayuki Watanabe
 (Maebashi Inst. of Tech.)

概要 Kamei showed some kind of the additivity for relative operator entropy $S(A|A \sharp_t B) = tS(A|B)$ for $t \in [0, 1]$. In regard to this, we show some results which have been obtained for operator valued α -divergence. Recently, we have introduced an notion of noncommutative ratio on the path $A \natural_w B$. In this talk, we also show some results obtained by applying noncommutative ratio to operator valued α -divergence.

- 26 渚 勝 (千葉大理) 作用素の対角性の特徴付け 10
 アルバニアイマム (千葉大理)
 Masaru Nagisa (Chiba Univ.) Characterization of diagonality of operators
 Albania Nugraha Imam (Chiba Univ.)

概要 We can consider Shur product of bounded linear operators on a Hilbert space. Using this product, we have already get the condition for an operator, which implies its diagonality. In this talk, we consider the algebra $B(\ell^p, \ell^q)$ ($1 \leq q \leq p \leq \infty$) of bounded linear operators which has a usual product (composition of maps) and a Schur product, and has two structures as a Banach algebra. So we can also consider the similar condition of an operators in $B(\ell^p, \ell^q)$, which implies its diagonality.

14:15~16:45

- 27 松本 健吾 (上越教育大)* 位相的マルコフシフトの軌道同型と力学系のゼータ関数 15
 松井 宏樹 (千葉大理)
 Kengo Matsumoto Continuous orbit equivalence of topological Markov shifts and dynamical zeta functions
 (Joetsu Univ. of Edu.)
 Hiroki Matui (Chiba Univ.)

概要 For continuously orbit equivalent one-sided topological Markov shifts (X_A, σ_A) and (X_B, σ_B) , we directly construct an isomorphism between their ordered cohomology groups (\bar{H}^A, \bar{H}_+^A) and (\bar{H}^B, \bar{H}_+^B) . We show that the cocycle functions for the continuous orbit equivalences give rise to positive elements of their ordered cohomology groups, so that the zeta functions of continuously orbit equivalent topological Markov shifts are related. The set of Borel measures is shown to be invariant under continuous orbit equivalence of one-sided topological Markov shifts.

- 28 永井康史 (慶大理工) タイリングにおけるパッチの分布と対応する力学系のスペクトルの性質
 15
 Yasushi Nagai (Keio Univ.) Distribution of patches in tilings and properties of spectrum of the corresponding dynamical systems

概要 Tiling is a cover of Euclidean space by tiles such as polygons that overlap only on their borders. Given a tiling, one can construct a topological dynamical system. This dynamical system is often uniquely ergodic, namely it admits a unique invariant measure. In this talk we investigate a relation between tiling and the spectrum of this measure theoretic dynamical system. It is known that we can deduce pure discrete spectrum of this dynamical system from information of distribution of patches in certain tiling. Here a patch is a configuration that appears in a tiling. In this talk we conversely deduce a property on distribution of patches from a spectral property of the corresponding dynamical system.

- 29 武石拓也 (東大数理)^b Bost–Connes system for local fields of characteristic zero 15
 Takuya Takeishi (Univ. of Tokyo) Bost–Connes system for local fields of characteristic zero

概要 The Bost–Connes system, which describes the relation between quantum statistical mechanics and class field theory, was first constructed by Bost and Connes for the rational field, and generalized for arbitrary number fields by the contribution of many researchers. In this talk, we will introduce a generalization of the Bost–Connes system for local fields of characteristic zero, and introduce some properties.

- 30 鈴木悠平 (東大数理) Amenable minimal Cantor systems of free groups arising from diagonal actions 15
 Yuhei Suzuki (Univ. of Tokyo) Amenable minimal Cantor systems of free groups arising from diagonal actions

概要 We study amenable minimal Cantor systems of free groups. We show for every free group, (explicitly given) continuum many Kirchberg algebras are realized as the crossed product of an amenable minimal Cantor system of it. In particular this shows there are continuum many Kirchberg algebras such that each of which is decomposed to the crossed products of amenable minimal Cantor systems of any virtually free group. We also give computations of K-groups for the diagonal actions of the boundary action and the odometer transformations. These computations with Matui’s theorem classify their topological full groups.

- 31 大坂博幸 (立命館大理工) The Jiang–Su absorption for inclusions of unital C*-algebras 15
 照屋保 (群馬大教育)
 Hiroyuki Osaka (Ritsumeikan Univ.) The Jiang–Su absorption for inclusions of unital C*-algebras
 Tamotsu Teruya (Gunma Univ.)

概要 In this talk we will introduce the tracial Rokhlin property for an inclusion of separable simple unital C*-algebras $P \subset A$ with finite index in the sense of Watatani, and prove theorems of the following type. Suppose that A belongs to a class of C*-algebras characterized by some structural property, such as the Jiang–Su absorption. Then P belongs to the same class. We show that an action α from a finite group G on a simple unital C*-algebra A has the tracial Rokhlin property in the sense of Phillips if and only if the canonical conditional expectation $E: A \rightarrow A^G$ has the tracial Rokhlin property for an inclusion $A^G \subset A$.

- 32 綿谷安男 (九大数理) 複素力学系からつくられる C^* -環の極大可換環と連続軌道同値 15
 梶原毅 (岡山大環境理工)
 Yasuo Watatani (Kyushu Univ.) Maximal abelian subalgebras of C^* -algebras generated by complex dy-
 Tsuyoshi Kajiwara (Okayama Univ.) namical systems and continuous orbit equivalence

概要 We consider a pair of the C^* -algebra generated by a complex dynamical system and its maximal abelian subalgebra. We study a relation between isomorphisms of such pairs and continuous orbit equivalence.

- 33 綿谷安男 (九大数理) Hilbert 空間の 3 つの部分空間の配置の Brenner 型分解 15
 榎本雅俊
 Yasuo Watatani (Kyushu Univ.) Relative position of three subspaces in a Hilbert space and Brenner type
 Masatoshi Enomoto decomposition

概要 We study the relative position of three subspaces in a infinite-dimensional Hilbert space. We give a condition that it has a Brenner type decomposition with a finite-dimensional double triangle part.

- 34 岡安類 (大阪教育大教育) Haagerup approximation property and positive cones associated with a
 戸松玲治 (北大理) von Neumann algebra 15
 Rui Okayasu (Osaka Kyoiku Univ.) Haagerup approximation property and positive cones associated with a
 Reiji Tomatsu (Hokkaido Univ.) von Neumann algebra

概要 We discuss various definitions of the Haagerup approximation property for an arbitrary von Neumann algebra. As a consequence, we give a simple and direct proof that the definition given by M. Caspers and A. Skalski is equivalent to our original one defined by using the standard form. Our strategy is to use the one-parameter family of positive cones due to H. Araki. This is based on a joint work with Reiji Tomatsu.

17:00~18:00 特別講演

佐藤康彦 (京大理)^b C^* 環の分類理論と Toms–Winter 予想

Yasuhiko Sato (Kyoto Univ.) Classification theorem of C^* -algebras and the Toms–Winter conjecture

概要 In 1989, G. A. Elliott showed that a certain class of amenable C^* -algebras can be classified by their K-groups. Following this success, he initiated the program to classify amenable C^* -algebras via K-theoretic invariants. In his program, it has become necessary to invoke some regularity property of the classifiable C^* -algebras in various manners, after the appearance of pathological examples constructed by Villadsen, Rørdam and Toms.

With the aim of characterizing classifiable C^* -algebras, in 2008, Toms and Winter have conjectured that the following three fundamental properties are equivalent for all separable, simple, amenable C^* -algebras: strict comparison, absorption of the Jiang–Su algebra, and finite nuclear dimension (or finite decomposition rank for stably finite cases). Recently, this conjecture has attracted a fair amount of attention from experts in operator algebras. Actually, it is known that an affirmative answer to this conjecture induces a solution of Rosenberg’s conjecture and also of the Blackadar–Kirchberg conjecture, which are concerned with quasidiagonality of amenable C^* -algebras.

In my talk, I report on the recent progress of the Toms–Winter conjecture and its application to the classification theorem for amenable C^* -algebras.

統計数学

9月25日(木) 第VII会場

9:30~11:50

- 1 長谷部高広 (北 大 理) 自由自己分解可能分布の単峰性 15
 S. Thorbjørnsen (Univ. of Aarhus)
 Takahiro Hasebe (Hokkaido Univ.) Unimodality of freely selfdecomposable distributions
 Steen Thorbjørnsen (Univ. of Aarhus)

概要 We show that any freely selfdecomposable probability distribution is unimodal. This is the free probabilistic analog of Yamazato's result in [Ann. Probab. 6 (1978), 523–531].

- 2 道工 勇 (埼玉大教育) ある確定的積分方程式の確率解の構成 15
 Isamu Dôku (Saitama Univ.) Construction of probabilistic solutions to a class of deterministic integral equations

概要 We consider a class of deterministic nonlinear integral equations. We begin with constructing a branching model, define a star-product and construct a tree-based star-product functional. By studying mathematical structure of the functional, we prove that the function given by expectation of the functional with respect to the law of a branching process satisfies the original integral equations.

- 3 尾張圭太 (東大経済) On the Lebesgue property of monotone convex functions on Orlicz-like spaces 15
 Keita Owari (Univ. of Tokyo) On the Lebesgue property of monotone convex functions on Orlicz-like spaces

概要 In financial mathematics, it is known as Jouini–Schachermayer–Touzi's theorem that for any convex risk measure on L^∞ with the σ -additive dual representation, (1) the Lebesgue property (order-continuity), (2) the weak compactness of sublevels of the conjugate on L^1 and the attainment of the supremum in the dual representation are equivalent. We provide a couple of generalization of this result which applies to any finite-valued monotone convex function on a lattice ideal of L^0 (including all Orlicz spaces) forming a dual pair with its order-continuous dual space with slightly different a priori lower semicontinuity assumptions and the choice of penalty function.

- 4 結城 郷 Consistency of the positive semi-definite Fourier type estimators 10
 (立命館大理工・JST CREST)
 Go Yuki Consistency of the positive semi-definite Fourier type estimators
 (Ritsumeikan Univ./JST CREST)

概要 To estimate volatilities of d -dimensional Itô semi-martingales, Malliavin and Mancino proposed Fourier series type estimators composed of the finite observations of d -dimensional Itô semi-martingale. Recently, Akahori et al. introduced Fourier series type estimators which are positive semi-definite. We discuss about the consistency of these positive semi-definite Fourier type estimators for the volatility.

- 5 小川重義 (立命館大理工) A direct inversion formula for natural SFT 10
 Shigeyoshi Ogawa (Ritsumeikan Univ.) A direct inversion formula for natural SFT

概要 We are concerned with the question whether and how a random function $f(t, \omega)$ is determined by its image of a stochastic transformation called SFT (stochastic Fourier transformation) $\mathcal{T}_*f(t, \omega)$, which was introduced by the author in 1990, in the study of stochastic integral equation of Fredholm type. The question of invertibility of the SFT has been studied again in the framework of homogeneous chaos and some affirmative answers as well as the inversion schemes were given (cf. Ogawa 2013, Ogawa–Uemura 2013, 2014). In this talk, by limiting ourselves to the case of causal functions and trigonometric basis, we are to show an elementary approach to the question, an approach that does not rely on the homogeneous chaos framework, and give a direct formula for the inversion of the natural SFT.

- 6 橋本大哉 (三和化学研)* A note on convergence rates for stability problems of SDEs under Nakao–
 土屋貴裕 Le Gall condition 15
 (会津大コンピュータ理工)
 Hiroya Hashimoto A note on convergence rates for stability problems of SDEs under Nakao–
 (Sanwa Kagaku Kenkyusho Co.) Le Gall condition
 Takahiro Tsuchiya (Univ. of Aizu)

概要 We consider the stability problems of one dimensional stochastic differential equations when the diffusion coefficients satisfy the so called Nakao–Le Gall condition. A bounded rate of strong convergence in the sense of L^1 are given by the Yamada–Watanabe method.

- 7 種村秀紀 (千葉大理) Airy random point field に対応する無限次元確率微分方程式 10
 長田博文 (九大数理)
 Hideki Tanemura (Chiba Univ.) Infinite-dimensional stochastic differential equations arising from Airy
 Hirofumi Osada (Kyushu Univ.) random point fields

概要 We identify infinite-dimensional stochastic differential equations (ISDEs) describing the stochastic dynamics related to Airy random point fields with $\beta = 1, 2, 4$. We prove the existence and uniqueness of the ISDEs.

- 8 種村秀紀 (千葉大理) 末尾 σ -加法族と無限次元確率微分方程式の強解の存在と一意性 10
 長田博文 (九大数理)
 Hideki Tanemura (Chiba Univ.) Strong solutions of infinite-dimensional stochastic differential equations
 Hirofumi Osada (Kyushu Univ.) and tail σ -fields

概要 We present a new method to construct unique strong solutions of infinite-dimensional stochastic differential equations (ISDEs) describing interacting Brownian motions (IBMs). Our method can be applied to IBMs related to random matrix theory.

- 9 天羽隆史 (立命館大理工) An integration by parts on space of loops 10
 Takafumi Amaba (Ritsumeikan Univ.) An integration by parts on space of loops

概要 We consider to construct a measure on space of loops in \mathbb{C}^* (strictly speaking, a space of paths in a coefficient body), surrounding the origin by employing the utility of the (alternate) Loewner–Kufarev equation. We discuss about a simple integration by parts formula under the measure.

- 10 矢野孝次 (京大理) 一次元拡散過程に対する修正零レゾルベント 15
 矢野裕子 (京都産大理)
 Kouji Yano (Kyoto Univ.) Renormalized zero resolvents for one-dimensional diffusions
 Yuko Yano (Kyoto Sangyo Univ.)

概要 For a one-dimensional diffusion on an interval for which 0 is the regular reflecting left boundary, the renormalized zero-resolvent, denoted by h_0 , is studied. It is reported that, for the process stopped upon hitting 0, the function h_0 is excessive and the h_0 -transform can be regarded as a conditional process to avoid zero.

14:15~15:15 特別講演

- 竹内敦司 (阪市大理) ジャンプ型確率過程に対する部分積分公式
 Atsushi Takeuchi (Osaka City Univ.) Integration by parts formula for jump processes

概要 It is well known that the Malliavin calculus is a powerful tool in stochastic analysis, and the integration by parts formula plays a crucial role in the argument. Most of works concern only diffusion processes without any jumps. Although it is a natural question whether a similar approaches can be taken in case of jump processes, there are a lot of difficulties to do it. Moreover, there are some approaches to the Malliavin calculus for jump processes, depending on the interests. In this talk, we shall give a brief review of some recent results concerning the integration by parts formula for jump processes.

15:30~16:30 特別講演

- 高橋博樹 (慶大理工)* エノン写像におけるスメールの馬蹄の崩壊と、その後におこる現象について
 Hiroki Takahashi (Keio Univ.) On the destruction of Smale’s horseshoe in the Henon map (and what comes afterwards)

概要 An important problem in dynamics (qualitative theory of ordinary differential equations) is to describe how Smale’s horseshoe loses its stability through continuous modifications of the system. The loss of stability of Smale’s horseshoe through homoclinic bifurcations is modeled by a certain parametrized family of planar diffeomorphisms, called the Henon maps. I will talk about recent results on the dynamics of the strongly dissipative Henon maps around the first bifurcation parameter, with special emphasis on their geometric and probabilistic aspects.

9月26日(金) 第VII会場

9:30~11:30

- 11 小林正弘 (東京理大理工) 2次元反射型ランダムウォークにおける構造的可逆性 15
清水宏 (日本ユニシス)
宮沢政清 (東京理大理工)
Masahiro Kobayashi (Tokyo Univ. of Sci.) Structure-reversibility of a two dimensional reflecting random walk
Hiroshi Shimizu (Nihon Unisys)
Masakiyo Miyazawa (Tokyo Univ. of Sci.)

概要 We consider a two dimensional reflecting random walk on the nonnegative integer quadrant. It is assumed that this reflecting random walk has skip free transitions. We are concerned with its time reversed process assuming that the stationary distribution exists. In general, the time reversed process may not be a reflecting random walk. In this paper, we derive necessary and sufficient conditions for the time reversed process also to be a reflecting random walk. These conditions are different from but closely related to the product form of the stationary distribution.

- 12 岡田いず海 (東工大理工) ランダムウォークの訪問点集合の境界点について 15
Izumi Okada (Tokyo Tech) The inner boundary of random walk range

概要 We deal with the inner boundary of random walk range, that is, the set of those points in a random walk range which have at least one neighbor site outside the range. We consider the number of the inner boundary points of random walk range and the question; how many times does a simple random walk revisit the most frequently visited site among the inner boundary points?

- 13 梶野直孝 (神戸大理) 点再帰的強局所対称 Dirichlet 空間における内部一様領域上の Neumann
J. Lierl (Univ. Illinois UC) 熱核評価 15
Naotaka Kajino (Kobe Univ.) Neumann heat kernel estimates on inner uniform domains in point-
Janna Lierl (Univ. Illinois UC) recurrent strongly local symmetric Dirichlet spaces

概要 For a strongly local regular Dirichlet space arising from a resistance form and with a complete geodesic metric and a *sub-Gaussian* type two-sided heat kernel estimate, we present a two-sided estimate of the Neumann heat kernel on inner uniform subdomains in terms of the natural inner geodesic metric of the domains. Besides the change of the metric, the Neumann heat kernel estimate involves no difference from the global heat kernel estimate. This result is meant mainly for inner uniform domains of fractals with spectral dimension less than 2 and is applicable to Brownian motion on affined nested fractals and 2-dimensional generalized Sierpiński carpets.

- 14 C. Aistleitner (Graz Univ. of Tech.)* 有界間隙列の重複対数の法則について II 5
 福山克司 (神戸大理)
 Christoph Aistleitner The law of the iterated logarithm for lacunary series with bounded gaps
 (Graz Univ. of Tech.) II
 Katusi Fukuyama (Kobe Univ.)

概要 For every positive number σ , there exists a sequence $\{n_k\}$ of integers satisfying $n_{k+1} - n_k \in \{0, 1\}$ such that the law of the iterated logarithm for $\sum \cos(2\pi n_k x)$ with limsup constant σ holds.

- 15 竹居正登 (横浜国大工) 三角格子上の Ising percolation における横断確率について 10
 Masato Takei (Yokohama Nat. Univ.) On crossing probabilities for Ising percolation on the triangular lattice

概要 We consider the percolation problem for spin configurations of the Ising model on the triangular lattice. We derive an RSW-type bound for crossing probabilities when the external field is zero and the inverse temperature is not more than the critical one. As an application, we can obtain power estimates of the one-arm probability and the two-point connectivity function from below, at the critical inverse temperature.

- 16 楠岡誠一郎 (東北大理) 半自己相似性を持つランダム媒質中の多次元拡散過程の再帰性と非再帰性
 高橋弘 (日大理工) 15
 田村要造 (慶大理工)
 Seiichiro Kusuoka (Tohoku Univ.) Recurrence and transience properties of multi-dimensional diffusion processes in semi-selfsimilar random environments
 Hiroshi Takahashi (Nihon Univ.)
 Yozo Tamura (Keio Univ.)

概要 We consider limiting behaviors of multi-dimensional diffusion processes in multi-parameter random environments, which are sets of values at different d points of one-dimensional α -stable or (r, α) -semi-stable Lévy processes. They imply conditions of random environments for the dichotomy of recurrence and transience of multi-dimensional diffusion processes formed by d independent one-dimensional diffusion processes in α -stable or (r, α) -semi-stable Lévy processes. Their limiting behaviors are quite different from those of multi-dimensional standard Brownian motions. We also consider a direct product of a one-dimensional diffusion process in a reflected non-positive Brownian environment and a one-dimensional standard Brownian motion. For the two-dimensional diffusion process, we show the transience property.

- 17 中島 誠 (筑波大数理物質) 1+2 次元ランダム媒質中のディレクティドポリマーの自由エネルギーの高温度での評価 15
 Makoto Nakashima (Univ. of Tsukuba) On the estimates of the free energy of directed polymers in random environment in 1+2 dimension at high temperature

概要 We consider the free energy of the directed polymers in random environment in 1+2 dimension. It is known that the free energy is strictly negative if the inverse temperature β is not zero when spatial dimension d is 1 or 2. In this talk, we give new bounds of the free energy as $\beta > 0$ small.

- 18 笹井健行 (東大情報理工) EFKP 型の重複対数法則のゲーム論的証明 15
 宮部賢志 (明大理工)
 竹村彰通 (東大情報理工)
 Takeyuki Sasai (Univ. of Tokyo) A game-theoretic proof of Erdős–Feller–Kolmogorov–Petrowsky law of
 Kenshi Miyabe (Meiji Univ.) the iterated logarithm for fair coin tossing
 Akimichi Takemura (Univ. of Tokyo)

概要 We give a game-theoretic proof of the celebrated Erdős–Feller–Kolmogorov–Petrowsky law of the iterated logarithm for fair coin tossing. As many other game-theoretic proofs of the laws in probability theory, our game-theoretic proof is explicit and gives insights on what happens when the law is violated.

11:45～12:15 統計数学科会総会

9月27日(土) 第VII会場

9:30～12:00

- 19 鈴木 聡 (島根大総合理工) Surrogate 双対性と制約想定について 15
 黒岩大史 (島根大総合理工)
 Satoshi Suzuki (Shimane Univ.) Surrogate duality and its constraint qualifications
 Daishi Kuroiwa (Shimane Univ.)

概要 In this talk, we study a constraint qualification which completely characterizes surrogate strong and min-max duality for quasiconvex programming. We show that the closed cone constraint qualification for surrogate duality is a necessary and sufficient constraint qualification for surrogate strong and min-max duality via quasiconvex programming. Also, we compare our constraint qualification with previous ones for Lagrange and surrogate duality.

- 20 藤田敏治 (九工大工) 結合型評価をもつ相互依存型マルコフ決定過程 10
 Toshiharu Fujita Mutually dependent Markov decision processes with associative criteria
 (Kyushu Inst. of Tech.)

概要 In this study, we consider in mutually dependent Markov decision processes (MDMDP) with associative criteria. The MDMDP model is structured upon finite-stage Markov decision processes. At each stage, the reward in one decision process is given by the optimal values of other decision process, whose initial state is determined by the current state and decision in the original process. We introduce an associative criterion to each MDMDP and derive recursive equations by dynamic programming with an invariant imbedding technique.

- 21 須藤慶大 (早大理工) 補間誤差をコントラスト関数とする母数推定 15
 劉言 (早大理工)
 谷口正信 (早大理工)
 Yoshihiro Suto (Waseda Univ.) Parameter estimation by a contrast function based on interpolation error
 Yan Liu (Waseda Univ.)
 Masanobu Taniguchi (Waseda Univ.)

概要 Interpolation is an important issue for a variety of fields in statistics (e.g., missing data analysis). In this research, the asymptotics of a contrast function estimator defined by pseudo interpolation error for Gaussian stationary process are investigated. We estimate parameters of the process by minimizing the pseudo interpolation error written in terms of a fitted parametric spectral density and the periodogram based on observed stretch. The estimator has the consistency and asymptotic normality. Although the criterion for the interpolation problem is known as the best in the sense of smallest mean square error for past and future extrapolation, it is shown that the estimator is asymptotically inefficient in general, which leads to an unexpected result.

- 22 明石郁哉 (早大理工) 周波数領域における GMM 推定量の高次漸近有効性 15
 Fumiya Akashi (Waseda Univ.) On the second-order asymptotic efficiency of frequency domain GMM estimators

概要 We elucidate the second-order asymptotic properties of frequency domain generalized method of moments (GMM) estimators for second order stationary processes. The process concerned is essentially nonparametric one, and we focus on the estimation problem of the pivotal quantity of the process, which is defined as a solution of over-identified spectral restrictions. The Edgeworth expansion for the GMM estimator up to second-order is given. We also discuss asymptotic efficiency of the GMM estimator by making a comparison between the limit distributions of the GMM estimator and a quasi-maximum likelihood estimator.

- 23 劉言 (早大理工) Quantile estimation in frequency domain 15
 Yan Liu (Waseda Univ.) Quantile estimation in frequency domain

概要 Nowadays, the quantile estimation becomes a notable method in statistics for its robustness against the existence of moments of random variables. In this talk, we extend the idea of quantile in time domain to that in frequency domain. The objective function for the quantile estimator in time domain can be naturally extended into frequency domain. The quantile estimator in frequency domain has the consistency. However, asymptotic normality of the quantile estimator based on the bare periodogram does not hold, which is obviously different from the quantile theory for time domain. We give the asymptotic properties of the estimator. The modified estimator for asymptotic normality will also be provided.

- 24 佃 康 司 (総合研究大学院大) L^2 空間アプローチによるエルゴード的確率過程のパラメータ経時的不均
 西 山 陽 一 性の検定 15
 (統計数理研・総合研究大学院大)

Koji Tsukuda Testing the time-homogeneity of ergodic stochastic processes by an L^2
 (Grad. Univ. for Adv. Stud.) space approach

Yoichi Nishiyama
 (Inst. of Stat. Math./Grad. Univ. for Adv. Stud.)

概要 A general approach to test the time-homogeneity of ergodic stochastic process models is presented. It is based on weak convergences of stochastic integrals taking values in L^2 spaces, and it enables us to use the Anderson–Darling type weight function which cannot be treated by a weak convergence theory for martingales taking values in ℓ^∞ spaces. The result is applied to some time-homogeneity testing problems of ergodic diffusion processes and stress release models with continuous observations.

- 25 柿 沢 佳 秀 (北 大 経 済) Bootstrap-based Bartlett-type adjustment 15
 Yoshihide Kakizawa (Hokkaido Univ.) Bootstrap-based Bartlett-type adjustment

概要 For the LR statistic, the Bartlett adjustment is applicable, i.e., a simple mean adjustment through multiplication by a constant of the form $1+b/N$ yields an improvement of the null distribution. On the other hand, the Bartlett-type adjustment is a higher-order asymptotic method for improving the chi-squared approximation to the null distributions of various statistics rather than the LR statistic. In this talk, we propose a bootstrap-based Bartlett-type adjustment, which is an extension of a bootstrap-based Bartlett adjustment for the LR statistic.

- 26 前 園 宜 彦 (九 大 数 理) ハザード関数のカーネル型推定量の高次漸近表現 10
 Yoshihiko Maesono (Kyushu Univ.) Higher order asymptotic representation of kernel estimator of hazard
 function

概要 In this talk we obtain an asymptotic representation a kernel type estimator of a hazard function. The hazard function estimator takes a form of a ratio of two kernel type estimators. The convergence rates of them are different, and so we cannot see an effect of the slow order rate estimator under an ordinal asymptotic setting. Using the higher order asymptotic representation, we can discuss an effect of the slow order rate estimator.

- 27 柳原宏和 (広島大理) On asymptotically KL loss efficiency of a log-likelihood-based information criterion in high-dimensional normal multivariate linear regression models 15
 Hirokazu Yanagihara (Hiroshima Univ.) On asymptotically KL loss efficiency of a log-likelihood-based information criterion in high-dimensional normal multivariate linear regression models

概要 This paper deals with a variable selection procedure in a multivariate linear regression model with normality assumption, which is called a normal multivariate linear regression model, by minimizing a model selection criterion. A model selection criterion considered in this paper is a log-likelihood-based information criterion which is defined by adding a penalty term to the negative twofold maximum log-likelihood. A purpose of this paper is to clarify a sufficient condition of the penalty term in the log-likelihood-based information criterion to satisfy an asymptotically Kullback–Leibler (KL) loss efficiency property from the HD asymptotic framework, such that $(n, p) \rightarrow \infty$ simultaneously under the condition $p/n \rightarrow c_0 \in (0, 1)$. Then, we can study whether AIC, AIC_c, BIC, CAIC and HQC are asymptotically KL loss efficient or not, when the HD asymptotic framework is used.

- 28 綾野孝則 (阪大理) 連続データに対する MDL 規準の漸近的な性質 10
 鈴木 讓 (阪大理)
 Takanori Ayano (Osaka Univ.) Asymptotic property of MDL information criterion for continuous data
 Joe Suzuki (Osaka Univ.)

概要 It is very important to estimate the probability of the data series accurately for applying MDL information criterion. For discrete data, it is known that the methods based on universal coding in information theory have high precision (universal Bayesian measures) and they are used in MDL. Recently, Ryabko extended the universal Bayesian measures for discrete data to continuous data and one gets to be able to apply MDL for not only discrete data but also continuous data. In this presentation, we give the rates of convergence for the generalization error of the universal Bayesian measures for continuous data and show that they achieve the optimal rate under a certain condition.

14:15～15:15 特別講演

- 藤澤洋徳 (統計数理研) ダイバージェンスに基づいたロバスト統計
 Hironori Fujisawa (Inst. of Stat. Math.) Divergence-based robust statistics

概要 Many estimation methods have been proposed for robust statistics. In this talk, some divergence-based methods are discussed. The density power divergence was proposed by Basu et al. (1998). This is based on the idea that the density values for outliers are small. The largest characteristic is that the density power divergence is easily applicable for any continuous probability density function. The density power divergence belongs to a separable class of Bregman divergences. Fujisawa and Eguchi (2008) proposed another type of divergence, which was called the gamma-divergence. This divergence belongs to a larger class of Bregman divergence. The robust estimator based on the gamma divergence can show a sufficiently small latent bias even if the ratio of outlier is not small. This solves an important open problem that has not been unsolved for a long time. Kanamori and Fujisawa (2014) proposed an extended class of divergence from the point of view of affine invariance of data transformation, which was called the Hölder divergence. This divergence does not belong to a class of Bregman divergence. Recently, Kanamori and Fujisawa (2014) pointed out a close relation between the gamma and Hölder divergences by virtue of unnormalized model.

15:30~16:30 特別講演

宮田 庸一 (高崎経大経済) ベイズ型推定量の漸近的な性質について

Yoichi Miyata On asymptotic properties of Bayesian type estimators
(Takasaki City Univ. of Econ.)

概要 Roughly speaking, nonstationary processes can be classified into three main groups: (i) nonstationary mixing processes, (ii) processes with deterministic trends, and (iii) unit root processes. In this talk, we will consider general conditions and rather easily verified conditions that ensure asymptotic posterior normality, strong consistency of the Bayesian type estimators and their asymptotic normality in possibly misspecified models for stationary processes, (i), and (ii). We will illustrate our main results with a heterogeneous AR model, a Logit model, and a finite mixture model.

9月28日(日) 第VII会場

9:30~11:20

29 田中 秀和 (阪府大工) Some results on gamma parameter estimation 15

N. Pal
(Univ. Louisiana at Lafayette)Wooi K. Lim
(William Paterson Univ.)

Hidekazu Tanaka (Osaka Pref. Univ.) Some results on gamma parameter estimation

Nabendu Pal
(Univ. Louisiana at Lafayette)

Wooi K. Lim (William Paterson Univ.)

概要 This talk deals with improved estimation of gamma parameters from a decision-theoretic point of view. First, we study the second order properties of three estimators of shape parameter —(i) the maximum likelihood estimator (MLE), (ii) a bias corrected version of the MLE, and (iii) an improved version (in terms of mean squared error) of the MLE. It is shown that all the three estimators mentioned above are second order inadmissible. Also, we obtain second order admissible estimators which are second order better than the above three estimators. Similarly, in estimating scale parameter, we consider the second order admissibilities of some estimators and propose second order admissible estimators.

30 生 亀 清 貴 (東京理大理工) 多変量密度関数の点対称性と分解 10

富 澤 貞 男 (東京理大理工)

Kiyotaka Iki (Tokyo Univ. of Sci.) Point-symmetric multivariate density function and decomposition

Sadao Tomizawa (Tokyo Univ. of Sci.)

概要 For a T -variate density function, we define the point-symmetry, quasi-point-symmetry of order k ($< T$) and the marginal point-symmetry of order k , and gives the theorem that the density function is T -variate point-symmetric if and only if it is quasi-point-symmetric and marginal point-symmetric of order k . The theorem is illustrated for the multivariate normal density function.

- 31 猪口 真 (広島大理) 高次元 GMANOVA モデルにおける係数行列の LSE と MLE の漸近比較
 柳原 宏和 (広島大理) 15
 Makoto Inokuchi (Hiroshima Univ.) Asymptotical comparison with LSE and MLE of coefficient matrices in
 Hirokazu Yanagihara (Hiroshima Univ.) high-dimensional GMANOVA model

概要 In this paper, we asymptotically compare with the least square estimator (LSE) and the maximum likelihood estimator (MLE) of coefficient matrices in high-dimensional GMANOVA model, when the sample size n and the number of response variables p simultaneously approach ∞ under the condition that $c_{n,p} = p/n \rightarrow c_0 \in [0, 1)$.

- 32 石井 晶 (筑波大数理物質) 高次元小標本における共分散行列の同等性検定 15
 矢田 和善 (筑波大数理物質)
 青嶋 誠 (筑波大数理物質)
 Aki Ishii (Univ. of Tsukuba) Equality test of covariance matrices in high-dimension, low-sample-size
 Kazuyoshi Yata (Univ. of Tsukuba) context
 Makoto Aoshima (Univ. of Tsukuba)

概要 A common feature of high-dimensional data is the data dimension is high, however, the sample size is relatively low. We call such data HDLSS data. Ishii et al. (2014) gave an asymptotic distribution of the largest eigenvalue estimator derived by the noise-reduction methodology that was created by Yata and Aoshima (2012). In this talk, we provide an estimator of the first eigenvector by using the noise-reduction methodology. We show that the estimator enjoys the consistency property in a mild condition. We consider testing the equality of covariance matrices between two classes and propose a new test statistic by applying both the largest eigenvalue estimator and the first eigenvector estimator. We verify the proposed test statistic improves the power as the dimension grows.

- 33 八木 文香 (東京理大理) k -step 単調欠測データの下での平均ベクトルの検定 15
 瀬尾 隆 (東京理大理)
 Ayaka Yagi (Tokyo Univ. of Sci.) A test for the mean vector with k -step monotone missing data
 Takashi Seo (Tokyo Univ. of Sci.)

概要 We consider the problem of testing the mean vector in one-sample problem when the data have k -step monotone pattern missing observations. Jinadasa and Tracy (1992) obtained closed form expressions for the maximum likelihood estimators of the mean vector and the covariance matrix of a multivariate normal distribution with a k -step monotone missing data pattern. We propose a simplified Hotelling's T^2 type statistic by evaluating the covariance matrix of the maximum likelihood estimator of the mean vector in the case of a k -step monotone missing data. Further, we give an approximate upper percentile of the T^2 type statistic and investigate the accuracy by Monte Carlo simulation.

- 34 矢田 和善 (筑波大数理物質) 高次元混合分布における PCA とその応用 15
 青嶋 誠 (筑波大数理物質)
 Kazuyoshi Yata (Univ. of Tsukuba) High-dimensional PCA for a mixture model and its applications
 Makoto Aoshima (Univ. of Tsukuba)

概要 In this talk, we consider PCA (Principle Component Analysis) for mixture models in high-dimension, low-sample-size (HDLSS) settings. We show that the first true eigenvalue, eigenvector and its PC scores hold consistency properties when the underlying distribution has a two-class mixture model. Next, we derive a geometric representation of the dual sample covariance matrix constructed by HDLSS data from the mixture model. With the help of the geometric representation, we propose a new clustering method for a high-dimensional mixture distribution. Finally, we extend the clustering method to a general case when three or more classes are mixed.

- 35 長縄 真学 (東京理大理工) 順序カテゴリ正方分割表における拡張線形非対称モデル 10
 田畑 耕治 (東京理大理工)
 富澤 貞男 (東京理大理工)
 Masato Naganawa (Tokyo Univ. of Sci.) Extended linear asymmetry model for square contingency tables with
 Kouji Tahata (Tokyo Univ. of Sci.) ordered categories
 Sadao Tomizawa (Tokyo Univ. of Sci.)

概要 The issues of various symmetry rather than independence arise naturally for the analysis of square contingency tables. So the models, that indicate the structure of symmetry and asymmetry, have been proposed, for example, the symmetry model (Bowker, 1948), the quasi-symmetry model (Causinus, 1965), and the marginal homogeneity model (Stuart, 1955). In this talk, we propose the models that indicate the structure of asymmetry. The model includes various symmetry and asymmetry models in the special cases. An example is given.

14:15~16:10

- 36 澤 正憲 (神戸大システム情報) 測度空間上のデザイン理論の構築に向けて, I 15
 Masanori Sawa (Kobe Univ.) Developing the theory of designs on measure spaces, I

概要 In this talk, I introduce the notion of cubature formulas or designs on measure spaces, generalizing various related objects in statistical experiments, numerical analysis, and algebraic combinatorics. Many examples will be given emphasizing how general our notion is. The idea of cubature on measure spaces originally goes back to an outstanding work by Kôno in 1962 on the construction of optimal experimental designs on the cube.

- 37 平尾 将剛 (愛知県立大情報) Characterizing optimum designs in terms of finite irreducible reflection
 澤 正憲 (神戸大システム情報) groups, II 10
 Masatake Hirao (Aichi Pref. Univ.) Characterizing optimum designs in terms of finite irreducible reflection
 Masanori Sawa (Kobe Univ.) groups, II

概要 In this talk we give a geometric characterization of D-optimal experimental designs on the unit ball that consist of corner vectors associated with finite irreducible reflection groups A_n and D_n .

- 38 山田 紘 頌 (名大情報) グラフの距離行列の符号数に関する Graham–Lovász の問題について .. 10
 Kohei Yamada (Nagoya Univ.) Some results related to a remark of Graham and Lovász

概要 Let G be an undirected connected graph without loops and multiedges, and let $n_+(G)$ and $n_-(G)$ be the number of positive and negative eigenvalues, respectively, of the distance matrix of G . Graham and Lovász [Adv. in Math., Vol.78, 1978] posed a problem of whether there exists a graph with $n_-(G) < n_+(G)$. Recently, Azarija [Discrete Math., 315-316, pp.65–68, 2014] proved that the Paley graph of order at least 13 is such an example. In this talk, I present some observations on graphs with $n_+(G) = n_-(G)$.

- 39 地 寄 頌 子 (東京理大理工) Difference systems of sets with size 2 10
 宮 本 暢 子 (東京理大理工)
 Shoko Chisaki (Tokyo Univ. of Sci.) Difference systems of sets with size 2
 Nobuko Miyamoto (Tokyo Univ. of Sci.)

概要 Difference systems of sets (DSS) are combinatorial structures introduced by Levenshtein in 1971, which are a generalization of cyclic difference sets and arise in connection with code synchronization. A DSS is a collection of t disjoint subsets $Q_i, 0 \leq i \leq t-1$, of any finite abelian group G of order $q = ef + 1$ such that every element of $G \setminus \{0\}$ appears at least ρ times in the multiset $\{a - b \mid a \in Q_i, b \in Q_j, 0 \leq i \neq j \leq t-1\}$. In this talk, we will talk about parameter ρ of DSS with $e = 4, 6$. Additionally, we define new blocks with size 2 and present the conditions that these blocks form a DSS.

- 40 盧 暁 南 (名大情報) Affine-invariant strictly cyclic Steiner quadruple systems and related
 神 保 雅 一 (名大情報) hypergraphs 15
 Xiao-Nan Lu (Nagoya Univ.) Affine-invariant strictly cyclic Steiner quadruple systems and related
 Masakazu Jimbo (Nagoya Univ.) hypergraphs

概要 A Steiner quadruple system denoted by $SQS(v)$, is a pair (V, \mathcal{B}) , where V is a finite set of v elements, and \mathcal{B} is a set of 4-subsets of V , called blocks or quadruples, such that each 3-subset of V appears exactly once in \mathcal{B} . An $SQS(v)$ admitting a cyclic permutation whose stabilizer of any block is trivial is said to be strictly cyclic, denoted by $sSQS(v)$. In this talk, we consider an $sSQS(2p)$ over \mathbb{Z}_{2p} admitting all the units of \mathbb{Z}_{2p} as multipliers, which is said to be affine-invariant. We show that the blocks of an affine-invariant $sSQS(2p)$ can be obtained from a rainbow 1-factor of a special kind of hypergraphs.

- 41 兵頭 義史 (岡山理大総合情報研・国際自然研) Existence conditions for balanced fractional 2^m factorial designs of resolution $R^*({1}|\Omega_\ell)$ with $N < \nu_\ell(m)$ 15
 栗田 正秀 (国際自然研)
 弓場 弘 (国際自然研)
 Yoshifumi Hyodo (Okayama Univ. of Sci./Int. Inst. for Nat. Sci.) Existence conditions for balanced fractional 2^m factorial designs of resolution $R^*({1}|\Omega_\ell)$ with $N < \nu_\ell(m)$
 Masahide Kuwada (Int. Inst. for Nat. Sci.)
 Hiromu Yumiba (Int. Inst. for Nat. Sci.)

概要 We consider a fractional 2^m factorial design derived from a simple array (SA) such that the $(\ell + 1)$ -factor and higher-order interactions are assumed to be negligible, where $\ell = 2, 3$. Under these situations, if the main effect is estimable, and furthermore some of the remaining non-negligible factorial effects may or may not be estimable, then a design is said to be of resolution $R^*({1}|\Omega_\ell)$. Using the algebraic structure of the TMDPB association scheme, we give a necessary and sufficient condition for an SA to be a balanced fractional 2^m factorial design of resolution $R^*({1}|\Omega_\ell)$, where the number of assemblies is less than the number of non-negligible factorial effects.

- 42 松原 和樹 (広島大理) An asymptotic existence of pairwise additive minimal BIB designs ... 15
 景山 三平 (広島工大環境)
 Kazuki Matsubara (Hiroshima Univ.) An asymptotic existence of pairwise additive minimal BIB designs
 Sanpei Kageyama (Hiroshima Inst. of Tech.)

概要 The existence of pairwise additive balanced incomplete block (BIB) designs, denoted by $PAB(v, k, \lambda)$, has been discussed with direct and recursive constructions in Matsubara et al. (2007, 2013). Especially, when k is an odd integer, $PAB(v, k, (k - 1)/2)$ are said to be *minimal*. On the other hand, Wilson (1975) proved an asymptotic existence of pairwise balanced designs. In this talk, an asymptotic existence of pairwise additive minimal BIB designs is shown by use of Wilson's theorem.

- 43 景山 三平 (広島工大環境)* An affine α -resolvable triangular design is not of simple type 10
 Sanpei Kageyama (Hiroshima Inst. of Tech.) An affine α -resolvable triangular design is not of simple type

概要 The validity of affine α -resolvability in 2-associate PBIB designs is considered. Some history is reviewed. Finally we will show that there does not exist an affine α -resolvable triangular design with $\lambda_1 = 0$ or $\lambda_2 = 0$.

応 用 数 学

9月25日(木) 第VIII会場

9:30~11:30

- 1 野 崎 寛 (愛知教育大教育) 正則グラフに対する線形計画限界について 15
 Hiroshi Nozaki (Aichi Univ. of Edu.) Linear programming bounds for regular graphs

概要 We develop a linear programming method to obtain bounds for the number of vertices of a connected regular graph. This method especially need only the information of the distinct eigenvalues of a graph. As an application of this bound, we prove that a connected k -regular graph satisfying $g > 2d - 1$ has the minimum second largest eigenvalue of all k -regular graphs of the same size, where d is the number of distinct non-trivial eigenvalues, and g is the girth.

- 2 足立俊明 (名 工 大)* ケーラーグラフ 10
ヤリママトルソンママト
 (名 工 大)

Toshiaki Adachi (Nagoya Inst. of Tech.) Kähler graphs
Tuerxunmaiti Yaermaiti
 (Nagoya Inst. of Tech.)

概要 As a discrete model of a Kähler manifold admitting Kähler magnetic fields, we introduce the notion of Kähler graphs. We show some examples of Kähler graphs and give a condition that we can construct a vertex-transitive finite Kähler graphs.

- 3 足立俊明 (名 工 大)* ケーラーグラフの (1,1)-ラプラシアン 10
ヤリママトルソンママト
 (名 工 大)

Toshiaki Adachi (Nagoya Inst. of Tech.) (1,1)-Laplacians for Kähler graphs
Tuerxunmaiti Yaermaiti
 (Nagoya Inst. of Tech.)

概要 For Kähler graphs we define their (p, q) -Laplacians by use of (p, q) -colored paths. We particularly consider (1, 1)-Laplacians and study eigenvalues for Kähler graphs defined by complement graphs and those of product types. We give examples of pairs of (1, 1)-isospectral Kähler graphs.

- 4 佐藤 巖 (小山工高専) グラフの離散時間量子ウォークの遷移行列に関するゼータ関数 …… 15
今野 紀雄 (横浜国大工)
樋口 雄介 (昭和大教養)
瀬川 悦生 (東北大情報)
Iwao Sato (Oyama Nat. Coll. of Tech.) A zeta function with respect to the transition matrix of a discrete-time
Norio Konno (Yokohama Nat. Univ.) quantum walk on a graph
Yusuke Higuchi (Showa Univ.)
Etsuo Segawa (Tohoku Univ.)

概要 From the viewpoint of quantum walks, the Ihara zeta function of a finite graph can be said to be closely related to its evolution matrix. In this note we introduce another kind of zeta function of a graph, which is closely related to, as to say, the square of the evolution matrix of a quantum walk. Then we give to such a function two types of determinant expressions and derive from it some geometric properties of a finite graph. As an application, we illustrate the distribution of poles of this function comparing with those of the usual Ihara zeta function.

- 5 鈴木 有祐 (新潟大理) 四角形分割から諸条件を満たす三角形分割への拡張 …… 15
野口 健太 (慶大理工)
Yuusuke Suzuki (Niigata Univ.) Extension to triangulations with some properties from quadrangulations
Kenta Noguchi (Keio Univ.)

概要 For a given triangulation G on a surface F^2 , we can extend G to a triangulation T by adding a diagonal edge in every face of G . Now the problem is to find a triangulation T with some properties. In this talk, we show that for a given simple 2-connected planar quadrangulation G with $|V(G)| \geq 6$, we can extend G to a 4-connected triangulation.

- 6 浜野 銀次 (東京電機大理工) 有限グラフに付随する辺凸多面体の正則単模三角形分割の存在 …… 15
日比 孝之 (阪大情報)
大杉 英史 (関西学院大理工)
Ginji Hamano (Tokyo Denki Univ.) Existence of a regular unimodular triangulation of the edge polytopes
Takayuki Hibi (Osaka Univ.) of finite graphs
Hidefumi Ohsugi
 (Kwansei Gakuin Univ.)

概要 Let G be a fundamental FHM-graph and P_G the edge polytope of G . Ohsugi has obtained a necessary and sufficient condition for P_G to possess a regular unimodular triangulation. However, this condition is not so easy to apply to a given graph only by looking at the graph.

In this lecture, for a fundamental FHM-graph G , we will give several criteria for the existence of a regular unimodular triangulation of P_G in terms of some simple data of the graph. We also apply our criteria to some examples, including the complete graph K_6 with 6 vertices, and show that their edge polytopes possess a regular unimodular triangulation.

- 7 須山雄介(阪市大理) 単体的2球面の頂点の番号づけ 10
 Yusuke Suyama (Osaka City Univ.) Numbering of vertices of simplicial 2-spheres

概要 Let K be a triangulation of the 2-sphere. We assign numbers to vertices of K on the basis of a certain rule. We determine the number of triangulations with at most 9 vertices, such that all vertices can be numbered by the rule.

- 8 小関健太 3-正則グラフの分割問題 15
 (国立情報学研・JST ERATO)

Dong Ye

(Middle Tennessee State Univ.)

Kenta Ozeki

A decomposition of cubic graphs

(Nat. Inst. of Information/JST ERATO)

Dong Ye (Middle Tennessee State Univ.)

概要 A decomposition of a graph G is a set E_1, \dots, E_k of disjoint edges such that the union of them form the edge set $E(G)$ of G . It was conjectured by Hoffmann–Ostenhof that every cubic graph has a decomposition E_1, E_2, E_3 such that E_1 is a spanning tree, E_2 is a matching, and E_3 is a family of cycles. (Note that a matching and a family of cycles desired here are not necessarily spanning subgraphs.) We prove that the conjecture is true for 3-connected plane cubic graphs.

14:15~17:00

- 9 潮和彦(近畿大理工) Balanced C_5 -foil designs and related designs 15
 Kazuhiko Ushio (Kinki Univ.) Balanced C_5 -foil designs and related designs

概要 In graph theory, the decomposition problem of graphs is a very important topic. Various type of decomposition of many graphs can be seen in the literature of graph theory. This paper gives balanced C_5 -foil designs and related designs.

- 10 藤沢潤(慶大商) 射影平面上のグラフのマッチング拡張問題について 15
 瀬野博紀(横浜国大環境情報)

Jun Fujisawa (Keio Univ.) Matching extension in projective planar graphs

Hiroki Seno (Yokohama Nat. Univ.)

概要 A matching M of G is said to be extendable in G if M is a subset of a perfect matching of G , and a graph with at least $2m + 2$ vertices in which every matching of size m is extendable is called m -extendable. Moreover, a graph with at least $2m + 2$ vertices is said to be distance d m -extendable if any matching M with $|M| = m$ in which the edges lie pair-wise distance at least d is extendable. In this talk we introduce the following result: Every 5-connected triangulation of the projective plane with an even order is distance 3 7-extendable, and there exist infinitely many 5-connected triangulations of the projective plane with an even order which are not distance 3 8-extendable.

- 11 善本 潔 (日大理工) Locating sets of vertices on Hamiltonian cycles 15
 R. J. Faudree (Univ. of Memphis)
 Hao Li (Univ. de Paris Sud)
 Kiyoshi Yoshimoto (Nihon Univ.) Locating sets of vertices on Hamiltonian cycles
 Ralph J. Faudree (Univ. of Memphis)
 Hao Li (Univ. de Paris Sud)

概要 Let $k \geq 2$ be a positive integer and $X = \{x_1, \dots, x_k\}$ and $Y = \{y_1, \dots, y_k\}$ be a pair of disjoint sets of vertices in a graph G . In this talk, we consider the existence of a Hamiltonian cycle of G such that all of the vertices of X precede the vertices of Y for appropriate initial vertex and orientation of the cycle.

- 12 斎藤 明 (日大文理) Spanning trees homeomorphic to a small tree 15
 佐野 和貴 (日大文理)
 Akira Saito (Nihon Univ.) Spanning trees homeomorphic to a small tree
 Kazuki Sano (Nihon Univ.)

概要 In this talk, we will present a sufficient condition for a connected graph to have a spanning tree which is homeomorphic to a tree of a bounded order. The result is a natural extension of a classical result for the existence of a hamiltonian path, known as Ore's Theorem. We also discuss the sharpness of the result.

- 13 安藤 清 A degree sum and forbidden subgraph condition for k -contractible edges 10
 (国立情報学研・JST ERATO)
 Kiyoshi Ando A degree sum and forbidden subgraph condition for k -contractible edges
 (Nat. Inst. of Information/JST ERATO)

概要 If a k -connected graph G has no $K_1 + C_4$ and the degree sum of each connected subgraph of G of order 3 is greater than $3k + 1$, then G has a k -contractible edge.

- 14 水澤 篤彦 (早大理工) A construction of smooth travel groupoids on finite graphs 15
 松本ディオゴけんじ
 (早大理工)
 Atsuhiko Mizusawa (Waseda Univ.) A construction of smooth travel groupoids on finite graphs
 Diogo Kendy Matsumoto
 (Waseda Univ.)

概要 In this talk, we show a way to construct smooth travel groupoids from given finite graphs. This gives an answer to Ladislav Nebeský's question "does there exists a connected graph that has no smooth travel groupoid?" on finite graphs.

- 15 藤田 慎也 (横浜市国際総合) A note on covering edge colored hypergraphs by monochromatic components 10
 古谷 倫貴 (東京理大理) A. Gyárfás
 (A. Rényi Inst. of Math.)
 Á. Tóth
 (A. Rényi Inst. of Math.)
 Shinya Fujita (Yokohama City Univ.) A note on covering edge colored hypergraphs by monochromatic components
 Michitaka Furuya (Tokyo Univ. of Sci.) András Gyárfás
 (A. Rényi Inst. of Math.)
 Ágnes Tóth (A. Rényi Inst. of Math.)

概要 For $r \geq 2$, $\alpha \geq r - 1$ and $k \geq 1$, let $c(r, \alpha, k)$ be the smallest integer c such that the vertex set of any non-trivial r -uniform k -edge-colored hypergraph \mathcal{H} with $\alpha(\mathcal{H}) = \alpha$ can be covered by c monochromatic connected components. Here $\alpha(\mathcal{H})$ is the maximum cardinality of a subset A of vertices in \mathcal{H} such that A does not contain any edges. An old conjecture of Ryser is equivalent to $c(2, \alpha, k) = \alpha(r - 1)$ and a recent result of Z. Király states that $c(r, r - 1, k) = \lceil \frac{k}{r} \rceil$ for any $r \geq 3$.

Here we make the first step to treat non-complete hypergraphs, showing that $c(r, r, r) = 2$ for $r \geq 2$ and $c(r, r, r + 1) = 3$ for $r \geq 3$.

- 16 松下 尚弘 (東大数理) 近傍複体のトポロジーと染色数について 15
 Takahiro Matsushita (Univ. of Tokyo) On the topology of neighborhood complexes and the chromatic numbers of graphs

概要 Lovasz introduced the neighborhood complex $N(G)$ of a graph G . He showed that for a graph G , the connectivity of $N(G)$ gives a lower bound for the chromatic number $\chi(G)$ of G . Then a natural question arises: how effective is to determine the topology of the neighborhood complex to compute the chromatic number? For example, is there a topological invariant which is equivalent to the chromatic number? In this talk, we discuss such a problem.

- 17 栗林 勝彦 (信州大理) 擬スキームの強ホモトピー 15
 Katsuhiko Kuribayashi (Shinshu Univ.) On the strong homotopy for quasi-schemoids

概要 A quasi-schemoid is a small category with a particular partition of the set of morphisms. We define a homotopy relation on the category of quasi-schemoids and study the fundamental properties of the homotopy.

- 18 須田 庄 (愛知教育大教育) グラフの再生核ヒルベルト空間とそのグラム行列 10
 瀬戸 道生 (島根大総合理工)
 谷口 哲至 (松江工高専)
 Sho Suda (Aichi Univ. of Edu.) Gram matrices of reproducing kernel Hilbert spaces over graphs I
 Michio Seto (Shimane Univ.)
 Tetsuji Taniguchi
 (Matsue Coll. of Tech.)

概要 We will define a structure of reproducing kernel Hilbert spaces for simple connected graphs. Since graphs are isomorphic if and only if corresponding Hilbert spaces are isomorphic, studying the space is important. In this talk, using the data of the corresponding Gram matrix, we obtain the inequalities for some values on graphs and characterize the some graphs as objects which attain the inequalities.

- 19 瀬戸 道生 (島根大総合理工)* グラフの再生核ヒルベルト空間とそのグラム行列 II 10
 須田 庄 (愛知教育大教育)
 谷口 哲至 (松江工高専)
 Michio Seto (Shimane Univ.) Gram matrices of reproducing kernel Hilbert spaces over graphs II
 Sho Suda (Aichi Univ. of Edu.)
 Tetsuji Taniguchi
 (Matsue Coll. of Tech.)

概要 We study graph homomorphisms from a viewpoint of reproducing kernel Hilbert space theory. In particular, de Branges–Rovnyak theory is introduced into graph theory. We describe graph homomorphisms with Gram matrices of de Branges–Rovnyak spaces.

9月26日(金) 第VIII会場

9:30~11:30

- 20 大堀 龍一 (東大数理) Walsh figure of merit is efficiently approximable 10
 芳木 武仁 (東大数理)
 Ryuichi Ohori (Univ. of Tokyo) Walsh figure of merit is efficiently approximable
 Takehito Yoshiki (Univ. of Tokyo)

概要 Walsh figure of merit (WAFOM) is a quality criterion for a quasi-Monte Carlo integration method, introduced by Matsumoto, Saito and Matoba. They define WAFOM as the sum over the orthogonal complement of the digital net, and compute WAFOM by iterating over the digital net itself. As a further improvement of their method we show an effective approximation method which is easier to implement and faster in computation.

- 21 鈴木航介(東大数理) On the decay of the Walsh coefficients of smooth functions 15
 芳木武仁(東大数理)
 Kosuke Suzuki (Univ. of Tokyo) On the decay of the Walsh coefficients of smooth functions
 Takehito Yoshiki (Univ. of Tokyo)

概要 For the study of quasi-Monte Carlo (QMC) rules for numerical integration of high smooth integrands defined over the s -dimensional unit cube, to analyze the decay of the b -adic Walsh coefficients is important. Josef Dick gave the decay of the Walsh coefficients of functions in Sobolev spaces in 2008. In this talk, we introduce another method to analyze the decay of the Walsh coefficients of smooth functions using iterate integration by parts which improves the bound on Walsh coefficients in some cases.

- 22 芳木武仁(東大数理) The mean square quasi-Monte Carlo error for digitally shifted point sets 15
 合田隆(東大工)
 大堀龍一(東大数理)
 鈴木航介(東大数理)
 Takehito Yoshiki (Univ. of Tokyo) The mean square quasi-Monte Carlo error for digitally shifted point sets
 Takashi Goda (Univ. of Tokyo)
 Ryuichi Ohori (Univ. of Tokyo)
 Kosuke Suzuki (Univ. of Tokyo)

概要 A digitally shifted point set is a point set which can be identified with a subset $P + \sigma$ for an element $\sigma \in Z_b^{s \times n}$ and a subgroup $P \subset Z_b^{s \times n}$, where Z_b denotes the cyclic group of order b . We shall choose σ uniformly and randomly from $\sigma \in Z_b^{s \times n}$. The randomized QMC integration by $P + \sigma$ is the approximation of $I(f) := \int_{x \in [0,1]^s} f(x) dx$ by $I_{P+\sigma}(f) := 1/|P + \sigma| \sum_{x \in P+\sigma} f(x)$. In this talk, we consider the root mean square error $ME_P(f) := 1/b^{sn} \sum_{\sigma \in Z_b^{s \times n}} (I(f) - I_{P+\sigma}(f))^2$. If $ME_P(f)$ is small, $I_{P+\sigma}(f)$ can be small averagely. For a sufficiently smooth integrand f , we obtain a quality measure called Walsh figure of merit $W(P; \mu)$, which gives an upper bound for $ME_P(f)$. We show that $W(P; \mu)$ is computable, and find a point set P with $W(P; \mu)$ small by computer search.

- 23 原瀬晋 Sobol' 列のプロジェクトションについて 15
 (東工大イノベーションマネジメント)
 Shin Harase (Tokyo Tech) Projections of Sobol' sequences

概要 We consider multivariate numerical integration by quasi-Monte Carlo methods. Typical quasi-Monte Carlo point sets are low-discrepancy point sets based on the t -values of (t, m, s) -nets. In particular, Sobol' sequences are famous examples and widely used in computational finance. In this talk, we focus on Sobol' sequences with better two-dimensional projections constructed by Joe and Kuo (2008). We investigate the t -values of higher dimensional projections, and give some remarks.

- 24 室 伏 俊 明 (東工大総合理工) 単調優モジュラ集合関数の加法分解可能性に関する計算機を用いた判定
堀 尾 尚 史 (東工大総合理工) 15
Toshiaki Murofushi (Tokyo Tech) A computational approach for testing additive decomposability of mono-
Takafumi Horio (Tokyo Tech) tone, supermodular set functions

概要 In this paper we consider additive decompositions by proper monotone set functions, i.e., monotone set functions with proper support, and investigate whether it is possible to have such decompositions for monotone, supermodular set functions, which we call a “decomposability question”. We propose a computational approach by formulating the polytope of monotone, supermodular set functions by means of hyperplanes, and by examining the corresponding vertices of polytope for further decomposability over proper monotone set functions. This enables us to numerically resolve the decomposability question in a finite steps of procedure. We have tested the proposed method for five- and six-element sets; we have obtained a positive answer to the question for a five-element set, and a negative one for a six-element set.

- 25 藤 本 実 (精華科学研) NP 完全問題の平方剰余による解法 15
上 原 邦 彦 (帝塚山大経営)
Minoru Fujimoto (Seika Science Lab.) NP Complete problem by quadratic residue problem
Kunihiko Uehara (Tezukayama Univ.)

概要 We study the quadratic residue problem known as an NP complete problem by way of the prime number and show that a nondeterministic polynomial process does not belong to the class P because of a random distribution of solutions for the quadratic residue problem.

- 26 生 田 卓 也 (神戸学院大法) Complex Hadamard matrices attached to some association schemes .. 15
Takuya Ikuta (Kobe Gakuin Univ.) Complex Hadamard matrices attached to some association schemes

概要 Recently, we classified type-II matrices attached to some association schemes. We have 6 infinite families of type-II matrices, among which are 4 families of our complex Hadamard matrices. To check whether our complex Hadamard matrices are inequivalent to the tensor product of known examples, we introduce the Haagerup set for type-II matrices and the Nomura algebra for our complex Hadamard matrices. In this talk, we mainly present how to compute the Haagerup set and the Nomura algebra for our complex Hadamard matrices. This is based on a joint work with Akihiro Munemasa.

13:00~14:00 特別講演

- 瀬 川 悦 生 (東北大情報) 量子ウォークの固有値写像について
Etsuo Segawa (Tohoku Univ.) Spectral mapping of quantum walks

概要 For a given finite graph $G = (V, E)$, we present a spectral mapping theorem onto a discrete-time quantum walk on $\ell^2(D)$ from the underlying cellular automaton on $\ell^2(V)$, where D is the set of the symmetric arcs generated by E . Firstly, we show that the cellular automaton satisfies a difference of a wave equation, and provides the real parts of the spectrum of the quantum walk at each time $n \in \mathbb{N}$. We denote this inherited eigenspace of the quantum walk from the underlying cellular automaton by \mathcal{L} . Secondly, we show that in the Grover walk case, which is intensively studied from the view points of quantum search algorithms and scattering theories, the orthogonal complement eigenspace \mathcal{L} is generated by a homological structure of the graph. Finally, we extend our notion to an infinite abelian covering graph. As an application, we discuss the following typical stochastic properties of quantum walks; localization and linear spreading.

9月27日(土) 第VIII会場

9:30~11:55 特別セッション「移動境界問題の数理解析」

- 石 渡 哲 哉 Motion of polygonal curves by crystalline curvature flow 45
 (芝浦工大システム理工)
- Tetsuya Ishiwata Motion of polygonal curves by crystalline curvature flow
 (Shibaura Inst. of Tech.)

概要 In this talk we consider the motion of planar polygonal curves governed by generalized crystalline curvature flow with a driving force: $\beta(N_j)V_j = U - g(H_j)$, where V_j , N_j and H_j denote an outward velocity, an outward normal vector and a crystalline curvature of the j -th *facet* of solution curve, respectively. Here, “facet” means a lines segment of solution curve $\Gamma(t)$. The positive function β and U describe an anisotropy of the mobility and a driving force, respectively. We consider 2 cases. One is the case when solution curves are closed. The other is the case when solution curves have infinity length. For both cases, we discuss a behavior of polygonal curves, especially, we mainly consider a deformation of solution curves.

- 木 村 正 人 (金 沢 大 理 工) Generalization of crystalline motion 45
 Masato Kimura (Kanazawa Univ.) Generalization of crystalline motion

概要 We consider a polygonal analogue of the Hele-Shaw moving boundary problem with surface tension based on a framework of polygonal motion proposed by Benes–Kimura–Yazaki (2009). This is a generalization of the crystalline motion for the curvature flow which was originally proposed by Angenent–Gurtin (1989) and J. E. Taylor (1993).

A key idea is to introduce a polygonal Dirichlet-to-Neumann map. We study variational properties of the polygonal Dirichlet-to-Neumann map and show that our polygonal Hele-Shaw problem is an almost complete analogue of the original problem. Local solvability of a polygonal Hele-Shaw problem is proved by means of the variational structure. We also give several polygonal versions of moving boundary problems such as the curvature flow, the Hele-Shaw flow and the Stokes flow etc. with some numerical examples. Our polygonal motions exactly satisfy properties of curve shortening and/or area preservation.

This is a joint work with Shigetoshi Yazaki (Meiji University) and Daisuke Tagami (Kyushu University).

- 田 端 正 久 (早 大 理 工) 混相流の数値解析 45
 Masahisa Tabata (Waseda Univ.) Numerical analysis of multiphase flows

概要 Multifluid and multiphase flows with surface tension are encountered frequently in scientific and engineering problems. Recently we have developed an energy-stable Galerkin-characteristics scheme to two-fluid flow problems with surface tension, where each flow is governed by the Navier–Stokes equations. After presenting excellent features of this scheme, we simulate numerically some multiphase flow problems such as rising bubbles and falling droplets by the use of it.

14:15~16:30

- 27 岩見真吾 (九大理) タイムラグを持つ微分方程式によるウイルス感染動態の定量的解析 15
 Shingo Iwami (Kyushu Univ.) Quantitative analysis of virus infection dynamics with distributed delay differential equations

概要 In this study, we were interested in quantifying and modeling the epliptic phase during virus infection. One simple mathematical approach describing this phase is modeling by delay differential equations (DDE). However, unfortunately, the distribution of the epliptic phase was poorly understood. Here, from cell culture experiments with simian/human immunodeficiency virus (SHIV), we quantified the distribution of the epliptic phase and found that the phase obeys gamma distribution. Based on this experimental result, we made a mathematical model including the distribution by DDE and estimated viral parameters from previously published experimental data. Our analyses showed that modeling the epliptic phase affected estimation of the virus infection rate (therefore, basic reproductive number) but not the virus production rate and the death rate of infected cells.

- 28 國谷紀良 (神戸大システム情報) 空間拡散を考慮に入れた年齢構造化 SIS 感染症モデルの解析 15
 Toshikazu Kuniya (Kobe Univ.) Analysis of an age-structured SIS epidemic model with spatial diffusion

概要 In this talk, we focus on an age-space-structured SIS epidemic model which is formulated as a nonlinear diffusion system of partial differential equations. In order to apply a traditional approach of integral operators, we use the Feynman-Kac formula in probability theory and express the solution of the system explicitly in terms of expected value. After that we reformulate the problem of existence of the solution to a fixed-point problem for a nonlinear operator and show that such a fixed-point exists if and only if the spectral radius of the next generation operator, that is, the well-known basic reproduction number R_0 is greater than unity.

- 29 平岡裕章 (九大IMI) 可換梯子籠上でのパーシステント加群 15
 E. G. Escolar (九大数理)
 Yasuaki Hiraoka (Kyushu Univ.) Persistence modules on commutative ladder quivers
 Emerson G. Escolar (Kyushu Univ.)

概要 In this talk, we generalize the concept of persistence modules on A_n quivers to commutative ladder quivers. This class of persistence modules frequently appears in topological data analysis, and the theory and algorithm proposed in this talk can be applied to these practical problems. A new algebraic framework deals with persistence modules as representations on associative algebras and the Auslander-Reiten theory is applied to develop the theoretical and algorithmic foundation.

- 30 E. G. Escolar (九大数理) Computing persistence modules on commutative ladders of finite type
平岡裕章 (九大IMI) 15
Emerson Gaw Escolar (Kyushu Univ.) Computing persistence modules on commutative ladders of finite type
Yasuaki Hiraoka (Kyushu Univ.)

概要 In topological data analysis, the idea of persistence modules on commutative ladders is a recent development extending the ideas of persistent homology to a more general setting. In particular, it can be used to study common robust topological features. In this talk, we discuss the computation of persistence modules on commutative ladders of finite type, focusing on the use of Morse reductions to reduce computation times.

- 31 草野元紀 (九大数理) センサーネットワーク被覆問題へのパーシステントホモロジーの応用 .. 15
平岡裕章 (九大IMI)
Genki Kusano (Kyushu Univ.) An application of persistent homology to a coverage problem in sensor
Yasuaki Hiraoka (Kyushu Univ.) networks

概要 A coverage problem in sensor networks was partially solved by V. de Silva and R. Ghrist by using homology. In this talk, we apply persistent homology to the coverage problem, and present a coverage theorem with perturbation.

- 32 高木 悟 (工学院大) 狩野モデルを応用した大学数学講義における学生ニーズ分析 15
上江洲弘明 (早大GEC)
Satoru Takagi (Kogakuin Univ.) An analysis of students' needs for undergraduate mathematics lectures
Hiroaki Uesu (Waseda Univ.) applying the Kano model

概要 At universities, mathematics lecturers have to change their teaching materials and methods according to students' mathematical skills, and also have to make them understand mathematics. In this paper, we apply the questionnaire analysis in some undergraduate mathematics lectures for the method of students' needs analysis using the Kano model, and report the results.

- 33 中根和昭 (阪大医) 位相幾何学的手法を用いた画像解析技術について I (病理組織に対する応用) 15
Kazuaki Nakane (Osaka Univ.) Image analysing method via homology —for cancer area detection—

概要 Development of technology for detecting of the cancer lesions from digitized pathological specimens has been studied for a long time. This type of system has been developed based on the pattern recognition technology. Because the cancer tissues have too various forms, this technology does not work well. Here we introduce a new method. By using a mathematical theory (homology theory), our system has been designed. Our method can estimate the contact degree of elements in the tissue of unit area. We may consider that, by the loss of contact inhibition, cancer lesions are included in high contact degree area. We have tried this method to pathological images and obtained satisfactory good results.

- 34 中根和昭(阪大医) 位相幾何学的手法を用いた画像解析技術についてII(一般組織に対する応用) 15
 Kazuaki Nakane (Osaka Univ.) Image analysing method via homology II —for general structures—

概要 There are structures that are formed with a certain order. For these kind of structures, research is progressing from both the theory and experiment. There are many examples we cannot find the rule of structures at first glance. In these cases, it is difficult to make mathematical description. Some of the structures, we apply the topological method and attempt to classify. The structures are constituted by the contact between the components. Homology can also be considered the concept of evaluating the degree of contact. We set the size of the unit area and calculate the indexes which are depended on the Betti numbers. This is the principle that quantify the degree of contact.

16:45~17:45 特別講演

- 大下承民(岡山大自然) Motion of droplets driven by curvature and potential
 Yoshihito Oshita (Okayama Univ.) Motion of droplets driven by curvature and potential

概要 We consider the motion of small droplets driven by the curvature and the potential in a bounded domain and show that the center of the droplets moves along a solution of some reduced equation in a small volume fraction limit and compare its homogenization limit as well.

9月28日(日) 第VIII会場

9:30~11:50

- 35 堀口俊二(新潟産大経済) 土倉・堀口法(村瀬義益・ニュートン型の第一拡張漸化式)の収束比較条件式III 10
 Shunzi Horiguchi The conditional expressions III which compare convergences of Tsuchikura–Horiguchi method (the first extended recurrence formula of Yoshimasu Murase–Newton’s type)
 (Niigata Sangyo Univ.)

概要 We obtained extension of Newton–Raphson method (Tsuchikura–Horiguchi method). We give the conditional expressions III which compare convergences of Tsuchikura–Horiguchi method (the first extended recurrence formula of Yoshimasu Murase–Newton’s type).

- 36 堀口俊二(新潟産大経済) 土倉・堀口法(村瀬義益・ニュートン型の拡張漸化式)の収束比較条件式IIIの数値計算 10
 Shunzi Horiguchi The numerical computations of the conditional expressions III which compare convergences of Tsuchikura–Horiguchi method (the first extended recurrence formula of Yoshimasu Murase–Newton’s type)
 (Niigata Sangyo Univ.)

概要 We give the numerical computations of the conditional expressions III which compare convergences of Tsuchikura–Horiguchi method (the first extended recurrence formula of Yoshimasu Murase–Newton’s type).

- 37 及川一誠 (早大理工) A hybridized discontinuous Galerkin method with reduced stabilization 15
 Issei Oikawa (Waseda Univ.) A hybridized discontinuous Galerkin method with reduced stabilization

概要 In this talk, we propose reduced stabilization of the hybridized discontinuous Galerkin methods. We show error estimates and numerical results. A new theoretical tool for analysis of the proposed method is also proposed.

- 38 内海晋弥 (早大理工) Navier-Stokes 方程式のための数値積分誤差を伴わない特性曲線有限要素
 田端正久 (早大理工) スキームの解析 15
 Shinya Uchiumi (Waseda Univ.) Analysis of a finite element scheme free from quadrature errors for the
 Masahisa Tabata (Waseda Univ.) Navier-Stokes equations

概要 We consider the characteristics finite element method for the Navier-Stokes equations. The conventional scheme cannot be implemented precisely because the composite function terms are not polynomials. In the analysis of the conventional scheme, however, it is assumed that the terms are integrated exactly. We have developed a scheme free from quadrature errors that can be precisely implemented for the convection-diffusion equations. Here we show that the scheme can be extended to the Navier-Stokes equations to achieve an optimal convergence result.

- 39 柏原崇人 (TU Darmstadt) Stokes 滑り境界問題に対するペナルティ法とその誤差解析 15
 周冠宇 (東大数理)
 及川一誠 (早大理工)
 Takahito Kashiwabara (TU Darmstadt) Penalty method to the Stokes problem with slip boundary condition
 Guanyu Zhou (Univ. of Tokyo)
 Issei Oikawa (Waseda Univ.)

概要 The penalty method for Stokes problem with slip boundary condition is concerned. We derive the optimal error estimate of penalty method. The smooth boundary is approximated by straight polygonal lines, then finite element method is applied to solve the penalty problem. The error estimate of finite element approximation is achieved. Some numerical experiments are presented to verify our theoretical results.

- 40 剣持智哉 (東大数理) 抽象的 Cauchy 問題に対する離散最大正則性と有限要素法への応用 15
 齊藤宣一 (東大数理)
 Tomoya Kemmochi (Univ. of Tokyo) Discrete maximal regularity for abstract Cauchy problems and its ap-
 Norikazu Saito (Univ. of Tokyo) plication to finite element methods

概要 Maximal regularity is one of fundamental concepts in theory of partial differential equations. It, for example, provides us a convenient way to establish the well-posedness for a wide class of nonlinear equations. Therefore, it is natural to ask whether a discrete analogue of maximal regularity is available. Furthermore, we are interested in application of discrete maximal regularity to error analysis of various discretization of nonlinear equations. In this paper, we first prove a maximal regularity result for time-discrete abstract Cauchy problem in a UMD space. Then, we also report an application of such discrete maximal regularity result to the finite element method for the heat equation and obtain reasonable a priori estimate.

- 41 杉谷 宜紀 (東大数理) Finite element approximation for the Stokes equations under a unilateral boundary condition 15
 周冠宇 (東大数理)
 齊藤 宣一 (東大数理)
 Yoshiki Sugitani (Univ. of Tokyo) Finite element approximation for the Stokes equations under a unilateral boundary condition
 Guanyu Zhou (Univ. of Tokyo)
 Norikazu Saito (Univ. of Tokyo)

概要 As an open boundary condition (a kind of artificial boundary conditions) for viscous incompressible fluids flow simulation, we propose a unilateral boundary condition. With our boundary condition, the Navier–Stokes equation satisfies an energy inequality so that we can avoid numerical instability in 3D simulations. In this paper, we restrict ourselves to a model Stokes problem and report some results on the finite element approximation.

- 42 上田 祐暉 (東大数理) B-spline に基づく高精度逐次的時間離散化法の解析 15
 齊藤 宣一 (東大数理)
 Yuuki Ueda (Univ. of Tokyo) On a high-accurate successive time-discretization method based on B-spline interpolation functions
 Norikazu Saito (Univ. of Tokyo)

概要 Recently, K. Takizawa and T. Tezduyar proposed a new time discretization method, which they call space-time computation techniques with continuous representation in time (ST-C), for fluid-structure interaction problem. The method is based on B-spline interpolation functions and, consequently, is highly accurate. In this paper, we examine the successive projection technique (SPT), which is a most basic technique among ST-C, and report the well-posedness of the algorithm and a certain error estimation.

- 43 大塚 厚二 一般 J 積分を用いた随伴法による形状最適化問題の解法 15
 (広島国際学院大情報デザイン)
 Khoji Ohtsuka Solution of the shape optimization problem with Generalized J-integral by adjoint method
 (Hiroshima Kokusai Gakuin Univ.)

概要 Let Ω be a domain on which the boundary value problem is defined and $u(\Omega)$ the solution. We study the shape optimization problem to find the shape which is optimal in that it minimizes the cost functional $\int_{\Omega} g(u(\Omega)) dx$ with the volume constraint, $|\Omega| = \text{constant}$. In this study, we talk how to solve shape optimization problems with Generalized J-integral coming from fracture mechanics, H1-gradient method proposed by Prof. Azegami and adjoint method.

14:15~16:15

- 44 穴田 浩一 (早大高等学院) ある非線形放物型偏微分方程式の解の特徴について 15
 石渡 哲哉
 (芝浦工大システム理工)
 Koichi Anada Some features for blow-up solutions of a nonlinear parabolic equations
 (Waseda Univ. Senior High School)
 Tetsuya Ishiwata
 (Shibaura Inst. of Tech.)

概要 In previous studies we have shown some conjectures for behavior of blow-up solutions to a nonlinear parabolic equations. They are very important features to investigate behavior of solutions near their blow-up time. The purpose of our paper is to prove one of them that we call weak eventual monotonicity.

- 45 佐々木多希子 (東大数理) 非線形シュレディンガー方程式の差分解の爆発について 15
齊藤宣一 (東大数理)
Takiko Sasaki (Univ. of Tokyo) Blow-up of finite difference solutions for nonlinear Schrödinger equations
Norikazu Saito (Univ. of Tokyo)

概要 The purpose of this paper is to establish a numerical method for computing blow-up solutions of one-dimensional nonlinear Schrödinger equations with power nonlinearities. There are various techniques for computing blow-up solutions of nonlinear heat and wave equations. However, a little is known for the nonlinear Schrödinger equations. In this paper, we propose a finite difference scheme that can approximate the blow-up phenomena. Thus, after having established convergence results, we show that the numerical blow-up time actually converges to the blow-up time of the nonlinear Schrödinger equation under consideration.

- 46 田上大助 (九大IMM) ゲージ条件を考慮した渦電流問題に対する反復型領域分割法 15
Daisuke Tagami (Kyushu Univ.) An iterative domain decomposition method for eddy current problems with the gauge condition

概要 An iterative domain decomposition method is introduced into eddy current problems, which is formulated by a mixed method with the Lagrange multiplier. By introducing the mixed method, a two-subdomain problem is equivalent to the conventional one-domain problem, and yields an iterative domain decomposition method of eddy current problems. Moreover, to mathematically justify the previous formulation, the formulation based on the mixed method has been established a connection with the previous one.

- 47 榊原航也 (東大数理) 一相 Hele-Shaw 流れの代用電荷法による数値計算 15
矢崎成俊 (明大理工)
Koya Sakakibara (Univ. of Tokyo) Numerical computation of the one phase Hele-Shaw flow based on the
Shigetoshi Yazaki (Meiji Univ.) charge simulation method

概要 We construct the variational structure-preserving numerical scheme for the one phase Hele-Shaw flow and examine its effectivity numerically. We modify the CSM invariant scheme in order to satisfy the area-preserving property by posing a condition that the weighted average of charges is equal to 0.

- 48 宮路智行 (京大数理研) 四つ葉状の軌道をもつ三次元常微分方程式系に対する計算機援用解析 .. 15
Tomoyuki Miyaji (Kyoto Univ.) Computer-assisted analysis to dynamical systems with four-leaf orbits

概要 We study a three-dimensional dynamical system defined by ordinary differential equations. Craik has shown that an exact solution to 3D Navier-Stokes equation is constructed by using a solution to the equations. Craik and Okamoto have found out that almost all orbits of the system are unbounded and an unstable periodic orbit plays an important role in determining the destination of an unbounded orbit. We prove the existence of such a periodic orbit by a method of numerical verification based on interval arithmetic. When passing through the vicinity of the periodic orbit, an orbit draws a four-leaf. If one adds damping terms to the system, one can observe a four-leaf chaotic attractor. We also study the route to four-leaf chaos.

- 49 三宅常時 (宇部工高専) セパトリックスループになるパラメータの計算法 15
勝田祐司 (宇部工高専)
 George Miyake A method of computing parameter values for shaping a limit cycle into
 (Ube Nat. Coll. of Tech.) a separatrix loop
 Yuji Katsuta (Ube Nat. Coll. of Tech.)

概要 Where there is an adhesion of a stable or unstable limit cycle to a saddle equilibrium point in nonlinear systems, the limit cycle is made up into separatrix loop. It is known the computational method of parameter values for shaping the limit cycle into the separatrix loop by using numerical integration on the alpha branch streaming out from the equilibrium point. The alpha branch is likely to happen because the instability of the alpha branch is to prevent the branch returning to in the neighborhood of equilibrium point, as a result a new computational method using bisection method is brought in.

- 50 関坂歩幹 (東北大理) 反応拡散系における進行波の安定性と絶対スペクトルの関係 15
 Ayuki Sekisaka (Tohoku Univ.) The relationship between the stability of pulse and the absolute spectrum in reaction diffusion system

概要 Can two unstable waves make a stable pulse? This problem was solved affirmatively by Nii and Sandstede–Scheel, independently. In particular, The stability of pulse is connected with absolute spectrum. But, their results depends on the codimension of a bifurcation. In this talk, we show that an extension between the stability of pulse and the absolute spectrum with higher codimension bifurcation.

16:30~17:30 特別講演

- 高石武史 フェーズフィールドによるき裂進展モデルとコンピュータによる検証
 (広島国際学院大総合教育センター)
 Takeshi Takaishi A phase-field model for crack growth and its numerical verification
 (Hiroshima Kokusai Gakuin Univ.)

概要 A phase field model for crack growth in two or three dimensional isotropic elastic material is proposed by the author and Kimura. A phase-field to represent the shape of the crack with a regularization parameter $\epsilon > 0$ is introduced in the model. The phase field model is derived as a gradient flow of this regularized energy that is approximated by the Francfort–Marigo type energy using the idea of Ambrosio and Tortorelli. The model is convenient to study the mathematical property of crack growth for theoretical and numerical analysis, such as the equilibrium profiles of the phase-field. Numerical results shows the two or three-dimensional structure of crack surface.

トポロジー

9月25日(木) 第VI会場

9:20~12:00

- 1 比嘉一晃 (高知大理) A surviving condition on permanent cycles in the Adams E_2 -term \cdots 10
 加藤 諒
 下村克己 (高知大理)
 Kazuaki Higa (Kochi Univ.) A surviving condition on permanent cycles in the Adams E_2 -term
 Ryo Kato
 Katsumi Shimomura (Kochi Univ.)

概要 For a nontrivial element in the Adams spectral sequence, we give a new condition for surviving to the stable homotopy groups of spheres. It may be seen as a generalization of many works of Xiugui Liu and collaborators.

- 2 宮沢健太郎 (高知大総合人間自然) On the action of Greek letter element β_1 in the stable homotopy groups
 下村克己 (高知大理) of spheres \cdots 10
 Kentaro Miyazawa (Kochi Univ.) On the action of Greek letter element β_1 in the stable homotopy groups
 Katsumi Shimomura (Kochi Univ.) of spheres

概要 In this talk, we consider the action of β_1 in the stable homotopy groups of spheres using the method's of Ravenel's small descent spectral sequence.

- 3 南 範彦 (名工大) Beilinson–Roeblyum の無限ループ空間 Hurewicz 関手 isogeny 定理につ
 いて \cdots 15
 Norihiko Minami On the Beilinson–Rosenblyum isogeny theorem of the Hurewicz functor
 (Nagoya Inst. of Tech.) for infinite loop spaces

概要 I shall offer a slightly streamlined proof of the Beilinson–Rosenblyum isogeny theorem of the Hurewicz functor for infinite loop spaces. Universal torsion order estimates of the kernel and the cokernel of the Hurewicz homomorphism functor are also slightly improved.

- 4 板垣早紀 (高知大総合人間自然) 有限位相空間の圏と有限単体的複体の圏の対応について \cdots 10
 逸見 豊 (高知大理)
 Saki Itagaki (Kochi Univ.) Correspondence between categories of finite topological spaces and finite
 Yutaka Hemmi (Kochi Univ.) simplicial complexes

概要 There is a functor from the category of finite topological spaces $\mathcal{F}top$ to the category of finite simplicial complexes. This functor is faithful but is not full. To make this functor full, we consider a weaker version of continuity of maps between finite topological spaces and consider an extend category $\mathcal{F}top_{ex}$ of $\mathcal{F}top$. Moreover, we can define the homotopy in $\mathcal{F}top_{ex}$ so that two finite topological spaces have the same homotopy type in $\mathcal{F}top_{ex}$ if and only if the corresponding simplicial complexes have the same strong homotopy type.

- 5 松下尚弘 (東大数理) The simplicial sets related to the Hom complexes of graphs 10
Takahiro Matsushita (Univ. of Tokyo) The simplicial sets related to the Hom complexes of graphs

概要 For a positive integer r , an r -set we say in this talk is a set X with a subset of the r -times direct product X^r of X . For r -sets X and Y , we construct the poset $\text{Hom}(X, Y)$ called the Hom complex, and the simplicial set $\text{Sing}(X, Y)$ called the singular complex. We show that their geometric realizations are homotopy equivalent. The Hom complex of r -sets is the generalization of the Hom complexes of graphs which have been applied to the graph coloring problem in graph theory.

- 6 鈴木直矢 (名大多元数理) 単体的ド・ラーム複体上の Dixmier–Douady 類の表示 15
Naoya Suzuki (Nagoya Univ.) The Dixmier–Douady class in a simplicial de Rham complex

概要 Let G be a Lie group and BG a classifying space of G . Then we can recognize a characteristic class of principal G -bundles as an element in $H^*(BG)$. In general BG is a very huge space so we can not use the ordinary de Rham cohomology theory on it. However if use the simplicial de Rham theory due to mainly Bott–Shulman–Stasheff and Dupont, we can exhibit a cocycle which represents an element in $H^*(BG)$ as a differential form on a simplicial manifold NG . In this presentation I will exhibit a cocycle in a simplicial de Rham complex which represents the Dixmier–Douady class in the sense of Carey–Crowley–Murray. It is a characteristic class of a principal G -bundle whose fiber has a central $U(1)$ -extension.

- 7 西信洋和 (高知大理) Certain examples of posets of rational Gottlieb subgroups 10
山口俊博 (高知大教育)
Hirokazu Nishinobu (Kochi Univ.) Certain examples of posets of rational Gottlieb subgroups
Toshihiro Yamaguchi (Kochi Univ.)

概要 Let $G_*^\xi(X)$ be the fibre-restricted Gottlieb group with respect to a fibration $\xi : X \rightarrow E \rightarrow Y$ in CW complexes. It is a subgroup of the Gottlieb group $G_*(X)$ of X . When $Y = BS^1$ and X is the product of odd-spheres, we illustrate a variety of Gottlieb poset, which is the poset of subspaces $G_*^\xi(X) \otimes \mathbb{Q}$ in $G_*(X) \otimes \mathbb{Q}$.

- 8 栗林勝彦 (信州大理) Hopf 空間のストリングトポロジー 10
Katsuhiko Kuribayashi (Shinshu Univ.) Loop products on Noetherian Hopf spaces

概要 We investigate the loop homology of a simply-connected Noetherian Hopf spaces. It turns out that non-triviality of the loop product characterizes the finiteness of the cohomology of such Hopf spaces.

- 9 畑中美帆 (阪市大理) グラフに対応するスピントリック多様体 10
Miho Hatanaka (Osaka City Univ.) Spin toric manifolds associated to graphs

概要 We describe a necessary and sufficient condition for a toric manifold to admit a spin structure. This implies that a toric manifold admits a spin structure if and only if its real part is orientable. It is known that a Delzant polytope can be constructed from a simple graph, so that one can associate a toric manifold to a simple graph. We characterize simple graphs whose associated toric manifolds admit spin structures.

- 10 阿部 拓 (阪市大数学研) ルート系から定まるトーリック多様体の交叉数とヤング図 15
 Hiraku Abe (Osaka City Univ.) Young diagrams and intersection numbers for toric manifolds arising from root systems

概要 We study intersection numbers in the toric manifold associated with the fan determined by the Weyl chambers for the root system of type A. We give a combinatorial formula for intersection numbers of certain subvarieties which are naturally indexed by elements of the Weyl group. These numbers describe the ring structure of the cohomology of the toric manifold.

- 11 堀口達也 (阪市大理) ピーターソン多様体の同変コホモロジー 10
 原田芽ぐみ (McMaster Univ.)
 栢田幹也 (阪市大理)
 Tatsuya Horiguchi (Osaka City Univ.) The equivariant cohomology rings of Peterson varieties
 Megumi Harada (McMaster Univ.)
 Mikiya Masuda (Osaka City Univ.)

概要 Let G be a complex semisimple linear algebra group and let Pet be the Peterson variety in the flag variety G/B . The main theorem gives an efficient presentation of the S -equivariant cohomology ring of the Peterson variety as a quotient of a polynomial ring by an ideal J generated by quadratic polynomials. Here the group S is a certain circle subgroup of a maximal torus T of G . Our description of the ideal J uses the Cartan matrix and is uniform across Lie types. Our result generalizes a previous theorem of Fukukawa–Harada–Masuda, which was only for Lie type A.

- 12 山口崇幸 (広島大理) Universal finite type invariant による rank が 5 以下の Gauss words の
 福永知則 (北大理) 完全な分類 10
 山野井隆晃 (北大理)
 Takayuki Yamaguchi (Hiroshima Univ.) Complete classification of Gauss words of rank less than or equal to 5
 Tomonori Fukunaga (Hokkaido Univ.) by universal finite type invariant
 Takaaki Yamanoi (Hokkaido Univ.)

概要 We show complete classification of Gauss words of rank less than or equal to 5, which is obtained by the universal finite type invariant of degree 7. We also show truncated Polyak algebras of Gauss words of rank less than or equal to 7. The invariant and the truncated Polyak algebras have been calculated by our computer program.

14:15~15:25

- 13 矢島幸信 (神奈川大工)* 無限積空間における D -空間性 10
 平田康史 (神奈川大工)
 Yukinobu Yajima (Kanagawa Univ.) The D -space property of infinite products
 Yasushi Hirata (Kanagawa Univ.)

概要 It is proved by an elementary submodel that any countable product of subparacompact DC -like spaces is a D -space. More generally, we prove that any uncountable product of subparacompact DC -like spaces is a D -space if and only if each factor of the product is compact except countably many ones.

- 14 矢ヶ崎達彦 (京都工繊大工芸) Homeomorphism groups of non-compact surfaces endowed with the Whitney topology 15
 T. Banakh
 (Ivan Franko Nat. Univ. of Lviv)
 嶺幸太郎 (東大数理)
 酒井克郎 (神奈川大工)
 Tatsuhiko Yagasaki (Kyoto Inst. Tech.) Homeomorphism groups of non-compact surfaces endowed with the Whitney topology
 Taras Banakh
 (Ivan Franko Nat. Univ. of Lviv)
 Kotaro Mine (Univ. of Tokyo)
 Katsuro Sakai (Kanagawa Univ.)

概要 We study topological type of the homeomorphism group $\mathcal{H}(M)$ of any non-compact connected surface M endowed with the Whitney topology and show that the identity connected component $\mathcal{H}_0(M)$ of $\mathcal{H}(M)$ is homeomorphic to the product $l_2 \times \mathbb{R}^\infty$.

- 15 下村尚司 (名古屋経済大) Ergodic measures and graph circuits of a sequence of covers 10
 Takashi Shimomura Ergodic measures and graph circuits of a sequence of covers
 (Nagoya Univ. of Econ.)

概要 We have shown that every 0-dimensional system is an inverse limit of a sequence of finite directed graph covers. In this expression, all invariant measures are expressed as a limit of barycenters of circuits of each finite directed graph. Consider the case in which each system of all circuits of a graph is linearly independent. Then, a condition for the invariant measure to be ergodic is given. This condition seems to mean that the time average and the space average coincide.

- 16 下村尚司 (名古屋経済大) Combinatorial construction of completely scrambled compact system · 10
 Takashi Shimomura Combinatorial construction of completely scrambled compact system
 (Nagoya Univ. of Econ.)

概要 Since Li and Yorke developed the notion of scrambled sets in the study of chaotic systems, there has been some discussion as to how large such sets can be. In 1997, Mai reported a non-compact example that is completely scrambled, i.e., the scrambled set is the whole space, and conjectured that there was no compact example. Huang and Ye later disputed this conjecture. They constructed a compact, 0-dimensional completely scrambled system. They also reported the existence of locally equicontinuous transitive examples. We construct a set of examples that are completely scrambled, transitive and not locally equicontinuous.

- 17 山内貴光 (愛媛大理) 整数群の可算直和の粗い幾何学における無限次元性について 15
 Takamitsu Yamauchi (Ehime Univ.) On coarse geometric infinite-dimensionality of the countable direct sum of the integers

概要 Asymptotic property C is an infinite-dimensional property in coarse geometry in the sense that every metric space with finite asymptotic dimension has asymptotic property C. It was introduced by Dranishnikov as a coarse geometric analogue of Haver's property C in dimension theory. In this talk, we discuss a question of Dranishnikov and Zarichnyi which asks whether the countable direct sum of the integers has asymptotic property C.

15:45~16:45 特別講演

- B. Bowditch * Rigidity results for spaces associated to a surface
(Univ. of Warwick・東工大情報理工)
- Brian Bowditch Rigidity results for spaces associated to a surface
(Univ. of Warwick/Tokyo Tech)

概要 A quasi-isometry between two metric spaces is a map which preserves their large-scale geometry, or more precisely, distances to within fixed linear bounds. An important aspect of geometric group theory is to describe the possible quasi-isometries between various naturally occurring spaces. For example, it was shown recently by Behrstock, Kleiner, Minsky and Mosher that (apart from a few exceptional cases) the mapping class group of a compact orientable surface is quasi-isometrically rigid. That is, any self quasi-isometry of (a Cayley graph of) the mapping class group agrees, up to bounded distance, with left multiplication by an element of the group. We aim to describe some of the ideas behind this, as well as some strengthenings and variations to other spaces such as the Weil–Petersson metric on Teichmüller space. Our approach makes use of the theory of median algebras.

9月26日(金) 第III会場

10:30~10:45 2014年度幾何学賞授賞式**10:50~11:50 2014年度幾何学賞受賞特別講演(幾何学分科会と合同) —倉西正武氏の受賞を記念して—**

後藤 竜司 (阪大 理) 倉西数学 —変形理論および CR 幾何について—
宮嶋 公夫 (鹿児島大*)

Ryushi Goto (Osaka Univ.) Kuranishi's masterpieces and their developments in deformation theory
Kimio Miyajima (Kagoshima Univ.*) and CR-geometry

概要 In this talk, we discuss Kuranishi's works and their developments focussing on deformation theory and CR-geometry.

第VI会場

13:00~14:15

- 18 鈴木 正明 (明大総合数理) Integral Euler characteristic of Out F_{11} 10
逆井 卓也 (東大数理)
森田 茂之 (東大*・東工大*)
- Masaaki Suzuki (Meiji Univ.) Integral Euler characteristic of Out F_{11}
Takuya Sakasai (Univ. of Tokyo)
Shigeyuki Morita
(Univ. of Tokyo*/Tokyo Tech*)

概要 The integral Euler characteristic of Out F_{11} is -1202 .

- 19 辻 俊輔 (東大数理)^b 非有向曲面におけるデーンツイストの対数 15
Shunsuke Tsuji (Univ. of Tokyo) The logarithms of Dehn twists on non-orientable surfaces

概要 We introduce a Lie algebra associated with a non-orientable surface, which is an analogue for the Goldman Lie algebra of an oriented surface. As an application, we deduce an explicit formula of the Dehn twist along an annulus circle on the surface as in Kawazumi–Kuno and Massuyeau–Turaev.

- 20 河澄響矢 (東大数理) Turaev 余括弧積, 榎本・佐藤トレースそして柏原 Vergne 問題の発散コサイクル 15
Nariya Kawazumi (Univ. of Tokyo) The Turaev cobracket, the Enomoto–Sato traces and the divergence cocycle in the Kashiwara–Vergne problem

概要 We interpret the divergence cocycle in the Kashiwara–Vergne problem and the Enomoto–Sato obstructions for the surjectivity of the Johnson homomorphisms as some part of a regular homotopy version of the Turaev cobracket.

- 21 石田智彦 (京大理) ハンドル体写像類群のねじれ係数 1 次元ホモロジー群について 10
佐藤正寿 (岐阜大教育)
Tomohiko Ishida (Kyoto Univ.) A twisted first homology of the handlebody mapping class group
Masatoshi Sato (Gifu Univ.)

概要 We compute first homology of the handlebody mapping class group with twisted coefficient.

- 22 石田智彦 (京大理) 体積に関するフラックス群の消滅について 10
Tomohiko Ishida (Kyoto Univ.) Vanishing of volume flux groups

概要 We give a sufficient condition for vanishing of volume flux groups, which was first proved by Kędra–Kotschick–Morita in another way.

9月27日(土) 第VI会場

9:20~11:50

- 23 中村伊南沙 (東大数理) 曲面絡み目上の 2 次元ブレイド 10
Inasa Nakamura (Univ. of Tokyo) Two-dimensional braids over a surface link

概要 For an oriented surface link S , we can take a satellite construction called a 2-dimensional braid over S , which is a surface link in the form of a covering over S . We demonstrate that 2-dimensional braids over surface links are useful for showing the distinctness of surface links.

- 24 佐藤 進 (神戸大理) 交差交換で溶接結び目をほどく 10
 Shin Satoh (Kobe Univ.) Crossing changes unknot a welded knot

概要 We prove that any descending diagram presents a trivial welded knot. This implies that the crossing change is an unknotting operation in the category of welded knots. As an application, we prove that there is an embedding of the product of a θ -graph and a circle whose constituents present three given ribbon knotted torus in 4-space.

- 25 鎌田 聖一 (阪市大理) 曲面結び目に接着するコードと1-ハンドルについて 10
 Seiichi Kamada (Osaka City Univ.) Cords and 1-handles attached to surface-knots

概要 Boyle classified 1-handles attached to surface-knots, that are closed and connected surfaces embedded in the Euclidean 4-space, in the case that the surfaces are oriented and 1-handles are orientable with respect to the orientations of the surfaces. We classify 1-handles attached to surface-knots in the case that the surface-knots are oriented and 1-handles are non-orientable, and in the case that the surface-knots are non-orientable.

- 26 安部 哲哉 (東工大理工) Annulus twist and diffeomorphic 4-manifolds II 15
 鄭 仁大 (近畿大理工)
 Tetsuya Abe (Tokyo Tech) Annulus twist and diffeomorphic 4-manifolds II
 In Dae Jong (Kinki Univ.)

概要 We solve a strong version of Kirby's problem 3.6 (D), that is, we show that for any integer γ , there exist infinitely many mutually distinct knots such that 2-handle additions along them with framing γ yield the same 4-manifold.

- 27 土屋 政統 (学習院大理) 2つの left handed trefoil knot から得られる homotopy $K3$ surface につ
 いて 10
 Masatsuna Tsuchiya (Gakushuin Univ.) On homotopy $K3$ surface constructed by two left handed trefoil knots

概要 We define X_n to be the 4-dimensional handlebody represented by the Kirby diagram which is a simple link of two left handed trefoil knots with framing $(0, n)$, and define M_n to be the boundary of X_n . Note that M_n is a homology 3-sphere. It is known that M_{-6} bounds a contractible 4-manifold. First we show that we can get a homotopy $K3$ surface by using M_{-6} and Kirby Calculus. Next we show that a knot is not a slice knot and if n is larger than 0, M_n does not bound any contractible 4-manifolds by using adjunction inequality.

- 28 濱田 法行 (東大数理) On sections of the Matsumoto–Cadavid–Korkmaz Lefschetz fibration · 15
Noriyuki Hamada (Univ. of Tokyo) On sections of the Matsumoto–Cadavid–Korkmaz Lefschetz fibration

概要 In the study of (topological, 4-dimensional) Lefschetz fibrations, to investigate (-1) -sections of a given Lefschetz fibration is a fundamental and important problem concerning Lefschetz pencils, hence symplectic topology. On the other hand, Yukio Matsumoto originally constructed a genus-2 Lefschetz fibration with eight singular fibers, and then, Cadavid and Korkmaz independently generalized it to the higher genera. Those Lefschetz fibrations has played a great role in the development of the theory. In this talk, we show explicit monodromies of those fibrations which describe a set of disjoint (-1) -sections. The number of the (-1) -sections involved is the largest ever known, and possibly maximal.

- 29 浮田 卓也 (東工大理工) A genus zero Lefschetz fibration on the Akbulut cork ··········· 10
Takuya Ukida (Tokyo Tech) A genus zero Lefschetz fibration on the Akbulut cork

概要 We construct a genus zero positive allowable Lefschetz fibration over the disk (a genus zero PALF for short) on a cork introduced by Akbulut and describe the monodromy as a positive factorization in the mapping class group of a fiber. We also examine the monodromies of genus zero PALFs on infinitely many exotic pairs of compact Stein surfaces such that one is obtained by applying a cork twist to the other.

- 30 大場 貴裕 (東工大理工) 写像類群を用いた Stein filling の微分同相型の決定 ··········· 15
Takahiro Oba (Tokyo Tech) Diffeomorphism types of Stein fillings and mapping class groups

概要 Classification of Stein fillings of a given contact manifold has been discussed as a crucial problem in contact geometry. In particular, it is important to examine which contact manifold has a unique Stein filling. Uniqueness of Stein fillings had often been clarified by symplectic geometry before 2010, while it is sometimes proved by using combinatorics in mapping class groups after 2010. In this talk, considering Lefschetz fibrations over D^2 and mapping class groups, I will present a condition for a Stein fillable integral homology 3-sphere to have a unique Stein filling. Moreover I will mention a result of weak fillings of the same manifold.

- 31 粕谷 直彦 (東大数理) 奇数次元ユークリッド空間の接触部分多様体について ··········· 15
Naohiko Kasuya (Univ. of Tokyo) On contact submanifolds of the odd dimensional Euclidean spaces

概要 We prove that a closed co-oriented contact 3-manifold can be a contact submanifold of some contact structure on \mathbb{R}^5 , if and only if the first Chern class is trivial. We also prove that a closed co-oriented contact $(2m+1)$ -manifold (M^{2m+1}, ξ) can be a contact submanifold of the standard contact structure on \mathbb{R}^{4m+1} , if it satisfies one of the following conditions: (1) m is odd ($m \geq 3$) and $H_1(M^{2m+1}; \mathbb{Z}) = 0$, (2) m is even ($m \geq 4$) and M^{2m+1} is 2-connected, (3) $m = 2$ and M^5 is simply-connected.

- 32 堀内 智広 (中大理工) Leafwise holomorphic automorphisms of Reeb components ··········· 15
Tomohiro Horiuchi (Chuo Univ.) Leafwise holomorphic automorphisms of Reeb components

概要 The aim of this study is to classify the structures of the Reeb component as a Levi flat CR manifold, by considering the leafwise holomorphic automorphisms.

14:15~15:40

- 33 高岡 邦行 (早大教育) 球面閉曲線の LR 数について 10
 Kuniyuki Takaoka (Waseda Univ.) LR number of spherical closed curves

概要 Given an oriented spherical closed curve with n transverse crossing points, we assign a cyclic word of length $2n$ on two letters L standing left and R standing right by reading the crossing sign so that each crossing point is read once L and once R . The LR number is the number of appearance of the subword LR . In this talk, we completely determine oriented spherical closed curves whose LR number is less than or equal to three.

- 34 伊藤 昇 (早大高等研) Triple chords and strong $(1, 2)$ homotopy 10
 瀧村 祐介 (学習院中)
 Noboru Ito (Waseda Univ.) Triple chords and strong $(1, 2)$ homotopy
 Yusuke Takimura
 (Gakushuin Boy's Junior High School)

概要 A triple chord is a circle and three chords whose endpoints are on the circle and any two chords are mutually intersect. A chord diagram of a spherical curve consists of a circle and chords, each of which connects the preimages for every double point. In this talk, we consider a relationship between the number of triple chords as sub-chord diagrams and an equivalence relation called strong $(1, 2)$ homotopy for spherical curves.

- 35 伊藤 昇 (早大高等研) Strong and weak $(1, 2)$ homotopies on spherical curves and new invari-
 瀧村 祐介 (学習院中) ants 10
 Noboru Ito (Waseda Univ.) Strong and weak $(1, 2)$ homotopies on spherical curves and new invari-
 Yusuke Takimura ants
 (Gakushuin Boy's Junior High School)

概要 The second Reidemeister move can be decomposed into two types, respectively called strong or weak. This talk gives a necessary and sufficient condition that two spherical curves are equivalent by a finite sequence consisting of the first and the strong second Reidemeister moves. Similarly, we obtain a necessary and sufficient condition in the weak case. We also define an integer-valued invariant with respect to the first and the strong second Reidemeister moves.

- 36 早野 健太 (北大理) A new aspect of the Arnold invariant J^+ from a global viewpoint 15
 伊藤 昇 (早大高等研)
 Kenta Hayano (Hokkaido Univ.) A new aspect of the Arnold invariant J^+ from a global viewpoint
 Noboru Ito (Waseda Univ.)

概要 The Arnold invariant J^+ for generic plane curves counts the number of local moves called direct self-tangency perestroika in a generic regular homotopy from a standard curve to a given one. In this talk we will give infinitely many regular homotopic curves with the same J^+ which cannot be mutually related by any generic regular homotopies without direct self-tangency perestroika.

- 37 境 圭一 (信州大理) Haefliger 不変量に対する Lin-Wang 型公式 15
 Keiichi Sakai (Shinshu Univ.) Lin-Wang type formula for Haefliger invariant

概要 Long embeddings $\mathbb{R}^{4k-1} \hookrightarrow \mathbb{R}^{6k}$ are classified by Haefliger invariant. We show that Haefliger invariant behaves as it is of order two. As a byproduct we obtain an invariant for long generic immersions $\mathbb{R}^{4k-1} \looparrowright \mathbb{R}^{6k-1}$ which can be lifted to embeddings $\mathbb{R}^{4k-1} \hookrightarrow \mathbb{R}^{6k}$.

- 38 溝田 裕介 (九大数理) Lowerable ベクトル場のなす加群は有限生成か? 10
 西村 尚史 (横浜国大環境情報)

Yusuke Mizota (Kyushu Univ.) Is the module of lowerable vector fields finitely generated?

Takashi Nishimura

(Yokohama Nat. Univ.)

概要 The notion of lowerable vector field was introduced by Arnol'd for studying bifurcations of wave front singularities. Is the module of lowerable vector fields finitely generated in general? In this talk, we give a partial affirmative answer to this problem.

16:00~17:00 特別講演

石川 昌治 (東北大理) 3次元多様体の安定写像と分岐シャドウについて

Masaharu Ishikawa (Tohoku Univ.) Stable maps and branched shadows of 3-manifolds

概要 The stable maps play an important role in the study of smooth manifolds, especially used for obtaining topological information of the source manifold from the types of their singularities as seen in Morse theory. We are studying stable maps from orientable 3-manifolds to a plane. A shadow is a polyhedron equipped with a half integer on each region. The 3-manifold represented by a shadow is a kind of circle bundle over the polyhedron. F. Costantino and D. Thurston regarded the Stein factorization of a stable map as a shadow and used it to construct a shadow from a triangulation of a 3-manifold.

The aim of our study is to get information of “complexity” of 3-manifolds from stable maps. We show that the stable map complexity is equal to the branched shadow complexity for a compact orientable 3-manifold possibly with torus boundaries; the former is the minimal number of singular fibers of codimension 2 of stable maps of the 3-manifold and the latter is the minimal number of vertices of its branched shadows. These complexities estimate the volume from above and below in case the manifold is hyperbolic. We also classify hyperbolic links in the 3-sphere whose exteriors have the complexity 1 in terms of Dehn surgeries.

This is joint work with Yuya Koda.

9月28日(日) 第VI会場

9:20~12:00

- 39 鎌田直子 (名古屋市大システム自然) Twisted knot の writhe 10

Naoko Kamada (Nagoya City Univ.) The writhes of a twisted knot

概要 The odd writhe is a numerical invariant of virtual knots defined by L. Kauffman. S. Satoh and K. Taniguchi introduced a series of numerical invariants of a virtual knot, called n -writhes. They are a refinement of the odd writhe. The n -writhes are related to the index polynomial defined by Y. Im, K. Lee and Y. Lee and A. Henrich and the affine index polynomial defined by Kauffman. We discuss a generalization of them to a twisted knot.

- 40 中村拓司 (大阪電通大工) 与えられた状態数を持つ仮想結び目の集合について 10
 中西康剛 (神戸大理)
 佐藤進 (神戸大理)

Takuji Nakamura (Osaka Electro-Comm. Univ.) On the set of virtual knots with a given state number

Yasutaka Nakanishi (Kobe Univ.)
 Shin Satoh (Kobe Univ.)

概要 A *state* of a virtual knot diagram D is a union of circles obtained from D by splicing all real crossings. Let $s_n(D)$ be the number of states of D consisting of n circles. The n -state number of a virtual knot K is the minimal number of $s_n(D)$ for all possible virtual knot diagrams of K . In this talk, we consider a set of virtual knots whose n -state number is equal to i for each non-negative integer i and study the finiteness of the set.

- 41 中西康剛 (神戸大理) Delta-crossing number for knots 10
 坂本遥子 (神戸大理)
 佐藤進 (神戸大理)

Yasutaka Nakanishi (Kobe Univ.) Delta-crossing number for knots

Yoko Sakamoto (Kobe Univ.)
 Shin Satoh (Kobe Univ.)

概要 A Delta-crossing tangle is a tangle of three arcs with three crossings, which is appeared in a Delta move. A Delta-crossing diagram is a diagram which can be decomposed into Delta-crossing tangles joined by simple arcs. We prove that every knot has a Delta-crossing diagram, and then investigate the Delta-crossing number which is the minimum number of Delta-crossing tangles among all Delta-crossing diagrams of the given knot. We obtain upper and lower bounds on the number in terms of the ordinal crossing number and genus.

- 42 松崎尚作 (早大教育) 各成分が自明である絡み目の配置について 15
 Shosaku Matsuzaki (Waseda Univ.) On arrangements of component-trivial links on planes

概要 Let $L = L_1 \cup L_2 \cup \dots \cup L_n$ be a link in \mathbb{R}^3 such that L_i is a trivial link for each $1 \leq i \leq n$. Let P_1, P_2, \dots, P_n be mutually distinct flat planes in \mathbb{R}^3 such that no two of them are parallel. Then there is a link $L' = L'_1 \cup L'_2 \cup \dots \cup L'_n$ in \mathbb{R}^3 such that L is ambient isotopic to L' and $L'_i \subset P_i$ for each $1 \leq i \leq n$.

- 43 石井 敦 (筑波大数理物質) Circulatory orientations and handlebody-links 10
 Atsushi Ishii (Univ. of Tsukuba) Circulatory orientations and handlebody-links

概要 A handlebody-knot is a handlebody embedded in the 3-sphere. It is a fundamental problem to give a suitable orientation for handlebody-knot. We introduce a circulatory orientation and discuss fundamental moves for handlebody-knots.

- 44 斎藤 敏夫 (上越教育大) Essential tangle spheres of knots 10
 Toshio Saito (Joetsu Univ. of Edu.) Essential tangle spheres of knots

概要 In 1998, it was shown by Ozawa that if a knot in the 3-sphere admits an essential free 2-tangle sphere, then its essential tangle sphere is unique. We discuss if the result above could be generalized.

- 45 中川 義行 (龍谷大経済) トーラス絡み目群の有理増大度関数 10
 山下 靖 (奈良女大理)
 田村 誠 (大阪産大教養)
 Yoshiyuki Nakagawa (Ryukoku Univ.) The growth of torus link groups
 Yasushi Yamashita
 (Nara Women's Univ.)
 Makoto Tamura (Osaka Sangyo Univ.)

概要 Let G be a finitely generated group with a finite generating set S . For $g \in G$, let $l_S(g)$ be the length of the shortest word over S representing g . The growth series of G with respect to S is the series $A(t) = \sum_{n=0}^{\infty} a_n t^n$, where a_n is the number of elements of G with $l_S(g) = n$. If $A(t)$ can be expressed as a rational function of t , then G is said to have a rational growth function.

We calculate explicitly the rational growth functions of (p, q) -torus link groups for any $p, q > 1$. As an application, we show that their growth rates are Perron numbers.

- 46 小鳥居 祐香 (東大数理) Milnor の μ 不変量と HOMFLYPT 多項式の関係について 10
 Yuka Kotorii (Univ. of Tokyo) On relation between the Milnor's μ -invariant and HOMFLYPT polynomial

概要 Milnor introduced a family of invariants for ordered oriented links, called $\bar{\mu}$ -invariants. Moreover, Habegger–Lin showed that Milnor's invariants are invariants of string link, called μ -invariants. Polyak showed a relation between the $\bar{\mu}$ -invariants of length 3 sequence and Conway polynomials. We show that any μ -invariant of length $\leq k$ can be represented as a combination of HOMFLYPT polynomials if all μ -invariant of length $\leq k - 2$ vanish. This result is an extension of Polyak's result.

- 47 堤 康嘉 (大島商船高専) Negativity of the third Ohtsuki invariants of the Brieskorn–Hamm homology 3-spheres 10
 Yasuyoshi Tsutsumi
 (Oshima Nat. Coll. of Maritime Tech.) Negativity of the third Ohtsuki invariants of the Brieskorn–Hamm homology 3-spheres

概要 We calculate the third Ohtsuki invariant of every Brieskorn–Hamm manifold which is a rational homology 3-sphere. By the result, we show that the third Ohtsuki invariants of Brieskorn–Hamm homology 3-spheres are negative.

- 48 森 藤 孝 之 (慶 大 経 済)* 二橋結び目のパラボリック表現とねじれ Alexander 多項式 10
 Anh T. Tran (Ohio State Univ.)
 Takayuki Morifuji (Keio Univ.) Parabolic representations of 2-bridge knots and twisted Alexander poly-
 Anh T. Tran (Ohio State Univ.) nomials

概要 In this talk we discuss the twisted Alexander polynomial associated to parabolic representations of 2-bridge knots. We also give an affirmative answer to a conjecture of Dunfield, Friedl and Jackson for infinitely many hyperbolic knots.

- 49 北 野 晃 朗 (創 価 大 工) トーラス結び目を Dehn 手術して得られるホモロジー球面の $SL(2, \mathbb{C})$ -既約表現に対応する Reidemeister torsion について 10
 Teruaki Kitano (Soka Univ.) Reidemeister torsion of a homology 3-sphere surgeried along a torus knot for $SL(2, \mathbb{C})$ -irreducible representaions

概要 In this talk we consider a homology 3-sphere M obtained by Dehn surgery along a torus knot and a $SL(2, \mathbb{C})$ -irreducible representation of $\pi_1(M)$. Here Reidemeister torsion $\tau_\rho(M)$ can be defined as a complex number. By using Reidemeister torsion $\tau_\rho(M)$ for ρ , we define the semi-torsion polynomial $\sigma_M(t) = \prod_\rho(t - \frac{1}{2}\tau_\rho)$. In 1980's D. Johnson computed this polynomial for the homology 3-sphere surgeried along (2,3)-torus knot. In this talk we give a formula for this polynomial for a homology 3-sphere along a torus knot. It is a generalization of D. Johnson's formula.

- 50 須 志 田 隆 道 (明 大 MIMS) Shape limit in Voronoi spiral multiple tilings 15
 日 詰 明 男 (龍 谷 大 理 工)
 山 岸 義 和 (龍 谷 大 理 工)
 Takamichi Sushida (Meiji Univ.) Shape limit in Voronoi spiral multiple tilings
 Akio Hizume (Ryukoku Univ.)
 Yoshikazu Yamagishi (Ryukoku Univ.)

概要 We study Voronoi diagrams with the spiral sequence $S = \{\zeta^j = r^j e^{\sqrt{-1}j\theta} : j \in \mathbb{Z}\}$ of a covering space M_v of $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$, where $v \in \mathbb{Z} \setminus \{0\}$. They are intimately related to the phyllotaxis and continued fractions. Generically, the tiles are hexagons, while they are quadrilaterals in the degenerate case. For each multiplicity v , the set B_v of generators $\zeta \in M_v$ of quadrilateral multiple tilings is the union of branches of real algebraic curves parameterized by θ . Their union $B = \bigcup_v B_v$ is a dense subset of the unit disk. Moreover, we consider the shapes of quadrilateral tiles when $\theta/2\pi v$ is fixed and r tends to 1. If $\theta/2\pi v$ is a quadratic irrational, then the limit set of the shape parameters is a finite set. In particular, if $\theta/2\pi v$ is linearly equivalent to the golden section $\tau = \frac{1+\sqrt{5}}{2}$, then the limit shape is the square.

- 51 秋 吉 宏 尚 (阪 市 大 理) 錐特異点を 1 点持つトーラスの side parameter 10
 Hirotaka Akiyoshi (Osaka City Univ.) Side parameter for the torus with a single cone point

概要 Let M be the cone manifold obtained as the product of the torus with a single cone point of cone angle $\theta \in (0, 2\pi)$ and the interval. We introduce the side parameter for the space of cone hyperbolic structures on M whose holonomy representations are real.

無 限 可 積 分 系

9月25日(木) 第IX会場

10:00~12:00

- 1 渋川元樹(九大数理) Multivariate Meixner, Charlier and Krawtchouk polynomials 15
 Genki Shibukawa (Kyushu Univ.) Multivariate Meixner, Charlier and Krawtchouk polynomials

概要 We introduce some multivariate analogues of Meixner, Charlier and Krawtchouk polynomials, and establish their main properties, that is, duality, degenerate limits, generating functions, orthogonality relations, difference equations, recurrence formulas and determinant expressions. A particularly important and interesting result is that “the generating function of the generating function” for the Meixner polynomials coincides with the generating function of the Laguerre polynomials, which has previously not been known even for the one variable case. Actually, main properties for the multivariate Meixner, Charlier and Krawtchouk polynomials are derived from some properties of the multivariate Laguerre polynomials by using this key result.

- 2 大久保勇輔(名大多元数理) AGT予想で現れる一般化 Jack 対称関数の存在と直交性とその q 変形 .. 15
 Yusuke Ohkubo (Nagoya Univ.) Existence and orthogonality of generalized Jack symmetric functions arising from AGT conjecture and its q -deformation

概要 We investigate the existence and the orthogonality of the generalized Jack symmetric functions which play an important role in the proof of the AGT conjecture. We show their orthogonality by deforming them to the generalized Macdonald symmetric functions. In addition we talk about a possibility to describe the q -deformed AGT conjecture with help of the generalized Macdonald symmetric functions.

- 3 竹村剛一(中大理工) Multi-indexed Jacobi polynomials and Maya diagrams 15
 Kouichi Takemura (Chuo Univ.) Multi-indexed Jacobi polynomials and Maya diagrams

概要 Multi-indexed Jacobi polynomials are defined by the Wronskian of four types of eigenfunctions of a deformed Pöschl–Teller Hamiltonian. We give a correspondence between multi-indexed Jacobi polynomials and pairs of Maya diagrams, and we show that any multi-indexed Jacobi polynomial is essentially equal to some multi-indexed Jacobi polynomial of two types of eigenfunction. As an application, we show a Wronskian-type formula of some special eigenstates of the deformed Pöschl–Teller Hamiltonian.

- 4 星野 歩 (香川高専) Askey–Wilson 多項式の四重級数表示 15
 野海正俊 (神戸大理)
 白石潤一 (東大数理)
 Ayumu Hoshino Fourfold series expression for Askey–Wilson polynomial
 (Kagawa Nat. Coll. of Tech.)
 Masatoshi Noumi (Kobe Univ.)
 Junichi Shiraishi (Univ. of Tokyo)

概要 We present a fourfold series expansion representing the Askey–Wilson polynomials. To obtain the result, a sequential use is made of several summation and transformation formulas for the basic hypergeometric series, including the Verma’s q -extension of the Field and Wimp expansion, Andrews’ terminating q -analogue of Watson’s ${}_3F_2$ sum, Singh’s quadratic transformation.

- 5 星野 歩 (香川高専) 一行型 Koornwinder 多項式の明示的公式と Lassalle の予想の証明 15
 野海正俊 (神戸大理)
 白石潤一 (東大数理)
 Ayumu Hoshino An explicit formula for Koornwinder polynomial with one row diagram
 (Kagawa Nat. Coll. of Tech.) and a proof of Lassalle’s conjectures
 Masatoshi Noumi (Kobe Univ.)
 Junichi Shiraishi (Univ. of Tokyo)

概要 We present an explicit formula for the Koornwinder polynomial of type BC_n with one row diagram. When the parameters are specialized, we recover Lassalle’s formula for Macdonald polynomials of type B_n , C_n and D_n with one row diagram, thereby proving his conjectures.

- 6 齋藤洋介 (東北大理) 2重サイン関数から得られる Ding–Iohara–Miki 代数の modular double
 的構造 15
 Yosuke Saito (Tohoku Univ.) Modular double of the Ding–Iohara–Miki algebra obtained from the
 double sine function

概要 From the kernel function defined by the double sine function, we obtain two free field realizations of the Ding–Iohara–Miki algebra which commute with each other. Similar situations have been observed in studies of modular doubles of quantum groups.

14:15~15:30

- 7 澁川陽一 (北大理) Hopf algebroids associated with dynamical Yang–Baxter maps 15
 Youichi Shibukawa (Hokkaido Univ.) Hopf algebroids associated with dynamical Yang–Baxter maps
- 8 澁川陽一 (北大理) Rigid tensor categories associated with dynamical Yang–Baxter maps
 15
 Youichi Shibukawa (Hokkaido Univ.) Rigid tensor categories associated with dynamical Yang–Baxter maps

概要 We introduce rigid tensor categories associated with dynamical Yang–Baxter maps.

- 9 安東 雅訓 (稚内北星学園大)* 分割と相異分割の母関数における相似性の理解とその応用 15
 Masanori Ando (Wakhol Univ.) An understanding and its application of the similarity in the generating functions of partitions and strict partitions

概要 The generating functions of partitions and strict partitions are similar. It is trivial in the generating function. However we understand this similarity as combinatorics using simple set theory. And we generalize it to q -identity called Uchimura type.

- 10 筧 三郎 (立教大理) 変形 KdV 階層による平面曲線の運動と戸田階層 15
 梶原 健司 (九大 I M I)
 Saburo Kakei (Rikkyo Univ.) Toda hierarchy and motion of plane curves by modified KdV hierarchy
 Kenji Kajiwara (Kyushu Univ.)

概要 A relation between the Goldstein–Petrich hierarchy for plane curves and the Toda lattice hierarchy is investigated. A representation formula for plane curves is given in terms of a special class of τ -functions of the Toda lattice hierarchy.

15:45~16:45 特別講演

- 岡田 聡一 (名大多元数理) Schur-type Pfaffians and their applications to symmetric function
 Soichi Okada (Nagoya Univ.) Schur-type Pfaffians and their applications to symmetric function

概要 The evaluation of determinants or Pfaffians plays a key role in combinatorics, representation theory, infinite analysis, and so on. In this talk, we consider generalizations of Schur's Pfaffians $\text{Pf}((x_i - x_j)/(x_i + x_j))$ and $\text{Pf}((x_i - x_j)/(1 - x_i x_j))$. We can express these Schur-type Pfaffians in terms of generalized Vandermonde determinants. By using these formulae, we derive King's column-length restricted version of Littlewood's identities for summations of Schur functions, and Worley's formulae expressing near-staircase Schur's P -functions in terms of Schur functions. Also we discuss extensions and variations of them.

9月26日(金) 第IX会場

10:00~12:00

- 11 森田 健 (阪大情報) A relation between the divergent bilateral basic hypergeometric series ${}_2\psi_2(a, 0; b_1, b_2; q, x)$ and the basic hypergeometric series 15
 Takeshi Morita (Osaka Univ.) A relation between the divergent bilateral basic hypergeometric series ${}_2\psi_2(a, 0; b_1, b_2; q, x)$ and the basic hypergeometric series

概要 We give the new relation between the bilateral basic hypergeometric series around the origin and the unilateral basic hypergeometric series around the origin by the using of the q -Borel–Laplace transformations.

- 12 磯島 伸 (法政大理工) パンルヴェ III 型方程式の特殊関数解の超離散極限 15
 Shin Isojima (Hosei Univ.) Ultradiscrete limit of special function solutions of the Painlevé III equation

概要 The Painlevé III equation possesses a class of special solutions in a determinantal form with the Bessel function elements. In this talk, its ultradiscrete analog is constructed by procedure of ultradiscretization with parity variables.

- 13 竹縄知之 (東京海洋大海洋工) Schlesinger 変換と差分 Painlevé 方程式 15
 Tomoyuki Takenawa Schlesinger transformations and difference Painlevé equations
 (Tokyo Univ. of Marine Sci. and Tech.)

概要 Schlesinger transformations are algebraic transformations of a Fuchsian system that preserve its monodromy representation and act on the characteristic indices of the system by integral shifts. One of the important reasons to study such transformations is the relationship between Schlesinger transformations and discrete Painlevé equations. Based on a recent work with A. Dzhamay and H. Sakai, we derive discrete Schlesinger evolution equations describing discrete dynamical systems generated by elementary Schlesinger transformations and give their discrete Hamiltonian description w.r.t. the standard symplectic structure on the space of Fuchsian systems. As an application, we explicitly derive difference Painlevé equations of type $D_4^{(1)}$, $A_2^{(1)*}$ and $A_1^{(1)*}$ from Schlesinger transformations.

- 14 岩木耕平 (京大数理研) 完全 WKB 解析と団代数 15
 中西知樹 (名大多元数理)
 Kohei Iwaki (Kyoto Univ.) Exact WKB analysis and cluster algebras
 Tomoki Nakanishi (Nagoya Univ.)

概要 We develop the mutation theory in the exact WKB analysis using the framework of cluster algebras. As we vary the phase of Stokes curves, the Stokes graph changes the topology when a Stokes segment appears. We call this phenomenon the mutation of Stokes graphs. Such a mutation of the Stokes graph causes a Stokes phenomenon for Voros symbols. We show that the Voros symbols mutate as variables of a cluster algebra with surface realization. As an application, we obtain the identities of Stokes automorphisms associated with periods of cluster algebras.

- 15 神吉雅崇 (立教大理) 互いに素条件による離散方程式の可積分性判定 15
 時弘哲治 (東大数理)
 間瀬崇史 (東大数理)
 間田潤 (日大生産工)
- Masataka Kanki (Rikkyo Univ.) Co-primeness condition as an integrability criterion for discrete equa-
 Tetsuji Tokihiro (Univ. of Tokyo) tions
 Takafumi Mase (Univ. of Tokyo)
 Jun Mada (Nihon Univ.)

概要 We study the Laurent property, the irreducibility and co-primeness of discrete integrable and non-integrable equations. First we study a discrete integrable equation related to the Somos-4 sequence, and also a non-integrable equation as a comparison. We prove that the conditions of irreducibility and co-primeness hold only in the integrable case. Next we apply this method to the nonlinear partial difference equations, such as the discrete KdV equation and the discrete Toda equation. We conclude that co-primeness of the terms can be used as a new integrability criterion, which is a mathematical re-interpretation of the confinement of singularities in discrete equations.

- 16 加藤晃史 (東大数理) Quiver mutation loops and partition q -series 15
 寺嶋郁二 (東工大情報理工)
- Akishhi Kato (Univ. of Tokyo) Quiver mutation loops and partition q -series
 Yuji Terashima (Tokyo Tech)

概要 A quiver mutation loop is a loop in a quiver exchange graph. We define partition q -series for quiver mutation loops, and show that they enjoy various remarkable properties such as pentagon identities, and modular properties, and relation with fermionic character formulas of certain conformal field theories. The partition q -series are defined solely in terms of combinatorial data and are independent of the details of the system under study.

13:00~14:00 特別講演

- 大山陽介 (阪大情報) q -パンルヴェ方程式の古典解析
 Yousuke Ohyama (Osaka Univ.) Classical Analysis on the q -Painlevé equations

概要 The Painlevé equations are found by Paul Painlevé in 1898. He classified all of second order nonlinear ordinary differential equations without movable branch points. Sakai showed that all of initial value spaces of the Painlevé equations are rational surfaces with some condition. He classified all of such surfaces, and showed that all of elliptic, q -analogue, difference and differential Painlevé equations can be unified in his classification.

There are many researches on the q -Painlevé equations in these twenty years, such that Bäcklund transformations, special solutions and their determinant representations. But the properties of generic transcendental solutions are unrevealed. In this talk, we study analytic properties of the q -Painlevé transcendents. We review the Riemann–Hilbert–Birkhoff correspondence and study connection formula of the q -Painlevé equations.